arXiv:1207.2180v1 [math.RA] 9 Jul 2012

OPERADIC TWISTING – WITH AN APPLICATION TO DELIGNE'S CONJECTURE

VASILY DOLGUSHEV AND THOMAS WILLWACHER

To the memory of Jean-Louis Loday

Abstract. We study categorial properties of the operadic twisting functor Tw. In particular, we show that Tw is a comonad. Coalgebras of this comonad are operads for which a natural notion of twisting by Maurer-Cartan elements exists. We give a large class of examples, including the classical cases of the Lie, associative and Gerstenhaber operads, and their infinity-counterparts Lie_∞, As_∞, Ger_∞. We also show that Tw is well behaved with respect to the homotopy theory of operads. As an application we show that every solution of Deligne's conjecture is homotopic to a solution that is compatible with twisting.

Contents

1. Introduction 2
1.1. Structure of the paper 4
Memorial note 5
Acknowledgements 5
2. Preliminaries 5
2.1. Trees 7
2.2. Operads, pseudo-operads, and their dual versions 10
2.3. Cobar construction 16
2.4. Convolution Lie algebra and its properties 17
2.5. What if $Q(n)$ is finite dimensional for all n? 18
2.6. Intrinsic derivations of an operad 20
3. Twisting of operads 21
3.1. The action of $\mathcal{L}_{\mathcal{O}}$ on $\widetilde{\mathrm{Tw}}\,\mathcal{O}$ 25
3.2. Algebras over $\mathrm{Tw}\,\mathcal{O}$ 27
3.3. Example: The dg operad $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$ 33
3.4. A useful modification $\mathrm{Tw}^{\oplus}\mathcal{O}$ 36
3.5. More general version of twisting 37
4. Categorial properties of twisting 37
4.1. Tw as a comonad 37
4.2. Coalgebras over the comonad Tw 42
4.3. Ger_∞ is canonically a Tw-coalgebra 43

5. Homotopy theoretic properties of the twisting procedure 50
5.1. The distributive law and the functor Tw 51
6. Twisting $\mathcal{O}$-algebra structures by Maurer-Cartan elements 62
7. The operad of brace trees BT 65
7.1. The operad structure on BT 66
8. The operad $\mathrm{Tw}\,\mathsf{BT}$ 68
8.1. The differential on $\mathrm{Tw}\,\mathsf{BT}$ 71
9. The operad of braces Br à la Kontsevich-Soibelman 72
9.1. A simpler description of the differential on Br 73
9.2. The inclusion $\mathsf{Br} \hookrightarrow \mathrm{Tw}\,\mathsf{BT}$ is a quasi-isomorphism 74
9.3. The dg operad Br is a homotopy fixed point of Tw 76
9.4. Br is a Tw-coalgebra 76
10. Every solution of Deligne's conjecture is homotopic to a one that is compatible with twisting 76
10.1. The proof of Proposition 10.2 79
10.2. Proof of Corollary 1.2 82
Appendix A. The operad Ger 83
Appendix B. Action of the operad $\mathrm{Tw}\,\mathsf{BT}$ on the Hochschild cochain complex of an A_∞-algebra 83
Appendix C. A direct computation of the cohomology of Br 87
C.1. Computing $H(V_\circ(n), \delta_0)$ 88
C.2. Computing $H(V_\bullet(n), \delta_0)$ 89
C.3. A technical claim 93
C.4. Analyzing the map (C.2) 97
Appendix D. Br is a homotopy Tw fixed point – combinatorial argument 99
Appendix E. A Lemma on a filtered complex 100
References 100

1. Introduction

The idea of twisting algebraic objects by Maurer-Cartan elements has been a folklore for very long time and the original nudge for this idea probably belongs to D. Quillen [24].

For example, let $\mathfrak{g}$ be a differential graded Lie algebra and let $m \in \mathfrak{g}$ be a Maurer-Cartan element, i.e., a degree 1 element satisfying the Maurer-Cartan equation

$$dm + \frac{1}{2}\,[m, m] = 0.$$

Then the twisted Lie algebra $\mathfrak{g}^m$ is the graded space $\mathfrak{g}$ with differential $d + [m, \cdot]$ and the same bracket $[\cdot, \cdot]$. In a similar manner the notion of Maurer-Cartan element and twisting can be defined for associative and Gerstenhaber algebras, and their homotopy versions, i.e., Lie_∞, As_∞ and Ger_∞ algebras (see, e. g. [4], [6], [7, Section 2.4], and [29]).

In this paper we formalize the notion of twisting by Maurer-Cartan elements, using the operadic twisting functor Tw introduced in [28]. Concretely, we show in Theorem 4.7 that Tw is naturally equipped with the structure of a comonad and this comonad has the following properties[1]:

- The twisting procedure by Maurer-Cartan elements can be defined for algebras over an operad which is, in turn, a coalgebra for the comonad Tw (see Section 6).
- A map $\mathcal{O} \to \mathcal{O}'$ of operads induces a functor from the category of $\mathcal{O}'$-algebras to the category of $\mathcal{O}$-algebras. If $\mathcal{O} \to \mathcal{O}'$ gives rise to a map of Tw-coalgebras, then this functor is compatible with the operation of twisting by Maurer-Cartan elements (see Theorem 6.1).
- The functor Tw preserves quasi-isomorphisms between operads (see Theorem 5.1).

We apply the developed machinery to study solutions of (our version of) the Deligne conjecture. By this we mean, abusing notation, a quasi-isomorphism

$$F\colon \mathsf{Ger}_\infty \to \mathsf{Br} \tag{1.1}$$

from the operad Ger_∞ governing homotopy Gerstenhaber algebras to the braces operad Br.

Both operads Ger_∞ and Br are Tw-coalgebras, i.e., their algebras can be twisted by Maurer-Cartan elements. However, in general, a quasi-isomorphism (1.1) does not need to be a morphism of Tw-coalgebras.

So, using our machinery, we established that every homotopy class of a quasi-isomorphism (1.1) has at least one representative which respects the structures of Tw-coalgebras on Ger_∞ and Br. More precisely,

Theorem 1.1. *Suppose that $F : \mathsf{Ger}_\infty \to \mathsf{Br}$ is a quasi-isomorphism of dg operads. Then there exists a quasi-isomorphism of dg operads $F' : \mathsf{Ger}_\infty \to \mathsf{Br}$ such that*

- *F' is a homomorphism of Tw-coalgebras, and*
- *F' is homotopy equivalent to F.*

Furthermore, the morphism F' is given by explicit formula (10.1)*.*

Let us recall that, since the operad Br acts on the Hochschild cochain complex $C^\bullet(A)$ of an A_∞-algebra A, a quasi-morphism (1.1) gives a Ger_∞-structure on $C^\bullet(A)$. Theorem 1.1 has an interesting consequence related to this Ger_∞-structure which we will describe now.

Let A be an A_∞-algebra and let F_A denote the composition

$$F_A = \mathfrak{a}_A \circ F : \mathsf{Ger}_\infty \to \mathsf{End}_{C^\bullet(A)}, \tag{1.2}$$

where $\mathsf{End}_{C^\bullet(A)}$ is the endomorphism operad of $C^\bullet(A)$ and

$$\mathfrak{a}_A : \mathsf{Br} \to \mathsf{End}_{C^\bullet(A)}$$

---

[1] Here we postpone the precise statements to later sections, since they require terminology introduced in the definition of Tw.

is the operad morphism coming from the action of Br on $C^\bullet(A)$. Let us assume, in addition, that the dg Lie algebra $C^\bullet(A)$ carries a complete descending filtration[2] $\mathcal{F}_\bullet C^\bullet(A)$ and let α be a Maurer-Cartan element in $\mathcal{F}_1 C^\bullet(A)$.

Such a Maurer-Cartan element gives us a new A_∞-structure on A and we denote this new A_∞-algebra by A^α. In addition, the Maurer-Cartan element α can be used to twist the dg Lie algebra structure on $C^\bullet(A)$ and form a new cochain complex

$$C^\bullet(A)^\alpha .$$

It is easy to see the cochain complexes $C^\bullet(A)^\alpha$ coincides with the Hochschild cochain complex $C^\bullet(A^\alpha)$ of the new A_∞-algebra A^α. Furthermore, there are two natural ways of obtaining a Ger_∞-structure on $C^\bullet(A^\alpha) = C^\bullet(A)^\alpha$. First, we may compose the operad morphism

$$\mathfrak{a}_{A^\alpha} : \mathsf{Br} \to C^\bullet(A^\alpha)$$

with $F : \mathsf{Ger}_\infty \to \mathsf{Br}$ and, second, we may twist F_A (1.2) by α and obtain

$$F_A^\alpha : \mathsf{Ger}_\infty \to C^\bullet(A)^\alpha .$$

We claim that

Corollary 1.2. *For every solution F (1.1) of Deligne's conjecture, the maps F_{A^α} and F_A^α are homotopic. Furthermore, if the morphism F is compatible with the coalgebra structures on Ger_∞ and Br over* Tw*, then F_{A^α} and F_A^α coincide.*

For the proof of this corollary we refer the reader to Section 10.2.

1.1. Structure of the paper.

In Section 2, we fix notation and recall some facts about operads and their dual versions. In this section, we also review the convolution Lie algebra and its properties. In Section 3, we introduce the functor Tw [28] and give an alternative description of the category of algebras over an operad $\mathrm{Tw}\,\mathcal{O}$. In this section, we also describe in detail the dg operad $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$.

Section 4 is devoted to categorical properties of the functor Tw . We show that the functor Tw forms a comonad on the under-category $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads*. At the end of this section, we prove that the dg operad Ger_∞ governing homotopy Gerstenhaber algebras has the canonical coalgebra structure over the comonad Tw .

In Section 5, we describe homotopy theoretic properties of the functor Tw and introduce the notion of homotopy fixed point for Tw (see Definition 5.3). In this section, we also describe a large class of dg operads which are simultaneously Tw-coalgebras and homotopy fixed points for Tw. This class includes the classical operads Lie, Ger and the operad governing Poisson algebras. At the end of this section, we show that the operads As and As_∞ are Tw-coalgebras and are homotopy fixed points for Tw .

---

[2] Such a filtration may come from a formal deformation parameter ε which enters the game when we extend the base field to the ring $\mathbb{K}[[\varepsilon]]$.

In Section 6, we consider a dg operad $\mathcal{O}$ which carries a coalgebra structure over Tw . We introduce the notion of twisting by Maurer-Cartan elements for $\mathcal{O}$-algebras and describe categorial and homotopy theoretical properties of this twisting procedure.

Sections 7, 8 and 9 treat the brace operad Br. In this section, we show how to recover the dg operad Br (essentially) as the twisted version of a simpler operad BT. The proofs of Theorem 1.1 and Corollary 1.2 are given in Section 10.

At the end of the paper we have several appendices. In Appendix A, we recall the operad Ger governing Gerstenhaber algebras. In Appendix B, we describe in detail the action of the operads BT, $\mathrm{Tw}\,\mathsf{BT}$ and Br on the Hochschild cochain complex $C^\bullet(A)$ of an A_∞-algebra A. In Appendix C, we give a "topology independent" proof of the fact that

$$H^\bullet(\mathsf{Br}) \cong \mathsf{Ger}\,.$$

This statement is not used directly in our text. However, we believe that our proof of this fact has an independent interest. In Appendix D, we sketch a different proof of the fact that the brace operad Br is a homotopy fixed point for Tw. Finally, in Appendix E, we prove a technical lemma about filtered complexes.

Memorial note. With great sadness, we learned that Jean-Louis Loday died as a consequence of an accident on June 6, 2012. He is widely known for his contributions to algebraic K-theory, for his research linking cyclic homology, K-theory, and combinatorics. His foundational work on operads has and will have a lasting impact on the development of algebra and topology. We devote our paper to the memory of this great mathematician.

Acknowledgements. We would like to thank Martin Markl, Ezra Getzler, Dmitry Tamarkin and Bruno Vallette for helpful discussions. V.D. acknowledges the NSF grants DMS-0856196, DMS-1161867 and the grant FASI RF 14.740.11.0347. V.D.'s NSF grant DMS-0856196 was supported by American Recovery and Reinvestment Act (ARRA) and it could not be transferred to Temple University from the UC Riverside by ARRA regulations. V.D. would like to thank Charles Greer, Tim LeFort, and Charles Louis from the UCR Office of Research for allowing V.D. to save this grant by creating a subaward from the UC Riverside to Temple University. V.D. also thanks John Baez for serving formally as the PI on DMS-0856196 starting July 1, 2010.

2. PRELIMINARIES

The underlying field $\mathbb{K}$ has characteristic zero. For a set X we denote by $\mathbb{K}\langle X\rangle$ the $\mathbb{K}$-vector space of finite linear combinations of elements in X .

The notation S_n is reserved for the symmetric group on n letters and $\mathrm{Sh}_{p_1,\dots,p_k}$ denotes the subset of $(p_1,\dots,p_k)$-shuffles in S_n, i.e. $\mathrm{Sh}_{p_1,\dots,p_k}$ consists of elements $\sigma\in S_n$, $n=p_1+p_2+\cdots+p_k$ such that

$$\sigma(1) < \sigma(2) < \cdots < \sigma(p_1),$$

$$\sigma(p_1+1) < \sigma(p_1+2) < \cdots < \sigma(p_1+p_2),$$

$$\dots$$

$$\sigma(n-p_k+1) < \sigma(n-p_k+2) < \cdots < \sigma(n)\,.$$

The underlying symmetric monoidal category $\mathfrak{C}$ is either the category $\mathsf{grVect}_{\mathbb{K}}$ of $\mathbb{Z}$-graded $\mathbb{K}$-vector spaces or the category $\mathsf{Ch}_{\mathbb{K}}$ of (possibly) unbounded cochain complexes of $\mathbb{K}$-vector spaces. We frequently use the ubiquitous combination "dg" (differential graded) to refer to algebraic objects in $\mathsf{Ch}_{\mathbb{K}}$. For a homogeneous vector v in a cochain complex, $|v|$ denotes the degree of v . Furthermore, we denote by $\mathbf{s}$ (resp. $\mathbf{s}^{-1}$) the operation of suspension (resp. desuspension) on $\mathsf{grVect}_{\mathbb{K}}$ or $\mathsf{Ch}_{\mathbb{K}}$.

The notation Lie (resp. As, Com, Ger) is reserved for the operad governing Lie algebras (resp. associative algebras without unit, commutative (and associative) algebras without unit, Gerstenhaber algebras without unit). Dually, the notation coLie (resp. coAs, coCom) is reserved for the cooperad governing Lie coalgebras (resp. coassociative coalgebras without counit, cocommutative (and coassociative) coalgebras without counit).

For an operad $\mathcal{O}$ (resp. a cooperad $\mathcal{C}$) and a cochain complex V, the notation $\mathcal{O}(V)$ (resp. $\mathcal{C}(V)$) is reserved for the free $\mathcal{O}$-algebra (resp. cofree $\mathcal{C}$-coalgebra). Namely,

$$\mathcal{O}(V) := \bigoplus_{n\ge 0} \Big(\mathcal{O}(n) \otimes V^{\otimes\, n} \Big)_{S_n}\,, \tag{2.1}$$

$$\mathcal{C}(V) := \bigoplus_{n\ge 0} \Big(\mathcal{C}(n) \otimes V^{\otimes\, n} \Big)^{S_n}\,. \tag{2.2}$$

For an operad $\mathcal{O}$, we denote by $\mathrm{Alg}_{\mathcal{O}}$ the category of $\mathcal{O}$-algebras.

Let $a_1, a_2, \dots, a_n$ be degree 0 dummy variables and $\mathcal{O}$ be a dg operad. It is clear that $\mathcal{O}(n)$ is naturally identified with the subspace of the free $\mathcal{O}$-algebra

$$\mathcal{O}\big(\mathbb{K}\langle a_1, a_2, \dots, a_n\rangle\big)$$

spanned by $\mathcal{O}$-monomials in which each variable from the set $\{a_1, a_2, \dots, a_n\}$ appears exactly once. We often use this identification in our paper.

For an augmented operad $\mathcal{O}$ (resp. a coaugmented cooperad $\mathcal{C}$) the notation $\mathcal{O}_\circ$ (resp. $\mathcal{C}_\circ$) is reserved for the kernel (resp. the cokernel) of the augmentation (resp. coaugmentation).

For an operad (resp. a cooperad) P in $\mathsf{Ch}_{\mathbb{K}}$ we denote by ΛP the operad (resp. the cooperad) with the spaces of n-ary operations:

$$\Lambda P(n) = \mathbf{s}^{1-n} P(n) \otimes \mathrm{sgn}_n\,, \tag{2.3}$$

where sgn_n denotes the sign representation of S_n . For example, an algebra over $\Lambda\mathsf{Lie}$ is a graded vector space V equipped with the binary operation:

$$\{\,,\,\} : V \otimes V \to V$$

of degree -1 satisfying the identities:

$$\{v_1, v_2\} = (-1)^{|v_1||v_2|}\{v_2, v_1\}$$

$$\{\{v_1, v_2\}, v_3\} + (-1)^{|v_1|(|v_2|+|v_3|)}\{\{v_2, v_3\}, v_1\} + (-1)^{|v_3|(|v_1|+|v_2|)}\{\{v_3, v_1\}, v_2\} = 0\,,$$

where v_1, v_2, v_3 are homogeneous vectors in V.

$\mathsf{Ger}^\vee$ denotes the Koszul dual cooperad [11], [14], [15] for Ger. It is known [16] that

$$\mathsf{Ger}^\vee = \Lambda^2 \mathsf{Ger}^*\,, \tag{2.4}$$

where Ger^* is obtained from the operad Ger by taking the linear dual. In other words, algebras over the linear dual $(\mathsf{Ger}^\vee)^*$ are very much like Gerstenhaber algebras except that the bracket carries degree 1 and the multiplication carries degree 2.

For a groupoid $\mathcal{G}$ the notation $\pi_0(\mathcal{G})$ is reserved for the set of its isomorphism classes.

2.1. **Trees.** In this paper all trees are rooted and the root vertex has always valency 1. (Such trees are sometimes called *planted*). The remaining vertices of valency 1 are called *leaves*. A vertex is called *internal* if it is neither a root nor a leaf. We always orient trees in the direction towards the root. Thus every internal vertex has at least 1 incoming edge and exactly 1 outgoing edge. An edge adjacent to a leaf is called *external*. We allow a degenerate tree, that is a tree with exactly two vertices (the root vertex and a leaf) connected by a single edge.

Let us recall that for every planar tree $\mathbf{t}$ the set $V(\mathbf{t})$ of its vertices is equipped with a natural total order. To define this total order on $V(\mathbf{t})$ we introduce the function

$$\mathcal{N} : V(\mathbf{t}) \to V(\mathbf{t})\,. \tag{2.5}$$

To a non-root vertex v the function $\mathcal{N}$ assigns the next vertex along the (unique) path connecting v to the root vertex. Furthermore, $\mathcal{N}$ sends the root vertex to the root vertex.

Let v_1, v_2 be two distinct vertices of $\mathbf{t}$. If v_1 lies on the path which connects v_2 to the root vertex then we declare that

$$v_1 < v_2\,.$$

Similarly, if v_2 lies on the path which connects v_1 to the root vertex then we declare that

$$v_2 < v_1\,.$$

If neither of the above options realize then there exist numbers k_1 and k_2 such that

$$\mathcal{N}^{k_1}(v_1) = \mathcal{N}^{k_2}(v_2) \tag{2.6}$$

and

$$\mathcal{N}^{k_1-1}(v_1) \neq \mathcal{N}^{k_2-1}(v_2)\,.$$

Since the tree $\mathbf{t}$ is planar the set of $\mathcal{N}^{-1}(\mathcal{N}^{k_1}(v_1))$ is equipped with a total order. Furthermore, since both vertices $\mathcal{N}^{k_1-1}(v_1)$ and $\mathcal{N}^{k_2-1}(v_2)$ belong to the set $\mathcal{N}^{-1}(\mathcal{N}^{k_1}(v_1))$, we may compare them with respect to this order.

We declare that, if $\mathcal{N}^{k_1-1}(v_1) < \mathcal{N}^{k_2-1}(v_2)$, then

$$v_1 < v_2\,.$$

Otherwise we set $v_2 < v_1$.

It is not hard to see that the resulting relation $<$ on $V(\mathbf{t})$ is indeed a total order. Keeping this order in mind, we can say things like "the first vertex", "the second vertex", and "the i-th vertex" of a planar tree $\mathbf{t}$. In fact, the first vertex of a tree is always its root vertex.

We have an obvious bijection between the set of edges $E(\mathbf{t})$ of a tree $\mathbf{t}$ and the subset of vertices:

$$V(\mathbf{t}) \setminus \{\text{root vertex}\}\,. \tag{2.7}$$

This bijection assigns to a vertex v in (2.7) its outgoing edge. Thus the canonical total order on the set (2.7) gives us a natural total order on the set of edges $E(\mathbf{t})$. For our purposes we also extend the total orders on the sets $V(\mathbf{t}) \setminus \{\text{root vertex}\}$ and $E(\mathbf{t})$ to the disjoint union

$$V(\mathbf{t}) \setminus \{\text{root vertex}\} \sqcup E(\mathbf{t}) \tag{2.8}$$

by declaring that a vertex is bigger than its outgoing edge. For example, the root edge is the minimal element in the set (2.8).

2.1.1. *Groupoid of labelled planar trees.* Let n be a non-negative integer. An n-labeled planar tree $\mathbf{t}$ is a planar tree equipped with an injective map

$$\mathfrak{l} : \{1, 2, \dots, n\} \to L(\mathbf{t}) \tag{2.9}$$

from the set $\{1, 2, \dots, n\}$ to the set $L(\mathbf{t})$ of leaves of $\mathbf{t}$. Although the set $L(\mathbf{t})$ has a natural total order we do not require that the map (2.9) is monotonous.

The set $L(\mathbf{t})$ of leaves of an n-labelled planar tree $\mathbf{t}$ splits into the disjoint union of the image $\mathfrak{l}(\{1, 2, \dots, n\})$ and its complement. We call leaves in the image of $\mathfrak{l}$ *labelled.*

A vertex x of an n-labelled planar tree $\mathbf{t}$ is called *nodal* if it is neither a root vertex, nor a labelled leaf. We denote by $V_{\text{nod}}(\mathbf{t})$ the set of all nodal vertices of $\mathbf{t}$. Keeping in mind the canonical total order on the set of all vertices of $\mathbf{t}$, we can say things like "the first nodal vertex", "the second nodal vertex", and "the i-th nodal vertex".

Example 2.1. *An example of a 4-labelled planar tree is depicted on figure 2.1. On figures we use*



FIG. 2.1. A 4-labelled planar tree

small white circles for nodal vertices and small black circles for labelled leaves and the root vertex.

For our purposes we need to upgrade the set of n-labelled planar trees to the groupoid $\mathsf{Tree}(n)$. Objects of $\mathsf{Tree}(n)$ are n-labelled planar trees and morphisms are non-planar isomorphisms of the corresponding (non-planar) trees compatible with labelling. The groupoid $\mathsf{Tree}(n)$ is equipped with an obvious left action of the symmetric group S_n.

There is exactly one labelled planar tree with no nodal vertices at all. This is the degenerate tree with a single edge connecting the root vertex with a single leaf labelled by 1. For our purposes it is convenient to discard this tree. However, we would still like to keep the degenerate tree depicted on figure 2.7.

The notation $\mathsf{Tree}_2(n)$ is reserved for the full sub-groupoid of $\mathsf{Tree}(n)$ whose objects are n-labelled planar trees with exactly 2 nodal vertices. It is not hard to see that every object in $\mathsf{Tree}_2(n)$ has at most $n+1$ leaves. Furthermore, isomorphism classes of $\mathsf{Tree}_2(n)$ are in bijection with the union

$$\bigsqcup_{0\le p\le n} \mathrm{Sh}_{p,n-p} \tag{2.10}$$

where $\mathrm{Sh}_{p,n-p}$ denotes the set of $(p,n-p)$-shuffles in S_n. The bijection assigns to a $(p,n-p)$-shuffle τ the isomorphism class of the planar tree depicted on figure 2.2.



FIG. 2.2. Here τ is a $(p,n-p)$-shuffle

2.1.2. *Insertions of trees.* Let $\widetilde{\mathbf{t}}$ be an n-labelled planar tree. If the i-th nodal vertex of $\widetilde{\mathbf{t}}$ has m_i incoming edges then for every m_i-labelled planar tree $\mathbf{t}$ we can define the insertion $\bullet_i$ of the tree $\mathbf{t}$ into the i-th nodal vertex of $\widetilde{\mathbf{t}}$. The resulting planar tree $\widetilde{\mathbf{t}}\bullet_i\mathbf{t}$ is also n-labelled.

If $m_i=0$ then $\widetilde{\mathbf{t}}\bullet_i\mathbf{t}$ is obtained via identifying the root edge of $\mathbf{t}$ with edge originating at the i-th nodal vertex.

If $m_i>0$ then the tree $\widetilde{\mathbf{t}}\bullet_i\mathbf{t}$ is built following these steps:

- First, we denote by $E_i(\widetilde{\mathbf{t}})$ the set of edges terminating at the i-th nodal vertex of $\widetilde{\mathbf{t}}$. Since $\widetilde{\mathbf{t}}$ is planar, the set $E_i(\widetilde{\mathbf{t}})$ comes with a total order;
- second, we erase the i-th nodal vertex of $\widetilde{\mathbf{t}}$;
- third, we identify the root edge of $\mathbf{t}$ with the edge of $\widetilde{\mathbf{t}}$ which originates at the i-th nodal vertex;
- finally, we identify the edges of $\mathbf{t}$ adjacent to labelled leaves with the edges in the set $E_i(\widetilde{\mathbf{t}})$ following this rule: the external edge with label j gets identified with the j-th edge in the set $E_i(\widetilde{\mathbf{t}})$. In doing this, we keep the same planar structure on $\mathbf{t}$, so, in general, branches of $\widetilde{\mathbf{t}}$ move around.

Example 2.2. *Let* $\widetilde{\mathbf{t}}$ *be the* 4*-labelled planar tree depicted on figure 2.1 and* $\mathbf{t}$ *be the* 3*-labelled planar tree depicted on figure 2.3. Then the insertion* $\widetilde{\mathbf{t}} \bullet_1 \mathbf{t}$ *of* $\mathbf{t}$ *into the first nodal vertex of* $\widetilde{\mathbf{t}}$ *is shown on figure 2.4.*



FIG. 2.3. A 3-labelled planar tree $\mathbf{t}$



FIG. 2.4. The 4-labelled planar tree $\widetilde{\mathbf{t}} \bullet_1 \mathbf{t}$

2.2. **Operads, pseudo-operads, and their dual versions.**

2.2.1. *Collections.* By a *collection* we mean the sequence $\{P(n)\}_{n\ge 0}$ of objects of the underlying symmetric monoidal category $\mathfrak{C}$ such that for each n, the object $P(n)$ is equipped with a left action of the symmetric group S_n .

Given a collection P we form covariant functors for $n \ge 0$

$$\underline{P}_n : \mathsf{Tree}(n) \to \mathfrak{C}\,.$$

To an n-labelled planar tree $\mathbf{t}$ the functor $\underline{P}_n$ assigns the object

$$\underline{P}_n(\mathbf{t}) = \bigotimes_{x\in V_{\mathrm{nod}}(\mathbf{t})} P(m(x))\,, \tag{2.11}$$

where $V_{\mathrm{nod}}(\mathbf{t})$ is the set of all nodal vertices of $\mathbf{t}$, the notation $m(x)$ is reserved for the number of edges terminating at the vertex x, and the order of the factors in the right hand side of the equation agrees with the natural order on the set $V_{\mathrm{nod}}(\mathbf{t})$.

To define the functor $\underline{P}_n$ on the level of morphisms we use the actions of the symmetric groups and the structure of the symmetric monoidal category $\mathfrak{C}$ in the obvious way.

Example 2.3. *Let* $\mathbf{t}_1$ *(resp.* $\mathbf{t}_2$*) be a* 3*-labelled planar tree depicted on figure 2.5 (resp. figure 2.6). There is a unique morphism* λ *from* $\mathbf{t}_1$ *to* $\mathbf{t}_2$ *in* $\mathsf{Tree}(3)$ *. For these trees we have*



FIG. 2.5. A 2-labelled planar tree $\mathbf{t}_1$



FIG. 2.6. A 2-labelled planar tree $\mathbf{t}_2$

$$\underline{P}_2(\mathbf{t}_1) = P(2) \otimes P(3) \otimes P(0) \otimes P(0)\,,$$

$$\underline{P}_2(\mathbf{t}_2) = P(2) \otimes P(0) \otimes P(3) \otimes P(0)\,,$$

and the morphism

$$\underline{P}_2(\lambda) : P(2) \otimes P(3) \otimes P(0) \otimes P(0) \to P(2) \otimes P(0) \otimes P(3) \otimes P(0)$$

is the composition

$$\underline{P}_2 = (1 \otimes \beta) \circ (\sigma_{12} \otimes \sigma_{13} \otimes 1 \otimes 1),$$

where σ_{12} *(resp.* σ_{13}*) is the corresponding transposition in* S_2 *(resp. in* S_3*) and* β *is the braiding*

$$\beta : \big(P(3) \otimes P(0)\big) \otimes P(0) \to P(0) \otimes \big(P(3) \otimes P(0)\big)\,.$$

2.2.2. *Plethysm of collections.* Let P and Q be collections in $\mathfrak{C}$. It is known [14] that the formula

$$P \odot Q(n) = \bigoplus_{r \geq 0} P(r) \otimes_{S_r} \Big(\bigoplus_{k_1 + \cdots + k_r = n} \mathsf{Ind}^{S_n}_{S_{k_1} \times \cdots \times S_{k_r}} Q(k_1) \otimes \cdots \otimes Q(k_r) \Big) \tag{2.12}$$

defines a monoidal structure on the category of collections. The product (2.12) is known as plethysm. The plethysm can be used to give a very concise definition of an operad. However, we will use it in a slightly different way.

2.2.3. *Pseudo-operads and operads.* We now recall that a pseudo-operad is a collection $\{P(n)\}_{n\geq 0}$ equipped with multiplication maps

$$\mu_{\mathbf{t}} : \underline{P}_n(\mathbf{t}) \to P(n) \tag{2.13}$$

for all n-labelled trees $\mathbf{t}$ and for all $n \geq 0$. These multiplications should satisfy the axioms which we list below.

First, we require that the standard corolla $\mathbf{q}_n$ (see figures 2.7 and 2.8) acts via identity



FIG. 2.7. The corolla $\mathbf{q}_0$



FIG. 2.8. The corolla $\mathbf{q}_n$ for $n \geq 1$

$$\mu_{\mathbf{q}_n} = \mathrm{id}_{P(n)}\,. \tag{2.14}$$

Second, we require that the operation $\mu_{\mathbf{t}}$ is S_n-equivariant

$$\mu_{\sigma(\mathbf{t})} = \sigma \circ \mu_{\mathbf{t}}\,. \tag{2.15}$$

Third, for every morphism $\lambda : \mathbf{t} \to \mathbf{t}'$ in $\mathsf{Tree}(n)$ we have

$$\mu_{\mathbf{t}'} \circ \underline{P}_n(\lambda) = \mu_{\mathbf{t}}\,. \tag{2.16}$$

Finally, we need to formulate the associativity axiom for multiplications (2.13). For this purpose we consider the following quadruple $(\widetilde{\mathbf{t}}, i, m_i, \mathbf{t})$ where $\widetilde{\mathbf{t}}$ is an n-labelled planar tree with k nodal vertices, $1 \leq i \leq k$, m_i is the number of edges terminating at the i-th nodal vertex of $\widetilde{\mathbf{t}}$, and $\mathbf{t}$ is an m_i-labelled planar tree.

The associativity axiom states that for each such quadruple $(\widetilde{\mathbf{t}}, i, m_i, \mathbf{t})$ we have

$$\mu_{\widetilde{\mathbf{t}}} \circ (\mathrm{id} \otimes \cdots \otimes \mathrm{id} \otimes \underbrace{\mu_{\mathbf{t}}}_{i\text{-th spot}} \otimes \mathrm{id} \otimes \cdots \otimes \mathrm{id}) \circ \beta_{\widetilde{\mathbf{t}}, i, m_i, \mathbf{t}} = \mu_{\widetilde{\mathbf{t}} \bullet_i \mathbf{t}} \tag{2.17}$$

where $\widetilde{\mathbf{t}} \bullet_i \mathbf{t}$ is the n-labelled planar tree obtained by inserting $\mathbf{t}$ into the i-th vertex of $\widetilde{\mathbf{t}}$ and $\beta_{\widetilde{\mathbf{t}}, i, m_i, \mathbf{t}}$ is the isomorphism in $\mathfrak{C}$ which is responsible for "putting tensor factors in the correct order".

Given integers $n \geq 1$, $k \geq 0$, $1 \leq i \leq n$ and a permutation $\sigma \in S_{n+k-1}$ we can form the $(n+k-1)$-labelled planar tree $\mathbf{t}^{n,k,i}_{\sigma}$ shown on figure 2.9.



FIG. 2.9. The $(n+k-1)$-labelled planar tree $\mathbf{t}^{n,k,i}_{\sigma}$

It is often convenient to use a different notation for the multiplication map $\mu_{\mathbf{t}^{n,k,i}_{\sigma}}$ corresponding to the tree $\mathbf{t}^{n,k,i}_{\sigma}$. More precisely, for a vector $v \in P(n)$ and $w \in P(k)$ of a pseudo-operad P we will use this notation

$$v\big(\sigma(1), \dots \sigma(i-1), w\big(\sigma(i), \dots, \sigma(i+k-1)\big), \sigma(i+k), \dots, \sigma(n+k-1)\big) := \mu_{\mathbf{t}^{n,k,i}_{\sigma}}(v, w)\,. \tag{2.18}$$

Recall that, for $\sigma = \mathrm{id} \in S_{n+k-1}$, the multiplications

$$\mu_{\mathbf{t}^{n,k,i}_{\mathrm{id}}} : P(n) \otimes P(k) \to P(n+k-1)$$

are called *the elementary insertion* and often denoted by $\circ_i$. Namely, for $v \in P(n)$ and $w \in P(k)$ we have[3]

$$v \circ_i w := \mu_{\mathbf{t}^{n,k,i}_{\mathrm{id}}}(v, w)\,. \tag{2.19}$$

It is not hard to see that a pseudo-operad structure on a collection P is uniquely determined by elementary insertions (2.19). All the remaining multiplications (2.13) can be expressed in terms of (2.19) using the axioms of a pseudo-operad.

Furthermore, to verify that (2.19) define a pseudo-operad structure[4] on a collection P one has to check that the operations $\circ_i$ are compatible with the differential and to show that for all homogeneous vectors a, b, c in $P(n_a), P(n_b)$, and $P(n_c)$, respectively and for all $1 \leq i \leq n_a$,

---

[3] Numbers n and k are suppressed from the notation.

[4] See the equivalent definition of a pseudo-operad in terms of the elementary insertions (2.19) in [20, Definition 17].

$1 \le j \le n_a + n_b - 1$, $1 \le p \le n_b - 1$, $1 \le k \le n_a - 1$

$$(a \circ_i b) \circ_j c = \begin{cases} (-1)^{|b||c|} (a \circ_j c) \circ_{i+n_c-1} b & \text{if } j < i\,, \\ a \circ_i (b \circ_{j-i+1} c) & \text{if } i \le j \le i + n_b - 1\,, \\ (-1)^{|b||c|} (a \circ_{j-n_b+1} c) \circ_i b & \text{if } j \ge i + n_b\,, \end{cases} \tag{2.20}$$

$$a \circ_i (\sigma_{p\,(p+1)} b) = \sigma_{(p+i-1)\,(p+i)} (a \circ_i b)\,, \tag{2.21}$$

$$(\sigma_{k\,(k+1)} a) \circ_i b = \begin{cases} \sigma_{k\,(k+1)} (a \circ_i b) & \text{if } k+1 < i \\ \varsigma_{i-1,\, i+n_b-1} (a \circ_{i-1} b) & \text{if } k+1 = i \\ \varsigma^{-1}_{i,\, i+n_b} (a \circ_{i+1} b) & \text{if } k+1 = i \\ \sigma_{(k+n_b-1)\,(k+n_b)} (a \circ_i b) & \text{if } k > i\,, \end{cases} \tag{2.22}$$

where $\sigma_{i_1\, i_2}$ denotes the transposition (i_1, i_2) and ς_{i_1, i_2} is the cycle $(i_1, i_1+1, \dots, i_2-1, i_2)$.

Let us now recall that an operad is a pseudo-operad P with unit, that is a map

$$\mathbf{u} : \mathbb{K} \to P(1) \tag{2.23}$$

for which the compositions

$$\begin{aligned} P(n) \cong P(n) \otimes \mathbb{K} &\overset{\mathrm{id}\otimes\mathbf{u}}{\longrightarrow} P(n) \otimes P(1) \xrightarrow{\circ_i} P(n) \\ P(n) \cong \mathbb{K} \otimes P(n) &\overset{\mathbf{u}\otimes\mathrm{id}}{\longrightarrow} P(1) \otimes P(n) \xrightarrow{\circ_1} P(n) \end{aligned} \tag{2.24}$$

coincide with the identity map on $P(n)$.

Let us remark that the map (2.23) is uniquely determined by the image $\mathbf{u}(1) \in P(1)$ of the generator $1 \in \mathbb{K}$. We call $\mathbf{u}(1) \in P(1)$ the identity element. By abuse of notation, we denote by $\mathbf{u}$ both the map (2.23) and the identity element in $P(1)$.

Morphisms of pseudo-operads and operads are defined in the obvious way.

2.2.4. *Pseudo-cooperads and cooperads.* Reversing the arrows in the definition of a pseudo-operad we get the definition of a pseudo-cooperad. More precisely, a pseudo-cooperad is a collection Q equipped with comultiplication maps

$$\Delta_{\mathbf{t}} : Q(n) \to \underline{Q}_n(\mathbf{t})\,, \tag{2.25}$$

which satisfy a similar list of axioms.

Just as for pseudo-operads, we have

$$\Delta_{\mathbf{q}_n} = \mathrm{id}_{Q(n)}\,, \tag{2.26}$$

where $\mathbf{q}_n$ is the standard corolla shown on figures 2.7 and 2.8.

We also require that the operations (2.25) are S_n-equivariant

$$\Delta_{\sigma(\mathbf{t})} \circ \sigma = \Delta_{\mathbf{t}} \,. \tag{2.27}$$

For every morphism $\lambda : \mathbf{t} \to \mathbf{t}'$ in $\mathsf{Tree}(n)$ we have

$$\Delta_{\mathbf{t}'} = \underline{Q}_n(\lambda) \circ \Delta_{\mathbf{t}} \,. \tag{2.28}$$

Finally, to formulate the coassociativity axiom for (2.25), we consider the following quadruple $(\widetilde{\mathbf{t}}, i, m_i, \mathbf{t})$ where $\widetilde{\mathbf{t}}$ is an n-labelled planar tree with k nodal vertices, $1 \le i \le k$, m_i is the number of edges terminating at the i-th nodal vertex of $\widetilde{\mathbf{t}}$, and $\mathbf{t}$ is an m_i-labelled planar tree.

The coassociativity axiom states that for each such quadruple $(\widetilde{\mathbf{t}}, i, m_i, \mathbf{t})$ we have

$$(\mathrm{id} \otimes \cdots \otimes \mathrm{id} \otimes \underbrace{\Delta_{\mathbf{t}}}_{i\text{-th spot}} \otimes \mathrm{id} \otimes \cdots \otimes \mathrm{id}) \circ \Delta_{\widetilde{\mathbf{t}}} = \beta_{\widetilde{\mathbf{t}}, i, m_i, \mathbf{t}} \circ \Delta_{\widetilde{\mathbf{t}} \bullet_i \mathbf{t}} \tag{2.29}$$

where $\widetilde{\mathbf{t}} \bullet_i \mathbf{t}$ is the n-labelled planar tree obtained by inserting $\mathbf{t}$ into the i-th nodal vertex of $\widetilde{\mathbf{t}}$ and $\beta_{\widetilde{\mathbf{t}}, i, m_i, \mathbf{t}}$ is the isomorphism in $\mathfrak{C}$ which is responsible for "putting tensor factors in the correct order".

Just as for pseudo-operads, a pseudo-cooperad structure on a collection Q is uniquely determined by the comultiplications:

$$\Delta_i := \Delta_{\mathbf{t}^{n,k,i}_{\mathrm{id}}} : Q(n+k-1) \to Q(n) \otimes Q(k) \,, \tag{2.30}$$

where $\{\mathbf{t}^{n,k,i}_{\sigma}\}_{\sigma \in S_{n+k-1}}$ is the family of labelled planar trees defined on figure 2.9.

The comultiplications (2.30) are called *elementary co-insertions.*

We now recall that a cooperad is a pseudo-cooperad Q with counit, that is a map

$$\mathbf{u}^* : Q(1) \to \mathbb{K} \tag{2.31}$$

for which the compositions

$$\begin{aligned} Q(n) &\xrightarrow{\Delta_i} Q(n) \otimes Q(1) \xrightarrow{\mathrm{id} \otimes \mathbf{u}^*} Q(n) \otimes \mathbb{K} \cong Q(n) \\ Q(n) &\xrightarrow{\Delta_1} Q(1) \otimes Q(n) \xrightarrow{\mathbf{u}^* \otimes \mathrm{id}} \mathbb{K} \otimes Q(n) \cong Q(n) \end{aligned} \tag{2.32}$$

coincide with the identity map on $Q(n)$.

Morphisms of pseudo-cooperads and cooperads are defined in the obvious way.

2.2.5. *Augmented operads and coaugmented cooperads.* Let us observe that the collection $*$ with

$$*\,(n) = \begin{cases} \mathbb{K} & \text{if } n = 1 \\ \mathbf{0} & \text{otherwise} \end{cases} \tag{2.33}$$

is equipped with the unique structure of an operad and a unique structure of a cooperad. In fact, $*$ is the initial object in the category of operads and $*$ is the terminal object in the category of cooperads.

We say that an operad $\mathcal{O}$ is augmented if $\mathcal{O}$ is equipped with an operad morphism

$$\varepsilon : \mathcal{O} \to * \,. \tag{2.34}$$

Similarly, a cooperad $\mathcal{C}$ is called coaugmented if $\mathcal{C}$ is equipped with a cooperad morphism

$$\varepsilon' : * \to \mathcal{C} \,. \tag{2.35}$$

It is not hard to see that a kernel of an augmentation (2.34) (resp. cokernel of a coaugmentation (2.35)) is naturally a pseudo-operad (resp. pseudo-cooperad).

This assignment can be easily upgraded to a functor. Furthermore, according to M. Markl [20, Proposition 21], this functor gives us an equivalence between the category of augmented operads and the category of pseudo-operads. Dually, we have an equivalence between the category of coaugmented cooperads and the category of pseudo-cooperads.

In this paper we denote by $\mathcal{O}_\circ$ (resp. $\mathcal{C}_\circ$) the kernel (resp. the cokernel) of the augmentation (resp. the coaugmentation) of an augmented operad $\mathcal{O}$ (resp. a coaugmented cooperad $\mathcal{C}$).

2.2.6. *Free operad.* Let Q be a collection in $\mathsf{Ch}_{\mathbb{K}}$. Following [3, Section 5.8] the spaces $\big\{\Psi\mathbb{OP}(Q)(n)\big\}_{n\ge 0}$ of the free pseudo-operad $\Psi\mathbb{OP}(Q)$ generated by the collection Q are

$$\Psi\mathbb{OP}(Q)(n) = \mathsf{colim}\, \underline{Q}_n \tag{2.36}$$

where $\underline{Q}_n$ is the functor from the groupoid $\mathsf{Tree}(n)$ to $\mathsf{Ch}_{\mathbb{K}}$ defined in Subsection 2.2.1.

The pseudo-operad structure on $\Psi\mathbb{OP}(Q)$ is defined in the obvious way using grafting of trees and the free operad $\mathbb{OP}(Q)$ generated by Q is obtained from $\Psi\mathbb{OP}(Q)$ by formally adjoining the unit.

Unfolding (2.36) we see that $\Psi\mathbb{OP}(Q)(n)$ is the quotient of the direct sum

$$\bigoplus_{\mathbf{t} \in \mathsf{Tree}(n)} \underline{Q}_n(\mathbf{t}) \tag{2.37}$$

by the subspace spanned by vectors of the form

$$(\mathbf{t}, X) - (\mathbf{t}', \underline{Q}_n(\lambda)(X))$$

where $\lambda : \mathbf{t} \to \mathbf{t}'$ is a morphism in $\mathsf{Tree}(n)$ and $X \in \underline{Q}_n(\mathbf{t})$.

Remark 2.4. *In view of the above description, generators $X \in Q(n)$ of the free operad $\mathbb{OP}(Q)$ can be also written in the form*

$$(\mathbf{q}_n, X)\,,$$

where $\mathbf{q}_n$ is the standard n-corolla (see figures 2.7, 2.8). We hope that the identification $X = (\mathbf{q}_n, X)$ does not lead to a confusion.

2.3. **Cobar construction.** The cobar construction Cobar [11], [14], [15] is a functor from the category of coaugmented dg cooperads to the category of augmented dg operads. It is used to construct free resolutions for operads.

Let $\mathcal{C}$ be a coaugmented cooperad in $\mathsf{Ch}_{\mathbb{K}}$. As an operad in the category $\mathsf{grVect}_{\mathbb{K}}$, $\mathrm{Cobar}(\mathcal{C})$ is freely generated by the collection $\mathbf{s}\,\mathcal{C}_{\circ}$

$$\mathrm{Cobar}(\mathcal{C}) = \mathbb{OP}(\mathbf{s}\,\mathcal{C}_{\circ})\,, \tag{2.38}$$

where $\mathcal{C}_{\circ}$ denotes the cokernel of the coaugmentation.

To define the differential $\partial^{\mathrm{Cobar}}$ on $\mathrm{Cobar}(\mathcal{C})$, we recall that $\mathsf{Tree}_2(n)$ is the full subcategory of $\mathsf{Tree}(n)$ which consists of n-labeled planar trees with exactly 2 nodal vertices and $\pi_0(\mathsf{Tree}_2(n))$ is the set of isomorphism classes in the groupoid $\mathsf{Tree}_2(n)$.

Since the operad $\mathrm{Cobar}(\mathcal{C})$ is freely generated by the collection $\mathbf{s}\,\mathcal{C}_{\circ}$, it suffices to define the differential $\partial^{\mathrm{Cobar}}$ on generators.

We have

$$\partial^{\mathrm{Cobar}} = \partial' + \partial''\,, \tag{2.39}$$

with

$$\partial'(X) = -\mathbf{s}\,\partial_{\mathcal{C}}\mathbf{s}^{-1}\,X\,, \tag{2.40}$$

and

$$\partial''(X) = -\sum_{z \in \pi_0(\mathsf{Tree}_2(n))} (\mathbf{s}\otimes\mathbf{s})\big(\mathbf{t}_z; \Delta_{\mathbf{t}_z}(\mathbf{s}^{-1}\,X)\big)\,, \tag{2.41}$$

where $X \in \mathbf{s}\,\mathcal{C}_{\circ}(n)$, $\mathbf{t}_z$ is any representative of the isomorphism class $z \in \pi_0(\mathsf{Tree}_2(n))$, and $\partial_{\mathcal{C}}$ is the differential on $\mathcal{C}$. The axioms of a pseudo-cooperad imply that the right hand side of (2.41) does not depend on the choice of representatives $\mathbf{t}_z$.

The identity

$$\partial' \circ \partial' = 0$$

readily follows from $(\partial_{\mathcal{C}})^2 = 0$. Identity

$$\partial' \circ \partial'' + \partial'' \circ \partial' = 0 \tag{2.42}$$

follows from the compatibility of the differential $\partial_{\mathcal{C}}$ with the cooperad structure and the identity

$$\partial'' \circ \partial'' = 0 \tag{2.43}$$

follows from the coassociativity axiom (2.29).

2.4. **Convolution Lie algebra and its properties.** In this section, we recall the convolution Lie algebra and describe its properties. All the statements given here were proved in the more general setting of properads by S. Merkulov and B. Vallette [23].

Let P (resp. Q) be a pseudo-operad (resp. pseudo-cooperad) in the category $\mathsf{Ch}_{\mathbb{K}}$ of unbounded cochain complexes of $\mathbb{K}$-vector spaces.

We consider the following cochain complex

$$\mathrm{Conv}(Q,P) = \prod_{n\geq 0} \mathsf{Hom}_{S_n}(Q(n), P(n))\,. \tag{2.44}$$

with the binary operation $\bullet$ defined by the formula

$$f \bullet g(X) = \sum_{z\in\pi_0\big(\mathsf{Tree}_2(n)\big)} \mu_{\mathbf{t}_z}(f\otimes g\ \Delta_{\mathbf{t}_z}(X))\,, \tag{2.45}$$

$$f,g\in \mathrm{Conv}(Q,P), \qquad X\in Q(n)\,,$$

where $\mathbf{t}_z$ is any representative of the isomorphism class $z\in\pi_0\big(\mathsf{Tree}_2(n)\big)$. The axioms of a pseudo-operad (resp. pseudo-cooperad) imply that the right hand side of (2.45) does not depend on the choice of representatives $\mathbf{t}_z$.

It follows directly from the definition that the operation $\bullet$ is compatible with the differential on $\mathrm{Conv}(Q,P)$ coming from Q and P . Furthermore, we claim that

Proposition 2.5 (Proposition 4.1, [8])**.** *The bracket*

$$[f,g] = \big(f\bullet g - (-1)^{|f|\,|g|} g\bullet f\big)$$

satisfies the Jacobi identity. □

In other words, $\mathrm{Conv}(Q,P)$ is a dg Lie algebra. We refer to $\mathrm{Conv}(Q,P)$ as the *convolution Lie algebra* of the pair (Q,P) .

Let $\mathcal{C}$ be a coaugmented dg cooperad with $\mathcal{C}_\circ$ being the cokernel of the coaugmentation.

We observe that for every a dg operad $\mathcal{O}$ the morphism (of dg operads)

$$F : \mathrm{Cobar}(\mathcal{C}) \to \mathcal{O} \tag{2.46}$$

is uniquely determined by its restriction

$$F\Big|_{\mathbf{s}\,\mathcal{C}_\circ} \ :\ \mathbf{s}\,\mathcal{C}_\circ \to \mathcal{O}\,. \tag{2.47}$$

Thus, to every morphism (2.46) we assign a degree 1 element

$$\alpha_F \in \mathrm{Conv}(\mathcal{C}_\circ, \mathcal{O}) \tag{2.48}$$

such that

$$F\Big|_{\mathbf{s}\,\mathcal{C}_\circ} = \alpha_F \circ \mathbf{s}^{-1}\,.$$

The construction of the dg Lie algebra $\mathrm{Conv}(\mathcal{C},\mathcal{O})$ is partially justified by the following statement:

Theorem 2.6 (Proposition 5.2, [8]). *Let $\mathcal{C}$ be a coaugmented dg cooperad with $\mathcal{C}_\circ$ being the cokernel of the coaugmentation. Then, for every dg operad $\mathcal{O}$, the above assignment*

$$F \ \mapsto\ \alpha_F$$

is a bijection between the set of maps (of dg operads) (2.46) *and Maurer-Cartan elements of the dg Lie algebra*

$$\mathrm{Conv}(\mathcal{C}_\circ, \mathcal{O})\,. \ \square$$

2.5. **What if $Q(n)$ is finite dimensional for all n?** Let us assume that the pseudo-cooperad Q satisfies the property

Property 2.7. *For each n the graded vector space $Q(n)$ is finite dimensional.*

Due to this property we have

$$\mathrm{Conv}(Q, P) \cong \prod_{n\ge 0} \big(P(n)\otimes Q^*(n)\big)^{S_n}\,. \tag{2.49}$$

where $Q^*(n)$ denotes the linear dual of the vector space $Q(n)$.

The collection $Q^* := \{Q^*(n)\}_{n\ge 0}$ is naturally a pseudo-operad and we can express the pre-Lie structure (2.45) in terms of elementary insertions on P and Q^*. Namely, given two vectors

$$X = \sum_{n\ge 0} v_n \otimes w_n, \qquad X' = \sum_{n\ge 0} v'_n \otimes w'_n$$

in

$$\prod_{n\ge 0} \big(P(n)\otimes Q^*(n)\big)^{S_n}\,,$$

we have

$$X \bullet X' = \sum_{n\ge 1, m\ge 0} (-1)^{|v'_m||w_n|} \sum_{\sigma\in \mathrm{Sh}_{m,n-1}} \sigma(v_n \circ_1 v'_m) \otimes \sigma(w_n \circ_1 w'_m)\,. \tag{2.50}$$

Let us now observe that the formula

$$\mathrm{Av}(v\otimes w) = \sum_{\sigma\in S_n} \sigma(v)\otimes\sigma(w) \tag{2.51}$$

defines a $\mathbb{K}$-linear map

$$\mathrm{Av} : P(n)\otimes Q^*(n) \to \big(P(n)\otimes Q^*(n)\big)^{S_n}\,.$$

Furthermore, since $\mathbb{K}$ has characteristic zero, Av induces an isomorphism (which we denote by the same letter):

$$\mathrm{Av} : \big(P(n)\otimes Q^*(n)\big)_{S_n} \xrightarrow{\cong} \big(P(n)\otimes Q^*(n)\big)^{S_n} \tag{2.52}$$

from the space

$$\big(P(n)\otimes Q^*(n)\big)_{S_n} \tag{2.53}$$

of S_n-coinvariants to the space $\big(P(n)\otimes Q^*(n)\big)^{S_n}$ of invariants.

We have the following proposition.

Proposition 2.8. *Let v, v', w, w' be homogeneous vectors in $P(n)$, $P(m)$, $Q(n)$, and $Q'(m)$, respectively. Then the formula*

$$(v \otimes w) \bullet' (v' \otimes w') = \sum_{i=1}^{n} (-1)^{|v'||w|} v \circ_i v' \otimes w \circ_i w' \tag{2.54}$$

defines a binary operation on

$$\bigoplus_{n \geq 0} \big(P(n) \otimes Q^*(n) \big)_{S_n} .$$

This operation extends in the obvious way to the infinite sum

$$\prod_{n \geq 0} \big(P(n) \otimes Q^*(n) \big)_{S_n} \tag{2.55}$$

and

$$\mathrm{Av}\big((v \otimes w) \bullet' (v' \otimes w') \big) = \mathrm{Av}(v \otimes w) \bullet \mathrm{Av}(v' \otimes w') . \tag{2.56}$$

Proof. Since the map (2.52) is an isomorphism of graded vector spaces, it suffices to prove that equation (2.56) holds.

This equation can be verified by a straightforward computation using the obvious identity[5]

$$\sum_{\sigma \in S_n} \sigma(v) = \sum_{\sigma' \in S_{\{2,3,\dots,n\}}} \sum_{i=1}^{n} \sigma' \circ (1, 2, \dots, i)(v)$$

and properties of the elementary insertions. □

Proposition 2.8 implies that the map (2.52) establishes an isomorphism between the pre-Lie algebra (2.49) with the pre-Lie product (2.50) and the pre-Lie algebra (2.55) with the pre-Lie product given by equation (2.54).

2.5.1. *Example: The dg operad $\Lambda\mathsf{Lie}_\infty$.* Let us recall from [15], that the dg operad $\Lambda\mathsf{Lie}_\infty$ governing homotopy $\Lambda\mathsf{Lie}$-algebras is

$$\Lambda\mathsf{Lie}_\infty = \mathrm{Cobar}(\Lambda^2 \mathsf{coCom}) , \tag{2.57}$$

where coCom is the cooperad governing cocommutative coalgebras without counit.

Let us denote by $1^{\mathfrak{c}}_n$ the canonical generator

$$1^{\mathfrak{c}}_n := \mathbf{s}^{2-2n}\, 1 \in \mathbf{s}^{2-2n}\, \mathbb{K} \cong \Lambda^2 \mathsf{coCom}(n) .$$

Using the identification between isomorphism classes of objects in $\mathsf{Tree}_2(n)$ and the set (2.10) of shuffles we get

$$\partial^{\mathrm{Cobar}}(\mathbf{s}\, 1^{\mathfrak{c}}_n) = - \sum_{p=2}^{n-1} \sum_{\sigma \in \mathrm{Sh}_{p,n-p}} \sigma\big((\mathbf{s}\, 1^{\mathfrak{c}}_{n-p+1}) \circ_1 (\mathbf{s}\, 1^{\mathfrak{c}}_p) \big) , \qquad n \geq 2 . \tag{2.58}$$

Let us also recall from [8, Section 5.2] that the canonical quasi-isomorphism of dg operads

$$U_{\Lambda\mathsf{Lie}} : \Lambda\mathsf{Lie}_\infty = \mathrm{Cobar}(\Lambda^2 \mathsf{coCom}) \ \to\ \Lambda\mathsf{Lie} \tag{2.59}$$

---

[5] Here $S_{\{2,3,\dots,n\}}$ is the subgroup of elements $\sigma \in S_n$ satisfying the condition $\sigma(1) = 1$.

corresponds to the Maurer-Cartan element

$$\alpha_{\Lambda\mathsf{Lie}} = \{a_1, a_2\} \otimes b_1 b_2 \in \mathrm{Conv}(\Lambda^2 \mathsf{coCom}_{\circ}, \Lambda\mathsf{Lie}) \cong \prod_{n \ge 2} \Big(\Lambda\mathsf{Lie}(n) \otimes \Lambda^{-2}\mathsf{Com}(n) \Big)^{S_n} ,$$

where $\{a_1, a_2\}$ (resp. $b_1 b_2$) denotes the natural generator of $\Lambda\mathsf{Lie}(2)$ (resp. $\Lambda^{-2}\mathsf{Com}(2)$).

In other words,

$$U_{\Lambda\mathsf{Lie}}(\mathbf{s} 1^{\mathfrak{c}}_n) = \begin{cases} \{a_1, a_2\} & \text{if } n = 2\,, \\ 0 & \text{otherwise}\,. \end{cases}$$

2.6. **Intrinsic derivations of an operad.** Let P be a dg operad. Then the operation $\circ_1$ equips $P(1)$ with a structure of an associative algebra. We consider $P(1)$ as a Lie algebra with the Lie bracket being the commutator.

We claim that

Proposition 2.9. *The formula*

$$\delta_b(a) = b \circ_1 a - (-1)^{|a||b|} \sum_{i=1}^{n} a \circ_i b \qquad (2.60)$$

with

$$b \in P(1), \qquad and \qquad a \in P(n)$$

defines an operadic derivation of P for every $b \in P(1)$.

Proof. Let $a_1 \in P(n_1)$ and $a_2 \in P(n_2)$. Then for every $b \in P(1)$ and $1 \le j \le n_1$ we have

$$\begin{aligned}
\delta_b(a_1 \circ_j a_2) &= b \circ_1 (a_1 \circ_j a_2) - (-1)^{(|a_1|+|a_2|)|b|} \sum_{i=1}^{n_1+n_2-1} (a_1 \circ_j a_2) \circ_i b \\
&= (b \circ_1 a_1) \circ_j a_2 - (-1)^{|a_1||b|} \sum_{\substack{1 \le i \le n_1 \\ i \ne j}} (a_1 \circ_i b) \circ_j a_2 - (-1)^{(|a_1|+|a_2|)|b|} \sum_{i=1}^{n_2} a_1 \circ_j (a_2 \circ_i b) \\
&= (b \circ_1 a_1) \circ_j a_2 - (-1)^{|a_1||b|} \sum_{i=1}^{n_1} (a_1 \circ_i b) \circ_j a_2 \\
&\qquad + (-1)^{|a_1||b|} (a_1 \circ_j b) \circ_j a_2 - (-1)^{(|a_1|+|a_2|)|b|} \sum_{i=1}^{n_2} a_1 \circ_j (a_2 \circ_i b) \\
&= (b \circ_1 a_1) \circ_j a_2 - (-1)^{|a_1||b|} \sum_{i=1}^{n_1} (a_1 \circ_i b) \circ_j a_2 \\
&\qquad + (-1)^{|a_1||b|} a_1 \circ_j (b \circ_1 a_2) - (-1)^{(|a_1|+|a_2|)|b|} \sum_{i=1}^{n_2} a_1 \circ_j (a_2 \circ_i b) \\
&= \delta_b(a_1) \circ_j a_2 + (-1)^{|a_1||b|} a_1 \circ_j \delta_b(a_2)\,.
\end{aligned}$$

Hence δ_b is indeed an operadic derivation of P .

The identity

$$[\delta_{b_1}, \delta_{b_2}] = \delta_{[b_1, b_2]}$$

follows from a similar direct computation. □

An operadic derivation of the form (2.60) is called *intrinsic*.

3. Twisting of operads

Let $\mathcal{O}$ be a dg operad equipped with a map

$$\widehat{\varphi} : \Lambda\mathsf{Lie}_\infty \to \mathcal{O} \tag{3.1}$$

and V be an algebra over $\mathcal{O}$.

Using the map $\widehat{\varphi}$, we equip V with a $\Lambda\mathsf{Lie}_\infty$-structure. If we assume, in addition, that V is equipped with a complete descending filtration

$$V \supset \mathcal{F}_1 V \supset \mathcal{F}_2 V \supset \mathcal{F}_3 V \supset \dots\,, \qquad V = \lim_k V \Big/ \mathcal{F}_k V \tag{3.2}$$

and the $\mathcal{O}$-algebra structure on V is compatible with this filtration then we may define *Maurer-Cartan elements* of V as degree 2 elements $\alpha \in \mathcal{F}_1 V$ satisfying the equation

$$\partial(\alpha) + \sum_{n \ge 2} \frac{1}{n!} \{\alpha, \alpha, \dots, \alpha\}_n = 0 \tag{3.3}$$

where ∂ is the differential on V and $\{\cdot, \cdot, \dots, \cdot\}_n$ are the operations of the $\Lambda\mathsf{Lie}_\infty$-structure on V. Given such a Maurer-Cartan element α we can twist the differential on V and insert α into various $\mathcal{O}$-operations on V. This way we get a new algebra structure on V. It turns out that this new algebra structure is governed by a dg operad $\mathrm{Tw}\,\mathcal{O}$ which is built from the pair $(\mathcal{O}, \widehat{\varphi})$. This section is devoted to the construction of $\mathrm{Tw}\,\mathcal{O}$.

First, we recall that $\Lambda\mathsf{Lie}_\infty = \mathrm{Cobar}(\Lambda^2\mathsf{coCom})$. Hence, due to Theorem 2.6, the morphism (3.1) is determined by a Maurer-Cartan element

$$\varphi \in \mathrm{Conv}(\Lambda^2\mathsf{coCom}_\circ, \mathcal{O})\,. \tag{3.4}$$

The n-th space of $\Lambda^2\mathsf{coCom}_\circ$ is the trivial S_n-modules placed in degree $2-2n$:

$$\Lambda^2\mathsf{coCom}(n) = \mathbf{s}^{2-2n}\mathbb{K}\,.$$

So we have

$$\mathrm{Conv}(\Lambda^2\mathsf{coCom}_\circ, \mathcal{O}) = \prod_{n \ge 2} \mathsf{Hom}_{S_n}(\mathbf{s}^{2-2n}\mathbb{K}, \mathcal{O}(n)) = \prod_{n \ge 2} \mathbf{s}^{2n-2} \big(\mathcal{O}(n)\big)^{S_n}\,.$$

For our purposes we will need to extend the dg Lie algebra $\mathrm{Conv}(\Lambda^2\mathsf{coCom}_\circ, \mathcal{O})$ to

$$\mathcal{L}_{\mathcal{O}} = \mathrm{Conv}(\Lambda^2\mathsf{coCom}, \mathcal{O}) = \prod_{n \ge 1} \mathsf{Hom}_{S_n}(\mathbf{s}^{2-2n}\mathbb{K}, \mathcal{O}(n))\,. \tag{3.5}$$

It is clear that

$$\mathcal{L}_{\mathcal{O}} = \prod_{n \ge 1} \mathbf{s}^{2n-2} \big(\mathcal{O}(n)\big)^{S_n}\,.$$

For $n, r \ge 1$ we realize the group S_r as the following subgroup of S_{r+n}

$$S_r \cong \big\{ \sigma \in S_{r+n} \mid \sigma(i) = i\,, \quad \forall\, i > r \big\}\,. \tag{3.6}$$

In other words, the group S_r may be viewed as subgroup of S_{r+n} permuting only the first r letters. We set S_0 to be the trivial group. Using this embedding of S_r into S_{n+r} we introduce the following collection $(n \geq 0)$

$$\widetilde{\mathrm{Tw}}\,\mathcal{O}(n) = \prod_{r \geq 0} \mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathcal{O}(r+n))\,. \tag{3.7}$$

It is clear that

$$\widetilde{\mathrm{Tw}}\,\mathcal{O}(n) = \prod_{r \geq 0} \mathbf{s}^{2r}\big(\mathcal{O}(r+n)\big)^{S_r}\,. \tag{3.8}$$

To define an operad structure on (3.7) we denote by 1_r the generator

$$1_r := \mathbf{s}^{-2r}\,1 \in \mathbf{s}^{-2r}\mathbb{K}\,.$$

Then the identity element $\mathbf{u}$ in $\widetilde{\mathrm{Tw}}\,\mathcal{O}(1)$ is given by

$$\mathbf{u}(1_r) = \begin{cases} \mathbf{u}_{\mathcal{O}} & \text{if } r = 0\,, \\ 0 & \text{otherwise}\,, \end{cases} \tag{3.9}$$

where $\mathbf{u}_{\mathcal{O}} \in \mathcal{O}(1)$ is the identity element for the operad $\mathcal{O}$. Next, for $f \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(n)$ and $g \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(m)$, we define the i-th elementary insertion $\circ_i$ $1 \leq i \leq n$ by the formula

$$f \circ_i g(1_r) = \sum_{p=0}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \mu_{\mathbf{t}_{\sigma,i}}\big(f(1_p) \otimes g(1_{r-p})\big)\,. \tag{3.10}$$

where the tree $\mathbf{t}_{\sigma,i}$ is depicted on figure 3.1.



FIG. 3.1. Here σ is a $(p, r-p)$-shuffle

Sometimes it is convenient to use a different but equivalent definition of the i-th elementary insertion $\circ_i$ for $\widetilde{\mathrm{Tw}}\,\mathcal{O}$. This definition is given by the formula[6]

$$f \circ_i g(1_r) = \tag{3.11}$$

$$\sum_{p=0}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \sigma\big(f_p(1,\dots,p,r{+}1,\dots,r{+}i{-}1, g_{r-p}(p{+}1,\dots,r,r{+}i,\dots,r{+}i{+}m{-}1), r{+}i{+}m,\dots,r{+}n{+}m{-}1)\big)\,,$$

where $f \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(n)$, $g \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(m)$, $f_p = f(1_p) \in \big(\mathcal{O}(p+n)\big)^{S_p}$ and $g_q = g(1_q) \in \big(\mathcal{O}(q+m)\big)^{S_q}$.

---

[6] Here we use the notation for operadic multiplications (2.18) from Section 2.2.3.

To see that the element $f \circ_i g(1_r) \in \mathcal{O}(r+n+m-1)$ is S_r-invariant one simply needs to use the fact that every element $\tau \in S_r$ can be uniquely presented as the composition $\tau_{sh} \circ \tau_{p,r-p}$, where τ_{sh} is a $(p, r-p)$-shuffle and $\tau_{p,r-p} \in S_p \times S_{r-p}$.

Let $f \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(n)$, $g \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(m)$, $h \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(k)$, $1 \le i \le n$, and $1 \le j \le m$. To check the identity

$$f \circ_i (g \circ_j h) = (f \circ_i g) \circ_{j+i-1} h \tag{3.12}$$

we observe that

$$\begin{aligned} f \circ_i (g \circ_j h)(1_r) &= \sum_{p=0}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \mu_{\mathbf{t}_{\sigma,i}} \big(f(1_p) \otimes (g \circ_j h)(1_{r-p}) \big) \\ &= \sum_{p_1+p_2+p_3=r} \; \sum_{\sigma \in \mathrm{Sh}_{p_1,p_2+p_3}} \; \sum_{\sigma' \in \mathrm{Sh}_{p_2,p_3}} \mu_{\mathbf{t}_{\sigma,i}} \circ (1 \otimes \mu_{\mathbf{t}_{\sigma',j}}) \big(f(1_{p_1}) \otimes g(1_{p_2}) \otimes h(1_{p_3}) \big) \\ &= \sum_{p_1+p_2+p_3=r} \; \sum_{\tau \in \mathrm{Sh}_{p_1,p_2,p_3}} \mu_{\mathbf{t}_{\tau,i,j}} \big(f(1_{p_1}) \otimes g(1_{p_2}) \otimes h(1_{p_3}) \big) \,, \end{aligned}$$

where the tree $\mathbf{t}_{\tau,i,j}$ is depicted on figure 3.2. Similar calculations show that



FIG. 3.2. Here τ is a (p_1, p_2, p_3)-shuffle and $r = p_1 + p_2 + p_3$

$$(f \circ_i g) \circ_{j+i-1} h = \sum_{p_1+p_2+p_3=r} \; \sum_{\tau \in \mathrm{Sh}_{p_1,p_2,p_3}} \mu_{\mathbf{t}_{\tau,i,j}} \big(f(1_{p_1}) \otimes g(1_{p_2}) \otimes h(1_{p_3}) \big) \,,$$

with $\mathbf{t}_{\tau,i,j}$ being the tree depicted on figure 3.2.

We leave the verification of the remaining axioms of the operad structure for the reader.

Our next goal is to define an auxiliary action of the dg Lie algebra $\mathcal{L}_{\mathcal{O}}$ on the operad $\widetilde{\mathrm{Tw}}\,\mathcal{O}$. For a vector $f \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(n)$ the action of $v \in \mathcal{L}_{\mathcal{O}}$ (3.5) on f is defined by the formula

$$v \cdot f(1_r) = -(-1)^{|v||f|} \sum_{p=1}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \mu_{\mathbf{t}_{\sigma,p,r-p}} \big(f(1_{r-p+1}) \otimes v(1^{\mathfrak{c}}_p) \big)\,, \tag{3.13}$$

where $1^{\mathfrak{c}}_p$ is the generator $\mathbf{s}^{2-2p}\,1 \in \Lambda^2\mathsf{coCom}(p) \cong \mathbf{s}^{2-2p}\,\mathbb{K}$ and the tree $\mathbf{t}_{\sigma,p,r-p}$ is depicted on figure 3.3.



FIG. 3.3. Here σ is a $(p, r-p)$-shuffle

We claim that

Proposition 3.1. *Formula* (3.13) *defines an action of the dg Lie algebra $\mathcal{L}_{\mathcal{O}}$ (3.5) on the operad $\widetilde{\mathrm{Tw}}\,\mathcal{O}$.*

Proof. A simple degree bookkeeping shows that the degree of $v \cdot f$ is $|v| + |f|$.

Then we need to check that for two homogeneous vectors $v, w \in \mathcal{L}_{\mathcal{O}}$ we have

$$[v,w] \cdot f(1_r) = (v \cdot (w \cdot f))(1_r) - (-1)^{|v||w|}(w \cdot (v \cdot f))(1_r) \tag{3.14}$$

Using the definition of the operation $\cdot$ and the associativity axiom for the operad structure on $\mathcal{O}$ we get

$$\begin{aligned} (v \cdot (w \cdot f))(1_r) - (-1)^{|v||w|}(w \cdot (v \cdot f))(1_r) = \\ (-1)^{|f|(|v|+|w|)+|v||w|} \sum_{p\ge 1\, q\ge 0} \sum_{\tau \in \mathrm{Sh}_{p,q,r-p-q}} \mu_{\mathbf{t}^{p,q}_{\tau}} (f(1_{r-p-q+1}) \otimes w(1^{\mathfrak{c}}_{q+1}) \otimes v(1^{\mathfrak{c}}_p)) \\ + (-1)^{|f|(|v|+|w|)+|v||w|} \sum_{p,q\ge 1} \sum_{\tau \in \mathrm{Sh}_{p,q,r-p-q}} \mu_{\widetilde{\mathbf{t}}^{p,q}_{\tau}} (f(1_{r-p-q+2}) \otimes w(1^{\mathfrak{c}}_{q}) \otimes v(1^{\mathfrak{c}}_p)) \\ - (-1)^{|v||w|}(v \leftrightarrow w)\,, \end{aligned} \tag{3.15}$$

where the trees $\mathbf{t}^{p,q}_{\tau}$ and $\widetilde{\mathbf{t}}^{p,q}_{\tau}$ are depicted on figures 3.4 and 3.5 , respectively.

Since $f(1_{r-p-q+2})$ is invariant with respect to the action of $S_{r-p-q+2}$ the sums involving $\mu_{\widetilde{\mathbf{t}}^{p,q}_{\tau}}$ cancel each other. Furthermore, it is not hard to see that the sums involving $\mu_{\mathbf{t}^{p,q}_{\tau}}$ form the expression

$$[v,w] \cdot f(1_r)\,.$$



FIG. 3.4. The tree $\mathbf{t}^{p,q}_{\tau}$



FIG. 3.5. The tree $\widetilde{\mathbf{t}}^{p,q}_{\tau}$

Thus equation (3.14) follows. It remains to check that the operation $f \mapsto v \cdot f$ is an operadic derivation and we leave this step as an exercise for the reader.

□

3.1. **The action of $\mathcal{L}_{\mathcal{O}}$ on $\widetilde{\mathrm{Tw}}\,\mathcal{O}$.** Let us view $\widetilde{\mathrm{Tw}}\,\mathcal{O}(1)$ as the Lie algebra with the bracket being commutator.

We have an obvious degree zero map

$$\kappa : \mathcal{L}_{\mathcal{O}} \to \widetilde{\mathrm{Tw}}\,\mathcal{O}(1)$$

defined by the formula:

$$\kappa(v)(1_r) = v(1^{\mathfrak{c}}_{r+1})\,. \qquad (3.16)$$

where, as above, 1_r is the generator $\mathbf{s}^{-2r}\,1 \in \mathbf{s}^{-2r}\,\mathbb{K}$ and $1^{\mathfrak{c}}_r$ is the generator $\mathbf{s}^{2-2r}\,1 \in \Lambda^2\mathsf{coCom}(r) \cong \mathbf{s}^{2-2r}\,\mathbb{K}$.

We have the following proposition.

Proposition 3.2. *Let us form the semi-direct product* $\mathcal{L}_{\mathcal{O}} \ltimes \widetilde{\mathrm{Tw}}\,\mathcal{O}(1)$ *of the dg Lie algebras* $\mathcal{L}_{\mathcal{O}}$ *and* $\widetilde{\mathrm{Tw}}\,\mathcal{O}(1)$ *using the action of* $\mathcal{L}_{\mathcal{O}}$ *on* $\widetilde{\mathrm{Tw}}\,\mathcal{O}$ *defined in Proposition 3.1. Then the formula*

$$\Theta(v) = v + \kappa(v) \tag{3.17}$$

defines a Lie algebra homomorphism

$$\Theta : \mathcal{L}_{\mathcal{O}} \to \mathcal{L}_{\mathcal{O}} \ltimes \widetilde{\mathrm{Tw}}\,\mathcal{O}(1)\,.$$

Proof. First, let us prove that for every pair of homogeneous vectors $v, w \in \mathcal{L}_{\mathcal{O}}$ we have

$$\kappa([v,w]) = [\kappa(v), \kappa(w)] + v \cdot \kappa(w) - (-1)^{|v||w|} w \cdot \kappa(v)\,. \tag{3.18}$$

Indeed, unfolding the definition of κ we get[7]

$$\kappa([v,w])(1_r) = \sum_{p=1}^{r} \sum_{\tau \in \mathrm{Sh}_{p,r-p}} v_{r-p+2}\big(w_p(\tau(1),\dots,\tau(p)), \tau(p+1), \dots, \tau(r), r+1\big) \tag{3.19}$$
$$+ \sum_{p=0}^{r} \sum_{\tau \in \mathrm{Sh}_{p,r-p}} v_{r-p+1}\big(w_{p+1}(\tau(1),\dots,\tau(p), r+1), \tau(p+1), \dots, \tau(r)\big)$$
$$-(-1)^{|v||w|}(v \leftrightarrow w)\,,$$

where $v_t = v(1_t)$ and $w_t = w(1_t)$. The first sum in (3.19) equals

$$-(-1)^{|v||w|}(w \cdot \kappa(v))(1_r)\,.$$

Furthermore, since $v(1_t)$ is invariant under the action of S_t, we see that the second sum in (3.19) equals

$$\big(\kappa(v) \circ_1 \kappa(w)\big)\,(1_r)\,.$$

Thus equation (3.18) holds. Now, using (3.18), it is easy to see that

$$[v + \kappa(v), w + \kappa(w)] = [v,w] + v \cdot \kappa(w) - (-1)^{|v||w|} w \cdot \kappa(v) + [\kappa(v), \kappa(w)] =$$
$$= [v,w] + \kappa([v,w])$$

and the statement of the proposition follows. □

The following corollaries are immediate consequences of Proposition 3.2.

Corollary 3.3. *For* $v \in \mathcal{L}_{\mathcal{O}}$ *and* $f \in \widetilde{\mathrm{Tw}}\,\mathcal{O}(n)$ *the formula*

$$f \to v \cdot f + \delta_{\kappa(v)}(f) \tag{3.20}$$

defines an action of the Lie algebra $\mathcal{L}_{\mathcal{O}}$ *on the operad* $\widetilde{\mathrm{Tw}}\,\mathcal{O}$. □

Corollary 3.4. *For every Maurer-Cartan element* $\varphi \in \mathcal{L}_{\mathcal{O}}$, *the sum*

$$\varphi + \kappa(\varphi)$$

is a Maurer-Cartan element of the Lie algebra $\mathcal{L}_{\mathcal{O}} \ltimes \widetilde{\mathrm{Tw}}\,\mathcal{O}(1)$. □

---

[7] Here we use the notation for operadic multiplications (2.18) from Section 2.2.3.

We finally give the definition of the operad $\mathrm{Tw}\,\mathcal{O}$.

Definition 3.5. Let $\mathcal{O}$ be an operad in $\mathsf{Ch}_{\mathbb{K}}$ and φ be a Maurer-Cartan element in $\mathcal{L}_{\mathcal{O}}$ (3.5) corresponding to an operad morphism $\widehat{\varphi}$ (3.1). Let us also denote by $\partial^{\mathcal{O}}$ the differential on $\widetilde{\mathrm{Tw}\,\mathcal{O}}$ coming from the one on $\mathcal{O}$. We define the operad $\mathrm{Tw}\,\mathcal{O}$ in $\mathsf{Ch}_{\mathbb{K}}$ by declaring that $\mathrm{Tw}\,\mathcal{O} = \widetilde{\mathrm{Tw}\,\mathcal{O}}$ as operads in $\mathsf{grVect}_{\mathbb{K}}$ and letting

$$\partial^{\mathrm{Tw}} = \partial^{\mathcal{O}} + \varphi\cdot + \delta_{\kappa(\varphi)} \tag{3.21}$$

be the differential on $\mathrm{Tw}\,\mathcal{O}$.

Corollaries 3.3 and 3.4 imply that ∂^{Tw} is indeed a differential on $\mathrm{Tw}\,\mathcal{O}$.

Remark 3.6. *It is easy to see that, if* $\mathcal{O}(0) = \mathbf{0}$ *then the cochain complexes* $\mathbf{s}^{-2}\mathrm{Tw}\,\mathcal{O}(0)$ *and* $\mathcal{L}_{\mathcal{O}}$ *(3.5) are tautologically isomorphic.*

3.2. **Algebras over** $\mathrm{Tw}\,\mathcal{O}$**.** Let us assume that V is an algebra over $\mathcal{O}$ equipped with a complete decreasing filtration (3.2). We also assume that the $\mathcal{O}$-algebra structure on V is compatible with this filtration.

Given a Maurer-Cartan element $\alpha \in \mathcal{F}_1 V$, the formula

$$\partial^{\alpha}(v) = \partial(v) + \sum_{r=1}^{\infty} \frac{1}{r!}\varphi(1^{\mathfrak{c}}_{r+1})(\alpha, \dots, \alpha, v) \tag{3.22}$$

defines a new (twisted) differential on V. We will denote by V^{α} the cochain complex V with this new differential. In this setting we have the following theorem:

Theorem 3.7. *If* V^{α} *is the cochain complex obtained from* V *via twisting the differential by the Maurer-Cartan element* α *then the formula*

$$f(v_1, \dots, v_n) = \sum_{r=0}^{\infty} \frac{1}{r!} f(1_r)(\alpha, \dots, \alpha, v_1, \dots, v_n) \tag{3.23}$$

$$f \in \mathrm{Tw}\,\mathcal{O}(n)\,, \qquad v_i \in V$$

defines a $\mathrm{Tw}\,\mathcal{O}$*-algebra structure on* V^{α}.

Proof. Let $f \in \mathrm{Tw}\,\mathcal{O}(n)$, $g \in \mathrm{Tw}\,\mathcal{O}(k)$,

$$f_r := f(1_r) \in \big(\mathcal{O}(r+n)\big)^{S_r}\,, \qquad \text{and} \qquad g_r = g(1_r) \in \big(\mathcal{O}(r+k)\big)^{S_r}\,.$$

Our first goal is to verify that

$$(-1)^{|g|(|v_i|+\dots+|v_{i-1}|)} f(v_1, \dots, v_{i-1}, g(v_i, \dots, v_{i+k-1}), v_{i+k}, \dots, v_{n+k-1}) \tag{3.24}$$
$$= f \circ_i g(v_1, \dots, v_{n+k-1})\,.$$

The left hand side of (3.24) can be rewritten as

$$(-1)^{|g|(|v_i|+\dots+|v_{i-1}|)} f(v_1, \dots, v_{i-1}, g(v_i, \dots, v_{i+k-1}), v_{i+k}, \dots, v_{n+k-1}) =$$

$$\sum_{p,q\ge 0} \frac{(-1)^{|g|(|v_i|+\dots+|v_{i-1}|)}}{p!q!} f_p(\alpha,\dots,\alpha,v_1,\dots,v_{i-1},g_q(\alpha,\dots,\alpha,v_i,\dots,v_{i+k-1}),v_{i+k},\dots,v_{n+k-1})\,.$$

Using the obvious combinatorial identity

$$|\mathrm{Sh}_{p,q}| = \frac{(p+q)!}{p!q!} \tag{3.25}$$

we rewrite the left hand side of (3.24) further

$$\text{L.H.S. of (3.24)} =$$

$$\sum_{p,q\ge 0} \frac{(-1)^{|g|(|v_i|+\dots+|v_{i-1}|)}}{(p+q)!} |\mathrm{Sh}_{p,q}| f_p(\alpha,\dots,\alpha,v_1,\dots,v_{i-1},g_q(\alpha,\dots,\alpha,v_i,\dots,v_{i+k-1}),v_{i+k},\dots,v_{n+k-1}) =$$

$$\sum_{r=0}^{\infty}\frac{1}{r!}\sum_{p=0}^{r}\sum_{\sigma\in \mathrm{Sh}_{p,r-p}} \sigma\circ\varrho_{r,p,i}(f_p\circ_{p+i} g_{r-p})(\underbrace{\alpha,\dots,\alpha}_{r \text{ arguments}},v_1,\dots,v_{n+k-1}),$$

where $\varrho_{r,p,i}$ is the following permutation in S_{r+i-1}

$$\varrho_{r,p,i} = \begin{pmatrix} p+1 & \dots & p+i-1 & p+i & \dots & r+i-1 \\ r+1 & \dots & r+i-1 & p+1 & \dots & r \end{pmatrix}. \tag{3.26}$$

Thus, using (3.11), we get

$$\text{L.H.S. of (3.24)} = f\circ_i g(v_1,\dots,v_{n+k-1})$$

and equation (3.24) holds.

Next, we need to show that

$$\begin{aligned} \partial^{\mathrm{Tw}}(f)(v_1,\dots,v_n) &= \partial^{\alpha} f(v_1,\dots,v_n) \\ &\quad - (-1)^{|f|}\sum_{i=1}^{n}(-1)^{|v_i|+\dots+|v_{i-1}|} f(v_1,\dots,v_{i-1},\partial^{\alpha}(v_i),v_{i+1},\dots,v_n)\,. \end{aligned} \tag{3.27}$$

The right hand side of (3.27) can be rewritten as

$$\text{R.H.S. of (3.27)} =$$

$$\sum_{p\ge 0}\frac{1}{p!}\partial f_p(\alpha,\dots,\alpha,v_1,\dots,v_n) + \sum_{p\ge 0,q\ge 1}\frac{1}{p!q!}\varphi_q(\alpha,\dots,\alpha,f_p(\alpha,\dots,\alpha,v_1,\dots,v_n))$$

$$-(-1)^{|f|}\sum_{i=1}^{n}\sum_{p\ge 0}\frac{(-1)^{|v_i|+\dots+|v_{i-1}|}}{p!} f_p(\alpha,\dots,\alpha,v_1,\dots,v_{i-1},\partial(v_i),v_{i+1},\dots,v_n)$$

$$-(-1)^{|f|}\sum_{i=1}^{n}\sum_{p\ge 0,q\ge 1}\frac{(-1)^{|v_i|+\dots+|v_{i-1}|}}{p!q!} f_p(\alpha,\dots,\alpha,v_1,\dots,v_{i-1},\varphi_q(\alpha,\dots,\alpha,v_i),v_{i+1},\dots,v_n)\,,$$

where $f_p = f(1_p)$ and $\varphi_q = \varphi(1^{\mathfrak{c}}_q)$.

Let us now add to and subtract from the right hand side of (3.27) the sum

$$-(-1)^{|f|}\sum_{p\ge 0}\frac{1}{p!} f_{p+1}(\partial\alpha,\alpha,\dots,\alpha,v_1,\dots,v_n)\,.$$

Using the symmetry of $f_p = f(1_p)$ with respect to the action of the subgroup $S_p \subset S_{p+n}$, we get

$$\text{R.H.S. of } (3.27) =$$

$$\sum_{p\ge 0} \frac{1}{p!} \partial f_p(\alpha, \dots, \alpha, v_1, \dots, v_n) - (-1)^{|f|} \sum_{p \ge 0} \frac{1}{p!} f_{p+1}(\partial\alpha, \alpha, \dots, \alpha, v_1, \dots, v_n)$$

$$-(-1)^{|f|} \sum_{i=1}^{n} \sum_{p\ge 0} \frac{(-1)^{|v_i|+\dots+|v_{i-1}|}}{p!} f_p(\alpha, \dots, \alpha, v_1, \dots, v_{i-1}, \partial(v_i), v_{i+1}, \dots, v_n)$$

$$+(-1)^{|f|} \sum_{p\ge 0} \frac{1}{p!} f_{p+1}(\partial\alpha, \alpha, \dots, \alpha, v_1, \dots, v_n)$$

$$+ \sum_{p\ge 0, q\ge 1} \frac{1}{p!q!} \varphi_q(\alpha, \dots, \alpha, f_p(\alpha, \dots, \alpha, v_1, \dots, v_n))$$

$$-(-1)^{|f|} \sum_{i=1}^{n} \sum_{p\ge 0, q\ge 1} \frac{(-1)^{|v_i|+\dots+|v_{i-1}|}}{p!q!} f_p(\alpha, \dots, \alpha, v_1, \dots, v_{i-1}, \varphi_q(\alpha, \dots, \alpha, v_i), v_{i+1}, \dots, v_n) =$$

$$(\partial^{\mathcal{O}} f)(v_1, \dots, v_n)$$

$$+(-1)^{|f|} \sum_{p\ge 0} \frac{1}{p!} f_{p+1}(\partial\alpha, \alpha, \dots, \alpha, v_1, \dots, v_n)$$

$$+ \sum_{p\ge 0, q\ge 1} \frac{1}{p!q!} \varphi_q(\alpha, \dots, \alpha, f_p(\alpha, \dots, \alpha, v_1, \dots, v_n))$$

$$-(-1)^{|f|} \sum_{i=1}^{n} \sum_{p\ge 0, q\ge 1} \frac{(-1)^{|v_i|+\dots+|v_{i-1}|}}{p!q!} f_p(\alpha, \dots, \alpha, v_1, \dots, v_{i-1}, \varphi_q(\alpha, \dots, \alpha, v_i), v_{i+1}, \dots, v_n)\,.$$

Due to the Maurer-Cartan equation for α

$$\partial(\alpha) + \frac{1}{q!} \varphi_q(\alpha, \alpha, \dots, \alpha) = 0$$

we have

$$+(-1)^{|f|} \sum_{p\ge 0} \frac{1}{p!} f_{p+1}(\partial\alpha, \alpha, \dots, \alpha, v_1, \dots, v_n) =$$

$$-(-1)^{|f|} \sum_{p\ge 0, q\ge 2} \frac{1}{p!q!} f_{p+1}(\varphi_q(\alpha, \dots, \alpha), \alpha, \dots, \alpha, v_1, \dots, v_n)\,.$$

Hence, using combinatorial formula (3.25), we get

$$\text{R.H.S. of } (3.27) =$$

$$(\partial^{\mathcal{O}} f)(v_1, \dots, v_n) + (\varphi \cdot f)(v_1, \dots, v_n)$$

$$\kappa(\varphi) \circ_1 f(v_1, \dots, v_n) - (-1)^{|f|} f \circ_1 \kappa(\varphi)(v_1, \dots, v_n)\,.$$

Theorem 3.7 is proven. □

Let us now observe that the dg operad $\mathrm{Tw}\,\mathcal{O}$ is equipped with a complete descending filtration. Namely,

$$\mathcal{F}_k \mathrm{Tw}\,\mathcal{O}(n) = \{ f \in \mathrm{Tw}\,\mathcal{O}(n) \mid f(1_r) = 0 \quad \forall\, r < k \}\,. \tag{3.28}$$

It is clear that the operad structure on $\mathrm{Tw}\,\mathcal{O}$ is compatible with this filtration. The endomorphism operad End_V also carries a complete descending filtration since so does V. This observation motivates the following definition:

Definition 3.8. A filtered $\mathrm{Tw}\,\mathcal{O}$-algebra is a cochain complex V equipped with a complete descending filtration for which the operad map

$$\mathrm{Tw}\,\mathcal{O} \to \mathsf{End}_V$$

is compatible with the filtrations. We denote by $\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}}$ the category of filtered $\mathrm{Tw}\,\mathcal{O}$-algebra.

It is easy to see that the $\mathrm{Tw}\,\mathcal{O}$-algebra V^{α} from Theorem 3.7 is a filtered $\mathrm{Tw}\,\mathcal{O}$-algebra in the sense of this definition. The next theorem provides us with an equivalent description of the category of filtered $\mathrm{Tw}\,\mathcal{O}$-algebras.

Theorem 3.9. *Let* $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$ *be the category of pairs* (V, α)*, where* V *is an* $\mathcal{O}$*-algebra equipped with a complete descending filtration as in* (3.2) *and* $\alpha \in \mathcal{F}_1 V$ *is a Maurer-Cartan element. Morphisms between pairs* (V, α) *and* (V', α') *are morphisms of filtered* $\mathcal{O}$*-algebras* $f : V \to V'$ *which satisfy the condition* $f(\alpha) = \alpha'$. *The category* $\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}}$ *of filtered* $\mathrm{Tw}\,\mathcal{O}$*-algebras is isomorphic to the category* $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$.

Proof. Theorem 3.7 yields a functor

$$\mathfrak{F} : \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}} \to \mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}} \tag{3.29}$$

from the category $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$ to the category $\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}}$ of filtered $\mathrm{Tw}\,\mathcal{O}$-algebras. The functor $\mathfrak{F}$ assigns to a pair (V, α) the cochain complex V^{α} with the differential ∂^{α} (3.22) and the $\mathrm{Tw}\,\mathcal{O}$-algebra structure defined by equation (3.23).

To define a functor in the opposite direction we produce a degree 2 element $u^{\circ} \in \mathcal{F}_1 \mathrm{Tw}\,\mathcal{O}(0)$ using the identity element $u_{\mathcal{O}} \in \mathcal{O}(1)$:

$$u^{\circ}(1_r) := \begin{cases} u_{\mathcal{O}} & \text{if} \quad r = 1 \\ 0 & \qquad \text{otherwise}\,. \end{cases} \tag{3.30}$$

Let, as above, φ be the Maurer-Cartan element in $\mathrm{Conv}(\Lambda^2\mathsf{coCom}, \mathcal{O})$ corresponding to the morphism $\hat{\varphi} : \Lambda\mathsf{Lie}_{\infty} \to \mathcal{O}$. Also, let $\varphi_r := \varphi(1^{\mathfrak{c}}_r)$.

Since S_r acts trivially on φ_r, and $\partial^{\mathcal{O}}(u_{\mathcal{O}}) = 0$ we get

$$\partial^{\mathrm{Tw}}(u^{\circ})(1_r) = \sum_{\sigma \in \mathrm{Sh}_{r-1,1}} \sigma\big(\varphi_r \circ_r u_{\mathcal{O}}\big) - u_{\mathcal{O}} \circ_1 \varphi_r = r\varphi_r - \varphi_r\,.$$

Thus

$$\partial^{\mathrm{Tw}}(u^{\circ})(1_r) = (r-1)\varphi_r\,. \tag{3.31}$$

Let us now consider a filtered $\mathrm{Tw}\,\mathcal{O}$-algebra W with the differential ∂_W. The element $u^{\circ} \in \mathcal{F}_1 \mathrm{Tw}\,\mathcal{O}(0)$ gives us a degree 2 vector in W. We denote this vector by α and observe that $\alpha \in \mathcal{F}_1 W$ since the map $\mathrm{Tw}\,\mathcal{O} \to \mathsf{End}_W$ is compatible with the filtrations.

Next, we remark that the formula

$$\iota(\gamma)(1_r) := \begin{cases} \gamma & \text{if } r = 0 \\ 0 & \text{otherwise}\,, \end{cases} \qquad \gamma \in \mathcal{O}(n) \tag{3.32}$$

defines an embedding

$$\iota : \mathcal{O} \hookrightarrow \mathrm{Tw}\,\mathcal{O} \tag{3.33}$$

of operads in the category $\mathsf{grVect}_{\mathbb{K}}$.

Thus, using ι and the $\mathrm{Tw}\,\mathcal{O}$-algebra structure on W, we get an $\mathcal{O}$-algebra structure on W. Since, in general, ι is not compatible with the differentials, this $\mathcal{O}$-algebra structure on W is not compatible with the differential ∂_W on W.

Let r be a non-negative integer. Then, for every vector

$$f_r \in \big(\mathcal{O}(r+n)\big)^{S_r}$$

the elements f_r, $\mathbf{s}^2 f_r$, $\mathbf{s}^4 f_r$, …, $\mathbf{s}^{2r} f_r$ may be viewed as vectors in $\mathrm{Tw}\,\mathcal{O}(r+n)$, $\mathrm{Tw}\,\mathcal{O}(r-1+n)$, $\mathrm{Tw}\,\mathcal{O}(r-2+n)$, …, $\mathrm{Tw}\,\mathcal{O}(n)$, respectively. Namely,

$$\mathbf{s}^{2k} f_r(1_p) := \begin{cases} f_r & \text{if } p = k\,, \\ 0 & \text{otherwise}\,. \end{cases} \tag{3.34}$$

Let us consider the vector $\mathbf{s}^{2k} f_r \circ_1 u^{\circ} \in \mathrm{Tw}\,\mathcal{O}(r-k-1+n)$ for $0 \le k \le r-1$. It is clear that for $p \neq k+1$

$$\big(\mathbf{s}^{2k} f_r \circ_1 u^{\circ}\big)(1_p) = 0\,.$$

For $p = k+1$ we use the S_r-invariance of f_r and get

$$\big(\mathbf{s}^{2k} f_r \circ_1 u^{\circ}\big)(1_{k+1}) = \sum_{\sigma \in \mathrm{Sh}_{k,1}} \sigma(f_r \circ_{k+1} u_{\mathcal{O}}) = (k+1) f_r\,.$$

Thus

$$\mathbf{s}^{2k} f_r \circ_1 u^{\circ} = (k+1)\mathbf{s}^{2k+2} f_r\,. \tag{3.35}$$

Applying identity (3.35) r times we get

$$(\dots((f_r \underbrace{\circ_1 u^{\circ}) \circ_1 u^{\circ}) \cdots \circ_1 u^{\circ})}_{r \text{ times}} = r!\mathbf{s}^{2r} f_r\,. \tag{3.36}$$

Combining this observation with the fact that the filtration on W is complete, we conclude that for every vector $f \in \mathrm{Tw}\,\mathcal{O}(n)$ with $f_r = f(1_r) \in \big(\mathcal{O}(r+n)\big)^{S_r}$ we have

$$f(w_1, w_2, \dots, w_n) = \sum_{r=0}^{\infty} \frac{1}{r!} \iota(f_r)(\alpha, \dots, \alpha, w_1, w_2, \dots, w_n)\,, \qquad w_i \in W\,, \tag{3.37}$$

where ι is the map of operads $\mathcal{O} \to \mathrm{Tw}\,\mathcal{O}$ in $\mathsf{grVect}_{\mathbb{K}}$ defined in (3.32).

Let us denote by ∂ the degree 1 operation

$$\partial : W \to W$$

defined by the equation

$$\partial(w) := \partial_W(w) - \sum_{r=1}^{\infty} \frac{1}{r!} \iota(\varphi_{r+1})(\alpha, \dots, \alpha, w)\,, \tag{3.38}$$

where, as above, $\varphi_q = \varphi(1^{\mathfrak{c}}_q)$.

To prove the identity $\partial^2 = 0$, we observe that, due to equation (3.37),

$$\partial(w) = \partial_W(w) - \kappa(\varphi)(w)\,. \tag{3.39}$$

Hence

$$\begin{gathered}
\partial^2(w) = -\partial_W(\kappa(\varphi)(w)) - \kappa(\varphi)(\partial_W(w)) + \big(\kappa(\varphi) \circ_1 \kappa(\varphi)\big)(w) = \\
-\big(\partial^{\mathrm{Tw}}\kappa(\varphi)\big)(w) + \big(\kappa(\varphi) \circ_1 \kappa(\varphi)\big)(w) = \\
-\big(\partial^{\mathcal{O}}\kappa(\varphi)\big)(w) - \big(\varphi \cdot \kappa(\varphi)\big)(w) - 2\big(\kappa(\varphi) \circ_1 \kappa(\varphi)\big)(w) + \big(\kappa(\varphi) \circ_1 \kappa(\varphi)\big)(w) = \\
-\big(\partial^{\mathcal{O}}\kappa(\varphi) + \varphi \cdot \kappa(\varphi) + \kappa(\varphi) \circ_1 \kappa(\varphi)\big)(w)\,.
\end{gathered}$$

On the other hand, the vector $\partial^{\mathcal{O}}\kappa(\varphi) + \varphi \cdot \kappa(\varphi) + \kappa(\varphi) \circ_1 \kappa(\varphi) \in \mathrm{Tw}\,\mathcal{O}(1)$ is zero due to Corollary 3.4 .

Thus equation (3.38) (or equation (3.39)) defines another differential on W .

Let us show that the differential ∂ is compatible with the $\mathcal{O}$-algebra structure given by ι (3.33).

Equation (3.39) implies that for every $\gamma \in \mathcal{O}(n)$ and $w_1, w_2, \dots, w_n \in W$, we have

$$\begin{gathered}
\partial\big(\iota(\gamma)(w_1, \dots, w_n)\big) - \sum_{i=1}^{n} (-1)^{|\gamma| + |w_1| + \dots + |w_{i-1}|} \iota(\gamma)(w_1, \dots, w_{i-1}, \partial(w_i), w_{i+1}, \dots, w_n) = \\
\partial_W\big(\iota(\gamma)(w_1, \dots, w_n)\big) - \sum_{i=1}^{n} (-1)^{|\gamma| + |w_1| + \dots + |w_{i-1}|} \iota(\gamma)(w_1, \dots, w_{i-1}, \partial_W(w_i), w_{i+1}, \dots, w_n) \\
-\kappa(\varphi)\big(\iota(\gamma)(w_1, \dots, w_n)\big) \\
+ \sum_{i=1}^{n} (-1)^{|\gamma| + |w_1| + \dots + |w_{i-1}|} \iota(\gamma)(w_1, \dots, w_{i-1}, \kappa(\varphi)(w_i), w_{i+1}, \dots, w_n) = \\
\big(\partial^{\mathrm{Tw}}\iota(\gamma)\big)(w_1, \dots, w_n) - \big(\kappa(\varphi) \circ_1 \iota(\gamma)\big)(w_1, \dots, w_n) + (-1)^{|\gamma|} \sum_{i=1}^{n} \big(\iota(\gamma) \circ_1 \kappa(\varphi)\big)(w_1, \dots, w_n) = \\
\big(\iota(\partial^{\mathcal{O}}\gamma)\big)(w_1, \dots, w_n)\,,
\end{gathered}$$

where, in the last step, we used the fact that $\iota(\gamma)(1_r) = 0$ for all $r \geq 1$.

Thus the differential ∂ (3.38) is indeed compatible with the $\mathcal{O}$-algebra structure on W.

It remains to prove that the vector $\alpha \in W$ is a Maurer-Cartan element (with respect to the differential ∂).

For this purpose we observe that, since the map $\mathrm{Tw}\,\mathcal{O} \to \mathsf{End}_W$ is compatible with the differentials, the vector $\partial^{\mathrm{Tw}}\,u^{\circ}$ maps to $\partial_W(\alpha)$ in W.

Therefore, combining equation (3.31) with equation (3.37) we get

$$\partial_W(\alpha) = \sum_{r=2}^{\infty} \frac{r-1}{r!} \iota(\varphi_r)(\alpha, \dots, \alpha)$$

or equivalently

$$\partial(\alpha) + \sum_{r=1}^{\infty} \frac{1}{r!} \iota(\varphi_{r+1})(\alpha, \dots, \alpha) = \sum_{r=2}^{\infty} \frac{r-1}{r!} \iota(\varphi_r)(\alpha, \dots, \alpha)\,.$$

Thus α indeed satisfies the Maurer-Cartan equation

$$\partial(\alpha) + \sum_{r=2}^{\infty} \frac{1}{r!} \iota(\varphi_r)(\alpha, \dots, \alpha) = 0\,. \tag{3.40}$$

Combining the performed work, we conclude that the cochain complex W with the differential (3.38) and the Maurer-Cartan element α is an object of the category $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$.

Equations (3.37) and (3.38) imply that the described construction, indeed, gives a (strict) inverse for the functor $\mathfrak{F}$ (3.29). □

3.3. **Example: The dg operad $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$.** In this subsection, we apply the twisting procedure to the pair $(\Lambda\mathsf{Lie}_\infty, \hat{\varphi})$, where

$$\hat{\varphi} = \mathrm{id} : \Lambda\mathsf{Lie}_\infty \to \Lambda\mathsf{Lie}_\infty\,. \tag{3.41}$$

Let us recall that $1^{\mathfrak{c}}_r$ denotes the canonical generator $\mathbf{s}^{2-2r}\,1 \in \Lambda^2\mathsf{coCom}_{\circ}(r)$. Thus, the Maurer-Cartan element φ corresponding to (3.41) is

$$\varphi(1^{\mathfrak{c}}_r) = \mathbf{s}\,1^{\mathfrak{c}}_r\,, \qquad r \geq 2 \tag{3.42}$$

and the element $\kappa(\varphi) \in \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty(1)$ takes the form

$$\kappa(\varphi)(1_r) = \begin{cases} \mathbf{s}\,1^{\mathfrak{c}}_{r+1} & \text{if } r \geq 1\,, \\ 0 & \text{if } r = 0\,. \end{cases} \tag{3.43}$$

According to general formula (3.21), the differential ∂^{Tw} on $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$ is given by the equation

$$\partial^{\mathrm{Tw}}(f)(1_r) = \partial^{\mathrm{Cobar}}(f_r) - (-1)^{|f|} \sum_{p=2}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \sigma\big(f_{r-p+1} \circ_1 (\mathbf{s}\,1^{\mathfrak{c}}_p)\big) \tag{3.44}$$

$$+ \sum_{p=1}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \sigma\big((\mathbf{s}\,1^{\mathfrak{c}}_{p+1}) \circ_{p+1} f_{r-p}\big) - (-1)^{|f|} \sum_{i=1}^{n} \sum_{p=1}^{r-1} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \sigma \circ \varrho_{r,p,i}\big(f_p \circ_{p+i} (\mathbf{s}\,1^{\mathfrak{c}}_{r-p+1})\big)\,.$$

where $f \in \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty(n)$, $f_r = f(1_r)$ and $\varrho_{r,p,i}$ is the permutation of S_{r+i-1} defined in (3.26).

We will need the following lemma:

Lemma 3.10. *If* $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$ *is the dg operad which is obtained from* $\Lambda\mathsf{Lie}_\infty$ *via applying the twisting procedure to the identity map* (3.41) *then the equation*

$$\mathfrak{T}(\mathbf{s}1^{\mathfrak{c}}_n)(1_r) = \mathbf{s}1^{\mathfrak{c}}_{r+n} \tag{3.45}$$

defines a map of dg operads

$$\mathfrak{T} : \Lambda\mathsf{Lie}_\infty \ \to \ \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty\,. \tag{3.46}$$

Proof. According to (2.57)

$$\Lambda\mathsf{Lie}_\infty = \mathrm{Cobar}(\Lambda^2\mathsf{coCom})\,.$$

Hence, due to Theorem 2.6, maps of dg operads from $\Lambda\mathsf{Lie}_\infty$ to $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$ are in bijection with Maurer-Cartan elements of the convolution Lie algebra

$$\mathrm{Conv}(\Lambda^2\mathsf{coCom}_\circ, \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty) = \prod_{n\ge 2} \mathsf{Hom}_{S_n}\big(\mathbf{s}^{2-2n}\mathbb{K}, \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty(n)\big)\,. \tag{3.47}$$

Let us denote by $\alpha_{\mathfrak{T}}$ a vector in (3.47) defined by the equation:

$$\alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_n)(1_r) = \mathbf{s}1^{\mathfrak{c}}_{r+n}\,. \tag{3.48}$$

Since the vectors $1^{\mathfrak{c}}_n$ and 1_n carry degrees

$$|1^{\mathfrak{c}}_n| = 2-2n\,, \qquad |1_n| = -2n\,,$$

the vector $\alpha_{\mathfrak{T}}$ has degree 1.

Using formula (3.44) for the differential ∂^{Tw} on $\mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$ we get (here $n\ge 2$)

$$\begin{gathered}\big(\partial^{\mathrm{Tw}}\alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_n)\big)(1_r) = \\ \partial^{\mathrm{Cobar}}(\mathbf{s}1^{\mathfrak{c}}_{r+n}) + \sum_{p=2}^{r}\sum_{\sigma\in \mathrm{Sh}_{p,r-p}} \sigma\big(\mathbf{s}1^{\mathfrak{c}}_{r-p+1+n}\circ_1 \mathbf{s}1^{\mathfrak{c}}_p\big) + \\ \sum_{p=1}^{r}\sum_{\sigma\in \mathrm{Sh}_{p,r-p}} \sigma\big(\mathbf{s}1^{\mathfrak{c}}_{p+1}\circ_{p+1} \mathbf{s}1^{\mathfrak{c}}_{r-p+n}\big) + \\ \sum_{i=1}^{n}\sum_{p=1}^{r-1}\sum_{\sigma\in \mathrm{Sh}_{p,r-p}} \sigma\circ\varrho_{r,p,i}\big(\mathbf{s}1^{\mathfrak{c}}_{p+n}\circ_{p+i} \mathbf{s}1^{\mathfrak{c}}_{r-p+1}\big)\,,\end{gathered} \tag{3.49}$$

where $\varrho_{r,p,i}$ is the permutation defined in (3.26).

Unfolding $\partial^{\mathrm{Cobar}}(\mathbf{s}1^{\mathfrak{c}}_{r+n})$, we get

$$\begin{gathered}\big(\partial^{\mathrm{Tw}}\alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_n)\big)(1_r) = \\ -\sum_{q=2}^{r+n-1}\sum_{\lambda\in \mathrm{Sh}_{q,r+n-q}} \lambda\big(\mathbf{s}1^{\mathfrak{c}}_{r+n-q+1}\circ_1 \mathbf{s}1^{\mathfrak{c}}_q\big) + \\ \sum_{p=2}^{r}\sum_{\sigma\in \mathrm{Sh}_{p,r-p}} \sigma\big(\mathbf{s}1^{\mathfrak{c}}_{r-p+1+n}\circ_1 \mathbf{s}1^{\mathfrak{c}}_p\big) +\end{gathered} \tag{3.50}$$

$$\sum_{p=1}^{r} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \sigma\big(\mathbf{s}1^{\mathfrak{c}}_{p+1} \circ_{p+1} \mathbf{s}1^{\mathfrak{c}}_{r-p+n}\big) +$$

$$\sum_{i=1}^{n} \sum_{p=1}^{r-1} \sum_{\sigma \in \mathrm{Sh}_{p,r-p}} \sigma \circ \varrho_{r,p,i}\big(\mathbf{s}1^{\mathfrak{c}}_{p+n} \circ_{p+i} \mathbf{s}1^{\mathfrak{c}}_{r-p+1}\big)\,.$$

Next, using the definition of the binary operation (2.45) we get

$$\big(\alpha_{\mathfrak{T}} \bullet \alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_n)\big)(1_r) = \tag{3.51}$$

$$\sum_{q=2}^{n-1} \sum_{\tau \in \mathrm{Sh}_{q,n-q}} \Big(\tau\big(\alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_{n-q+1}) \circ_1 \alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_q)\big)\Big)(1_r) =$$

$$\sum_{0 \le p \le r}^{2 \le q \le n-1} \sum_{\sigma \in \mathrm{Sh}_{p,r-p,q,n-q}} \sigma\big(\,\mathbf{s}1^{\mathfrak{c}}_{n+p-q+1} \circ_{p+1} \mathbf{s}1^{\mathfrak{c}}_{r-p+q}\,\big)\,.$$

We observe that the second sum in the right hand side of (3.50) is obtained from

$$\sum_{q=2}^{r+n-1} \sum_{\lambda \in \mathrm{Sh}_{q,r+n-q}} \lambda\big(\mathbf{s}1^{\mathfrak{c}}_{r+n-q+1} \circ_1 \mathbf{s}1^{\mathfrak{c}}_q\big) \tag{3.52}$$

by keeping only the terms for which

$$\lambda(r+1) = r+1, \quad \lambda(r+2) = r+2, \quad \dots, \quad \lambda(r+n) = r+n\,.$$

Next, using the fact that S_m acts trivially on the vector $1^{\mathfrak{c}}_m$, we see that the third sum in the right hand side of (3.50) is obtained from (3.52) by keeping only the terms for which $q \ge n$ and

$$\lambda(q-n+1) = r+1\,, \quad \lambda(q-n+2) = r+2\,, \quad \dots, \quad \lambda(q) = r+n\,.$$

Similarly, the last sum in the right hand side of (3.50) is obtained from the sum (3.52) by keeping only the terms for which

$$\lambda(q) \ge r+1 \qquad \text{and} \qquad \lambda(i) \le r \quad \forall\ 1 \le i \le q-1\,.$$

Finally, using the invariance with respect to the action of the symmetric group once again, we see that $\big(\alpha_{\mathfrak{T}} \bullet \alpha_{\mathfrak{T}}(1^{\mathfrak{c}}_n)\big)(1_r)$ is obtained from the sum (3.52) by keeping only the terms for which

$$\lambda(q-1) \ge r+1\,, \qquad \text{and} \qquad \lambda(r+n) \ge r+1\,.$$

Thus, the vector $\alpha_{\mathfrak{T}}$ satisfies the Maurer-Cartan equation

$$\partial^{\mathrm{Tw}} \alpha_{\mathfrak{T}} + \alpha_{\mathfrak{T}} \bullet \alpha_{\mathfrak{T}} = 0\,. \tag{3.53}$$

Since the morphism of dg operads

$$\mathfrak{T} : \Lambda\mathsf{Lie}_{\infty} \ \to\ \mathrm{Tw}\,\Lambda\mathsf{Lie}_{\infty}$$

corresponding to the Maurer-Cartan element $\alpha_{\mathfrak{T}}$ is defined by equation (3.45), the lemma is proved. □

Remark 3.11. *Let V be a $\Lambda\mathsf{Lie}_\infty$-algebra with a differential ∂ and the brackets*

$$\{\ ,\ ,\dots,\ \}_n : V^{\otimes\, n} \to V\,.$$

Let us assume that V is equipped with a complete descending filtration (3.2) *and $\alpha \in \mathcal{F}_1 V$ is a Maurer-Cartan element. Theorem 3.7 and Lemma 3.10 imply that the formula*

$$\{v_1, \dots, v_n\}^{\alpha}_n = \sum_{r=0}^{\infty} \frac{1}{r!} \{\underbrace{\alpha, \dots, \alpha}_{r\ \ times}, v_1, \dots, v_n\}_{r+n} \tag{3.54}$$

defines a $\Lambda\mathsf{Lie}_\infty$-structure on V with the differential given by equation (3.22). *We say that this latter $\Lambda\mathsf{Lie}_\infty$-structure on V is obtained from the former one via twisting by the Maurer-Cartan element α.*

3.4. **A useful modification $\mathrm{Tw}^{\oplus}\mathcal{O}$.** In practice the morphism (3.1) often comes from the map (of dg operads)

$$\mathfrak{i} : \Lambda\mathsf{Lie} \to \mathcal{O}\,.$$

In this case, the above construction of twisting is well defined for the suboperad $\mathrm{Tw}^{\oplus}(\mathcal{O}) \subset \mathrm{Tw}\,\mathcal{O}$ with

$$\mathrm{Tw}^{\oplus}(\mathcal{O})(n) = \bigoplus_{r \ge 0} \mathbf{s}^{2r} \big(\mathcal{O}(r+n)\big)^{S_r}\,. \tag{3.55}$$

It is not hard to see that the Maurer-Cartan element

$$\varphi \in \mathrm{Conv}(\Lambda^2\mathsf{coCom}, \mathcal{O})$$

corresponding to the composition

$$\mathfrak{i} \circ U_{\Lambda\mathsf{Lie}} : \mathrm{Cobar}(\Lambda^2\mathsf{coCom}) \to \mathcal{O}$$

is given by the formula:

$$\varphi(1^{\mathfrak{c}}_r) = \begin{cases} \mathfrak{i}(\{a_1, a_2\}) & \text{if } r = 2 \\ 0 & \text{otherwise}\,. \end{cases} \tag{3.56}$$

Hence

$$\mathcal{L}^{\oplus}_{\mathcal{O}} = \bigoplus_{r \ge 0} \mathbf{s}^{2r-2} \big(\mathcal{O}(r)\big)^{S_r} \tag{3.57}$$

is a sub- dg Lie algebra of $\mathcal{L}_{\mathcal{O}}$ (3.5).

Specifying general formula (3.21) to this particular case, we see that the differential ∂^{Tw} on (3.55) is given by the equation:

$$\partial^{\mathrm{Tw}}(v) = -(-1)^{|v|} \sum_{\sigma \in \mathrm{Sh}_{2,r-1}} \sigma\big(v \circ_1 \mathfrak{i}(\{a_1, a_2\})\big) + \sum_{\tau \in \mathrm{Sh}_{1,r}} \tau\big(\mathfrak{i}(\{a_1, a_2\}) \circ_2 v\big) \tag{3.58}$$

$$-(-1)^{|v|} \sum_{\tau' \in \mathrm{Sh}_{r,1}} \sum_{i=1}^{n} \tau' \circ \varsigma_{r+1, r+i} \big(v \circ_{r+i} \mathfrak{i}(\{a_1, a_2\})\big)\,,$$

where

$$v \in \mathbf{s}^{2r}\big(\mathcal{O}(r+n)\big)^{S_r}\,,$$

and $\varsigma_{r+1,r+i}$ is the cycle $(r+1, r+2, \dots, r+i)$.

Remark 3.12. *We should remark that, when we apply elementary insertions in the right hand side of* (3.58)*, we view v and $\mathsf{i}(\{a_1,a_2\})$ as vectors in $\mathcal{O}(r+n)$ and $\mathcal{O}(2)$ respectively. The resulting sum in the right hand side of* (3.58) *is viewed as a vector in* $\mathrm{Tw}\,\mathcal{O}(n)$.

3.5. **More general version of twisting.** The reader may object that the version of twisting we have presented so far does not even cover the most classical case of the operad $\mathcal{O} = \mathsf{Lie}$, since in (3.1) we required a map $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$. However, this is easily repaired:

Let k be an integer and let

$$\Lambda^k\mathsf{Lie}_\infty \to \mathcal{O}$$

be an operad map. Then, by functoriality of Λ, there is an operad map

$$\Lambda\mathsf{Lie}_\infty \to \Lambda^{1-k}\mathcal{O}.$$

We then define:

$$\mathrm{Tw}\,\mathcal{O} := \Lambda^{k-1}\mathrm{Tw}\,(\Lambda^{1-k}\mathcal{O}).$$

In this paper we will always assume $k = 1$ since (i) this is the relevant case for the Deligne conjecture and (ii) the signs are significantly simpler for odd k. One should however be aware that all important statements may be transcribed to the arbitrary-k case by application of the functor Λ.

4. Categorial properties of twisting

4.1. Tw **as a comonad.** The goal of this section is to show that the operation Tw defines a comonad on the under-category $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads*, where *Operads* is the category of dg operads. Recall that the under-category $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads* is the category of arrows $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ in *Operads*, with morphisms the commutative diagrams

$$\begin{tikzcd}
 & \Lambda\mathsf{Lie}_\infty \arrow[dl] \arrow[dr] & \\
\mathcal{O} \arrow[rr] & & \mathcal{O}'
\end{tikzcd}.$$

4.1.1. Tw *is an endofunctor.* Consider arrows $\Lambda\mathsf{Lie}_\infty \to \mathcal{O} \xrightarrow{f} \mathcal{O}'$ in the category *Operads*. Twisting $\mathcal{O}$ and $\mathcal{O}'$, we obtain new dg operads with

$$\mathrm{Tw}\,\mathcal{O}(n) = \prod_{r\geq 0} \mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathcal{O}(r+n))\,,$$

$$\mathrm{Tw}\,\mathcal{O}'(n) = \prod_{r\geq 0} \mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathcal{O}'(r+n))\,.$$

By composing morphisms on the right with (components of) the map $f : \mathcal{O} \to \mathcal{O}'$ we obtain a map (of collections)

$$\mathrm{Tw}\, f \colon \mathrm{Tw}\, \mathcal{O} \to \mathrm{Tw}\, \mathcal{O}' \,.$$

Lemma 4.1. *The above map* $\mathrm{Tw}\, f : \mathrm{Tw}\, \mathcal{O} \to \mathrm{Tw}\, \mathcal{O}'$ *is a map of dg operads. Furthermore, if*

$$\Lambda\mathsf{Lie}_\infty \to \mathcal{O} \xrightarrow{f} \mathcal{O}' \xrightarrow{g} \mathcal{O}''$$

are maps of dg operads, then $\mathrm{Tw}\,(f \circ g) = (\mathrm{Tw}\, f) \circ (\mathrm{Tw}\, g)$.

Proof. It is clear since we only used natural operations (i.e., the operad structure) in defining $\mathrm{Tw}\, \mathcal{O}$. □

Applying Tw to $\hat{\varphi} : \Lambda\mathsf{Lie}_\infty \to \mathcal{O}$, we obtain a morphism of dg operads

$$\mathrm{Tw}\,(\hat{\varphi}) : \mathrm{Tw}\, \Lambda\mathsf{Lie}_\infty \to \mathrm{Tw}\, \mathcal{O} \,. \tag{4.1}$$

Composing (4.1) with the canonical map of dg operads

$$\mathfrak{T} : \Lambda\mathsf{Lie}_\infty \to \mathrm{Tw}\, \Lambda\mathsf{Lie}_\infty$$

from Lemma 3.10, we obtain a morphism of dg operads

$$\beta_{\mathcal{O}} := \mathrm{Tw}\,(\hat{\varphi}) \circ \mathfrak{T} : \Lambda\mathsf{Lie}_\infty \to \mathrm{Tw}\, \mathcal{O} \,. \tag{4.2}$$

From Lemma 4.1 it is then evident that the following diagram commutes:

$$\begin{tikzcd}
 & \Lambda\mathsf{Lie}_\infty \arrow[dl, "\beta_{\mathcal{O}}"'] \arrow[dr, "\beta_{\mathcal{O}'}"] & \\
\mathrm{Tw}\, \mathcal{O} \arrow[rr, "\mathrm{Tw}\, f"] & & \mathrm{Tw}\, \mathcal{O}'
\end{tikzcd}$$

Here $f, \mathcal{O}, \mathcal{O}'$ are as above. Summarizing, we obtain the following result:

Corollary 4.2. *The operation* Tw *defines an endofunctor on the under-category* $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads.* □

Remark 4.3. *Let* $*$ *be the initial object* (2.33) *in the category of dg operads. It is easy to see that* $\mathrm{Tw}\, *$ *carries the obvious augmentation*

$$\mathrm{Tw}\, * \to * \,. \tag{4.3}$$

Thus, if the dg operad $\mathcal{O}$ *in an object of* $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads has an augmentation morphism* $\varepsilon : \mathcal{O} \to *$ *then* $\mathrm{Tw}\, \mathcal{O}$ *is canonically augmented. The desired augmentation* $\mathrm{Tw}\, \mathcal{O} \to *$ *is obtained by composing*

$$\mathrm{Tw}\,(\varepsilon) : \mathrm{Tw}\, \mathcal{O} \to \mathrm{Tw}\, *$$

with (4.3).

4.1.2. *The natural projection.* Let $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ be an arrow in *Operads*. There is the natural map

$$\eta_{\mathcal{O}} : \mathrm{Tw}\,\mathcal{O} \to \mathcal{O} \tag{4.4}$$

projecting the product

$$\mathrm{Tw}\,\mathcal{O}(n) = \prod_{r\geq 0} \mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathcal{O}(r+n))$$

to its first ($r = 0$) factor. It is easy to see that (4.4) is a map of dg operads.

We claim that

Lemma 4.4. *The maps $\eta_{\mathcal{O}}$ assemble to form a natural transformation $\eta : \mathrm{Tw} \Rightarrow id$, where id is the identity functor on $\Lambda\mathsf{Lie}_\infty \downarrow$ Operads.*

Proof. We have to show that for all arrows $\Lambda\mathsf{Lie}_\infty \to \mathcal{O} \to \mathcal{O}'$ the following diagram commutes.

$$\begin{tikzcd}
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,f"] \arrow[d, "\eta_{\mathcal{O}}"'] & \mathrm{Tw}\,\mathcal{O}' \arrow[d, "\eta_{\mathcal{O}'}"] \\
\mathcal{O} \arrow[r, "f"] & \mathcal{O}'
\end{tikzcd}$$

This is obvious. □

4.1.3. *The comultiplication $\mathfrak{D} : \mathrm{Tw} \to \mathrm{Tw} \circ \mathrm{Tw}$.* Let again $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ be an arrow in *Operads*. Consider

$$\mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} = \prod_{r,s\geq 0} \mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathsf{Hom}_{S_s}(\mathbf{s}^{-2s}\mathbb{K}, \mathcal{O}(s+r+n))) \cong \prod_{r,s\geq 0} \mathsf{Hom}_{S_s\times S_r}(\mathbf{s}^{-2s-2r}\mathbb{K}, \mathcal{O}(s+r+n)).$$

There are natural inclusions

$$\mathsf{Hom}_{S_{s+r}}(\mathbf{s}^{-2s-2r}\mathbb{K}, \mathcal{O}(s+r+n)) \to \mathsf{Hom}_{S_s\times S_r}(\mathbf{s}^{-2s-2r}\mathbb{K}, \mathcal{O}(s+r+n))$$

and they assemble to form a map

$$\mathfrak{D}_{\mathcal{O}} \colon \mathrm{Tw}\,\mathcal{O} \to \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}, \tag{4.5}$$

which is alternatively defined by the equation:

$$\mathfrak{D}_{\mathcal{O}}(f)(1_r \otimes 1_s) = f(1_{s+r}), \qquad f \in \mathrm{Tw}\,\mathcal{O}(n). \tag{4.6}$$

We claim that

Lemma 4.5. *The map $\mathfrak{D}_{\mathcal{O}}$ is a map of operads and the diagram*

$$\begin{tikzcd}
 & \Lambda\mathsf{Lie}_\infty \arrow[dl, "\beta_{\mathcal{O}}"'] \arrow[dr, "\beta_{\mathrm{Tw}\,\mathcal{O}}"] & \\
\mathrm{Tw}\,\mathcal{O} \arrow[rr, "\mathfrak{D}_{\mathcal{O}}"] & & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}
\end{tikzcd} \tag{4.7}$$

commutes.

Proof. Let us consider commutativity of diagram (4.7) first.

By definition of $\beta_{\mathcal{O}}$ (4.2), we have

$$\beta_{\mathcal{O}}(\mathbf{s}1^{\mathfrak{c}}_n)(1_r) = \varphi_{r+n}\,,$$

where $\varphi_n := \hat{\varphi}(\mathbf{s}\,1^{\mathfrak{c}}_n)\,.$

On the other hand,

$$\beta_{\mathrm{Tw}\,\mathcal{O}} = \mathrm{Tw}\,(\beta_{\mathcal{O}}) \circ \mathfrak{T}\,. \tag{4.8}$$

Hence

$$\beta_{\mathrm{Tw}\,\mathcal{O}}(\mathbf{s}\,1^{\mathfrak{c}}_n)(1_r \otimes 1_s) = \beta_{\mathcal{O}}(\mathbf{s}\,1^{\mathfrak{c}}_{r+n})(1_s) = \varphi_{s+r+n}\,. \tag{4.9}$$

Composing $\beta_{\mathcal{O}}$ with $\mathfrak{D}_{\mathcal{O}}$, we get

$$\mathfrak{D}_{\mathcal{O}} \circ \beta_{\mathcal{O}}(\mathbf{s}1^{\mathfrak{c}}_n)(1_r \otimes 1_s) = \beta_{\mathcal{O}}(\mathbf{s}1^{\mathfrak{c}}_n)(1_{r+s}) = \varphi_{s+r+n}\,.$$

Comparing this result with (4.9) we conclude that diagram (4.7) indeed commutes.

Let us now show that $\mathfrak{D}_{\mathcal{O}}$ is a map of operads.

For this purpose we consider a pair of vectors $f \in \mathrm{Tw}\,\mathcal{O}(n)$ and $g \in \mathrm{Tw}\,\mathcal{O}(m)$ and compute

$$\mathfrak{D}_{\mathcal{O}}(f \circ_i g)(1_r \otimes 1_s) \,=\, f \circ_i g(1_{r+s}) \,=\, \sum_{p=0}^{r+s-p} \sum_{\sigma \in \mathrm{Sh}_{p,r+s-p}} \sigma \circ \varrho_{r+s,p,i}\big(f(1_p) \circ_{p+i} g(1_{r+s-p})\big) \tag{4.10}$$

where the family of permutations $\{\varrho_{r,p,i}\}_{p \le r}$ is defined in (3.26)$\,.$

On the other hand,

$$\begin{aligned} \Big(\mathfrak{D}_{\mathcal{O}}(f) \circ_i \mathfrak{D}_{\mathcal{O}}(g)\Big)(1_r \otimes 1_s) &= \sum_{p_1=0}^{r} \sum_{\tau \in \mathrm{Sh}_{p_1,r-p_1}} \tau \circ \varrho_{r,p_1,i}\big(\mathfrak{D}_{\mathcal{O}}(f)(1_{p_1}) \circ_{p_1+i} \mathfrak{D}_{\mathcal{O}}(g)(1_{r-p_1})\big)\,(1_s) \\ &= \sum_{\substack{0 \le p_1 \le r\\ 0 \le p_2 \le s}} \sum_{\sigma \in T_{r,s,p_1,p_2}} \mu_{\mathbf{t}_\sigma}\Big(\mathfrak{D}_{\mathcal{O}}(f)(1_{p_1} \otimes 1_{p_2}) \otimes \mathfrak{D}_{\mathcal{O}}(g)(1_{r-p_1} \otimes 1_{s-p_2})\Big) \\ &= \sum_{\substack{0 \le p_1 \le r\\ 0 \le p_2 \le s}} \sum_{\sigma \in T_{r,s,p_1,p_2}} \mu_{\mathbf{t}_\sigma}\Big(f(1_{p_1+p_2}) \otimes g(1_{r+s-p_1-p_2})\Big)\,, \end{aligned} \tag{4.11}$$

where $\mathbf{t}_\sigma$ is the labeled planar tree depicted on figure 4.1 and the set T_{r,s,p_1,p_2} consists of shuffles $\sigma \in \mathrm{Sh}_{p_1+p_2,r+s-p_1-p_2}$ satisfying the conditions:

$$\begin{gathered} \sigma(1), \sigma(2), \dots, \sigma(p_2) \;\in\; \{1,2,\dots,s\}\,, \\ \sigma(p_2+p_1+1), \sigma(p_2+p_1+2), \dots, \sigma(p_1+s) \;\in\; \{1,2,\dots,s\}\,, \\ \sigma(p_2+1), \sigma(p_2+2), \dots, \sigma(p_2+p_1) \;\in\; \{s+1,s+2,\dots,s+r\}\,, \\ \sigma(p_1+s+1), \sigma(p_1+s+2), \dots, \sigma(s+r) \;\in\; \{s+1,s+2,\dots,s+r\}\,. \end{gathered}$$



FIG. 4.1. The labeled planar tree $\mathbf{t}_\sigma$

It is clear that for every $0 \le p \le r+s$ the (disjoint) union

$$\bigsqcup_{p_1+p_2=p} T_{r,s,p_1,p_2}$$

coincides with the set $\mathrm{Sh}_{p,r+s-p}$.

Hence,

$$\sum_{\substack{0\le p_1\le r\\ 0\le p_2\le s}} \sum_{\sigma\in T_{r,s,p_1,p_2}} \mu_{\mathbf{t}_\sigma}\Big(f(1_{p_1+p_2})\otimes g(1_{r+s-p_1-p_2})\Big) = \sum_{p=0}^{r+s-p} \sum_{\sigma\in \mathrm{Sh}_{p,r+s-p}} \sigma\circ\varrho_{r+s,p,i}\big(f(1_p)\circ_{p+i} g(1_{r+s-p})\big)\,.$$

Thus,

$$\mathfrak{D}_{\mathcal{O}}(f\circ_i g) = \mathfrak{D}_{\mathcal{O}}(f)\ \circ_i\ \mathfrak{D}_{\mathcal{O}}(g)\,.$$

In other words, $\mathfrak{D}_{\mathcal{O}}$ is compatible with all elementary operadic insertions.

It is easy to see that $\mathfrak{D}_{\mathcal{O}}$ is compatible with the units.

Lemma 4.5 is proved. □

Lemma 4.6. *The maps* $\mathfrak{D}_{\mathcal{O}}$ (4.6) *assemble to form a natural transformation* $\mathfrak{D} : \mathrm{Tw} \Rightarrow \mathrm{Tw}\circ\mathrm{Tw}$.

Proof. We have to show that for all arrows $f : \mathcal{O}\to\mathcal{O}'$ (respecting the maps from $\Lambda\mathsf{Lie}_\infty$) the following diagram commutes.

$$\begin{tikzcd}
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,f"] \arrow[d, "\mathfrak{D}_{\mathcal{O}}"'] & \mathrm{Tw}\,\mathcal{O}' \arrow[d, "\mathfrak{D}_{\mathcal{O}'}"] \\
\mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,\mathrm{Tw}\,f"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}'
\end{tikzcd}$$

Unravelling the definitions this is amounts to saying that the following diagrams commute

$$\begin{tikzcd}
\mathsf{Hom}_{S_{r+s}}(\mathbf{s}^{-2r-2s}\mathbb{K}, \mathcal{O}(r+s+n)) \arrow[r, "f\circ"] \arrow[d, hook] & \mathsf{Hom}_{S_{r+s}}(\mathbf{s}^{-2r-2s}\mathbb{K}, \mathcal{O}'(r+s+n)) \arrow[d, hook] \\
\mathsf{Hom}_{S_r\times S_s}(\mathbf{s}^{-2r-2s}\mathbb{K}, \mathcal{O}(r+s+n)) \arrow[r, "f\circ"] & \mathsf{Hom}_{S_r\times S_s}(\mathbf{s}^{-2r-2s}\mathbb{K}, \mathcal{O}'(r+s+n))
\end{tikzcd}$$

for all r, s, n. This is clear. □

Theorem 4.7. *The functor* Tw *together with the natural transformations* $\eta, \mathfrak{D}$ (4.4), (4.6)*, is a comonad on the under-category* $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads.*

Proof. We have to verify the defining relations for a comonad. The two co-unit relations boil down to the statement that for any operad $\mathcal{O}$ the compositions

$$\mathrm{Tw}\,\mathcal{O} \overset{\mathfrak{D}_{\mathcal{O}}}{\longrightarrow} \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} \overset{\eta_{\mathrm{Tw}\,\mathcal{O}}}{\longrightarrow} \mathrm{Tw}\,\mathcal{O}$$

$$\mathrm{Tw}\,\mathcal{O} \overset{\mathfrak{D}_{\mathcal{O}}}{\longrightarrow} \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} \overset{\mathrm{Tw}\,\eta_{\mathcal{O}}}{\longrightarrow} \mathrm{Tw}\,\mathcal{O}$$

are the identity maps on $\mathrm{Tw}\,\mathcal{O}$. This statement follows immediately from the definitions. Next consider the co-associativity axiom. In our case it boils down to the statement that for any operad $\mathcal{O}$ the diagram

$$\begin{tikzcd}
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathfrak{D}_{\mathcal{O}}"] \arrow[d, "\mathfrak{D}_{\mathcal{O}}"'] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} \arrow[d, "\mathfrak{D}_{\mathrm{Tw}\,\mathcal{O}}"] \\
\mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,(\mathfrak{D}_{\mathcal{O}})"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}
\end{tikzcd}$$

commutes. Unravelling the definitions, we have to show that the following diagram commutes

$$\begin{tikzcd}
\mathsf{Hom}_{S_{r+s+t}}(\mathbf{s}^{-2r-2s-2t}\mathbb{K}, \mathcal{O}(r+s+t+n)) \arrow[r, hook] \arrow[d, hook] & \mathsf{Hom}_{S_{r+s}\times S_t}(\mathbf{s}^{-2r-2s-2t}\mathbb{K}, \mathcal{O}(r+s+t+n)) \arrow[d, hook] \\
\mathsf{Hom}_{S_r\times S_{s+t}}(\mathbf{s}^{-2r-2s-2t}\mathbb{K}, \mathcal{O}(r+s+t+n)) \arrow[r, hook] & \mathsf{Hom}_{S_r\times S_s\times S_t}(\mathbf{s}^{-2r-2s-2t}\mathbb{K}, \mathcal{O}(r+s+t+n))
\end{tikzcd}$$

for all r, s, t, n. This is again clear. □

4.2. **Coalgebras over the comonad** Tw**.** Let us now consider coalgebras over the comonad Tw. These are arrows of operads $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ together with an operad map

$$c \colon \mathcal{O} \to \mathrm{Tw}\,\mathcal{O}$$

such that the following axioms hold:

- The following diagram commutes:

$$\begin{tikzcd}
 & \Lambda\mathsf{Lie}_\infty \arrow[dl] \arrow[dr] & \\
\mathcal{O} \arrow[rr, "c"] & & \mathrm{Tw}\,\mathcal{O}
\end{tikzcd} \tag{4.12}$$

- The composition

$$\mathcal{O} \overset{c}{\to} \mathrm{Tw}\,\mathcal{O} \overset{\eta_{\mathcal{O}}}{\to} \mathcal{O} \tag{4.13}$$

  is the identity.
- The following diagram commutes:

$$\begin{tikzcd}
\mathcal{O} \arrow[r, "c"] \arrow[d, "c"] & \mathrm{Tw}\,\mathcal{O} \arrow[d, "\mathfrak{D}_{\mathcal{O}}"] \\
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\, c"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}
\end{tikzcd} \tag{4.14}$$

Remark 4.8. *It often happens that a map from the dg operad* $\Lambda\mathsf{Lie}_\infty$ *to* $\mathcal{O}$ *is clear from the context. In this case, we abuse the notation and say that* $\mathcal{O}$ *is a* Tw *-coalgebra if the corresponding arrow* $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ *carries a coalgebra structure over the comonad* Tw *.*

Example 4.9. *In Subsection 5.1.1 below, we will show that the operads* $\Lambda\mathsf{Lie}$ *and* Ger *carry canonical structures of a* Tw *-coalgebra. Furthermore, it is not hard to see that, the canonical morphism of dg operads*

$$\mathfrak{T} : \Lambda\mathsf{Lie}_\infty \to \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$$

from Lemma 3.10 equips $\Lambda\mathsf{Lie}_\infty$ *with a structure of a* Tw *-coalgebra.*

Example 4.10. *If* $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ *is an arrow, then* $\mathrm{Tw}\,\mathcal{O}$ *is a* Tw *-coalgebra (a cofree* Tw *-coalgebra).*

Another example of a Tw-coalgebra is given in the next section.

4.3. Ger_∞ **is canonically a** Tw**-coalgebra.** Let us recall from [14], [15], [16] that

$$\mathsf{Ger}_\infty := \mathrm{Cobar}(\mathsf{Ger}^\vee)\,,$$

where $\mathsf{Ger}^\vee$ is the cooperad Koszul dual to Ger. Concretely, $\mathsf{Ger}^\vee$ is obtained from $\Lambda^{-2}\mathsf{Ger}$ by taking the linear dual. The operad $\Lambda\mathsf{Lie}_\infty$ is a sub-operad of Ger_∞, and we denote the inclusion by

$$\iota : \Lambda\mathsf{Lie}_\infty \to \mathsf{Ger}_\infty.$$

Our goal, in this section, is to show that Ger_∞ is a Tw-coalgebra. In order to do this, we have to complete two tasks:

1. Construct a map of dg operads $c : \mathsf{Ger}_\infty \to \mathrm{Tw}\,\mathsf{Ger}_\infty$.
2. Verify that c satisfies the axioms for a Tw-coalgebra.

To complete the first task we consider the free $\Lambda^{-2}\mathsf{Ger}$-algebra $\Lambda^{-2}\mathsf{Ger}_n$ in n dummy variables $b_1, \dots, b_n$ of degree zero. The n-th space $\Lambda^{-2}\mathsf{Ger}(n)$ of the operad $\Lambda^{-2}\mathsf{Ger}$ is spanned by $\Lambda^{-2}\mathsf{Ger}$-monomials in $b_1, \dots, b_n$ in which each dummy variable b_i appears exactly once. It is clear that the operad $\Lambda^{-2}\mathsf{Ger}$ is generated by the two vectors $b_1 b_2, \{b_1, b_2\} \in \Lambda^{-2}\mathsf{Ger}(2)$ of degrees 2 and 1, respectively.

Next, we consider the ordered partitions of the set $\{1, 2, \dots, n\}$

$$\{i_{11}, i_{12}, \dots, i_{1p_1}\} \sqcup \{i_{21}, i_{22}, \dots, i_{2p_2}\} \sqcup \dots \sqcup \{i_{t1}, i_{t2}, \dots, i_{tp_t}\} \tag{4.15}$$

satisfying the following properties:

- for each $1 \le \beta \le t$ the index $i_{\beta p_\beta}$ is the biggest among $i_{\beta 1}, \dots, i_{\beta p_\beta}$
- $i_{1p_1} < i_{2p_2} < \dots < i_{tp_t}$ (in particular, $i_{tp_t} = n$).

It is not hard to see that for each $n \ge 1$ the monomials

$$\{b_{i_{11}}, \dots, \{b_{i_{1(p_1-1)}}, b_{i_{1p_1}}\}..\} \dots \{b_{i_{t1}}, \dots, \{b_{i_{t(p_t-1)}}, b_{i_{tp_t}}\}..\} \tag{4.16}$$

corresponding to all ordered partitions (4.15) satisfying the above properties form a basis[8] of $\Lambda^{-2}\mathsf{Ger}(n)$. We denote by I_n the set of the ordered partitions (4.15) and reserve the notation

$$\big\{w_{n,i}\big\}_{i \in I_n} \tag{4.17}$$

and

$$\{w^*_{n,i}\}_{i \in I_n} \tag{4.18}$$

for the basis of $\Lambda^{-2}\mathsf{Ger}(n)$ formed by monomials (4.16) and the dual basis of $\mathsf{Ger}^\vee(n) = \big(\Lambda^{-2}\mathsf{Ger}(n)\big)^*$, respectively.

We observe that, for every $r \ge 0$ and for every basis vector $w_{n,i}$, the monomial

$$b_1 \dots b_r \, w_{n,i}(b_{r+1}, \dots, b_{r+n})$$

belongs to the basis of $\Lambda^{-2}\mathsf{Ger}(r+n)$. In particular, we denote by

$$\big(b_1 \dots b_r \, w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^*$$

the basis vector of $\mathsf{Ger}^\vee(r+n)$ which is dual to $b_1 \dots b_r \, w_{n,i}(b_{r+1}, \dots, b_{r+n})$ in $\Lambda^{-2}\mathsf{Ger}(r+n)$.

Since for every $r \ge 0$ and $n \ge 2$ the element $\mathbf{s}^{2r+1}\big(b_1 \dots b_r \, w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^*$ can be viewed as a vector in

$$\mathbf{s}^{2r+1}\big(\mathsf{Ger}^\vee(r+n)\big)^{S_r} \subset \mathrm{Tw}\,\mathsf{Ger}_\infty(n)\,,$$

the formula

$$\alpha_{\mathfrak{G}} = \sum_{n \ge 2,\, r \ge 0} \; \sum_{i \in I_n} \mathbf{s}^{2r+1}\left(b_1 \dots b_r \, w_{n,i}(b_{r+1}, \dots, b_{r+n})\right)^* \otimes w_{n,i} \tag{4.19}$$

---

[8] Using this fact, it is easy to see that $\dim \Lambda^{-2}\mathsf{Ger}(n) = n!$

defines a degree 1 element in the dg Lie algebra

$$\mathrm{Conv}(\mathsf{Ger}^\vee_\circ, \mathrm{Tw}\,\mathsf{Ger}_\infty) \cong \prod_{n\geq 2}\Big(\mathrm{Tw}\,\mathsf{Ger}_\infty(n)\otimes\Lambda^{-2}\mathsf{Ger}(n)\Big)^{S_n}. \tag{4.20}$$

We claim that

Proposition 4.11. *Equation* (4.19) *defines a Maurer-Cartan element of the dg Lie algebra* (4.20).

The proof of this proposition is quite technical so we postpone it to Subsection 4.3.1 given below.

Due to Theorem 2.6, maps from Ger_∞ to $\mathrm{Tw}\,\mathsf{Ger}_\infty$ are in bijection with Maurer-Cartan elements of the dg Lie algebra (4.20). Hence the Maurer-Cartan element $\alpha_{\mathfrak{G}}$ (4.19) defines a map of dg operads

$$c : \mathsf{Ger}_\infty \to \mathrm{Tw}\,\mathsf{Ger}_\infty\,. \tag{4.21}$$

To prove that c equips Ger_∞ with the structure of a Tw-coalgebra, we have to verify three conditions. The first condition states that the following diagram shall commute:

$$\begin{tikzcd}
& \Lambda\mathsf{Lie}_\infty \arrow[dl] \arrow[dr] & \\
\mathsf{Ger}_\infty \arrow[rr, "c"] & & \mathrm{Tw}\,\mathsf{Ger}_\infty
\end{tikzcd}.$$

Here the right hand arrow is defined as the composition

$$\Lambda\mathsf{Lie}_\infty \to \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty \xrightarrow{\mathrm{Tw}\,\iota} \mathrm{Tw}\,\mathsf{Ger}_\infty.$$

It sends the generator $\mathbf{s}\,1^{\mathfrak{c}}_n$ of $\Lambda\mathsf{Lie}_\infty$ to (cf. (3.45))

$$\sum_{r\geq 0}\mathbf{s}^{2r+1}(b_1b_2\cdots b_{r+n})^*.$$

On the other hand, the left and bottom morphisms in the diagram under consideration send $\mathbf{s}\,1^{\mathfrak{c}}_n$ to

$$c\circ\iota(\mathbf{s}\,1^{\mathfrak{c}}_n) = c\big(\mathbf{s}(b_1\cdots b_n)^*\big) = \sum_{r\geq 0}\mathbf{s}^{2r+1}(b_1b_2\cdots b_{r+n})^*\,.$$

Hence this condition is satisfied. The second condition for Tw-coalgebras to be checked says that the composition

$$\mathsf{Ger}_\infty \to \mathrm{Tw}\,\mathsf{Ger}_\infty \to \mathsf{Ger}_\infty$$

shall be the identity. It is obviously satisfied. The third condition states that the two compositions below shall be the same.

$$\mathsf{Ger}_\infty \xrightarrow{c} \mathrm{Tw}\,\mathsf{Ger}_\infty \xrightarrow{\mathfrak{D}_{\mathsf{Ger}_\infty}} \mathrm{Tw}\,\mathrm{Tw}\,\mathsf{Ger}_\infty$$

$$\mathsf{Ger}_\infty \xrightarrow{c} \mathrm{Tw}\,\mathsf{Ger}_\infty \xrightarrow{\mathrm{Tw}\,(c)} \mathrm{Tw}\,\mathrm{Tw}\,\mathsf{Ger}_\infty$$

In fact, unfolding the definitions one finds that both compositions operate on generators $\mathbf{s}w^*_{n,i}$ of Ger_∞ as follows:

$$\mathbf{s}\, w^*_{n,i} \;\mapsto\; \sum_{r\ge 0}\sum_{s\ge 0} \mathbf{s}^{2r+2s+1} b_1 \dots b_{r+s} w^*_{n,i}(b_{r+s+1}, \dots, b_{r+s+n})$$

Thus the third condition is also satisfied.

Summarizing, we obtain the following theorem:

Theorem 4.12. *The morphism of dg operads*

$$c : \mathsf{Ger}_\infty \to \mathrm{Tw}\,\mathsf{Ger}_\infty \tag{4.22}$$

corresponding to the Maurer-Cartan element $\alpha_{\mathfrak{G}}$ (4.19) *equips* Ger_∞ *with a* Tw *-coalgebra structure.* □

4.3.1. *The proof of Proposition 4.11.* The proof of Theorem 4.12 is based on Proposition 4.11. Here we prove this proposition.

Let n, r be a pair of integers with $r \ge 0$ and $n \ge 2$. In the groupoid $\mathsf{Tree}_2(r+n)$ we consider the objects $\mathbf{t}^{below}_\sigma$, $\mathbf{t}^{above}_\sigma$, $\mathbf{t}^{k}_\sigma$, $\mathbf{t}^{p,q}_{\sigma,\tau}$ shown on figures 4.2, 4.3, 4.4, and 4.5, respectively.



FIG. 4.2. The tree $\mathbf{t}^{below}_\sigma$. Here $2 \le p \le r$ and $\sigma \in \mathrm{Sh}_{p,r-p}$



FIG. 4.3. The tree $\mathbf{t}^{above}_\sigma$. Here $0 \le p \le r-1$ and $\sigma \in \mathrm{Sh}_{p,r-p}$



FIG. 4.4. The tree $\mathbf{t}^{k}_\sigma$. Here $2-n \le p \le r-1$, $1 \le k \le n$, and $\sigma \in \mathrm{Sh}_{p,r-p}$

It is clear that the trees $\mathbf{t}^{below}_\sigma$, $\mathbf{t}^{above}_\sigma$, $\mathbf{t}^{k}_\sigma$, $\mathbf{t}^{p,q}_{\sigma,\tau}$ are all mutually non-isomorphic. Furthermore, if both nodal vertices of a tree $\mathbf{t} \in \mathsf{Tree}_2(n)$ have valencies ≥ 3, then $\mathbf{t}$ is isomorphic to one of the trees in the list: $\mathbf{t}^{below}_\sigma$, $\mathbf{t}^{above}_\sigma$, $\mathbf{t}^{k}_\sigma$, $\mathbf{t}^{p,q}_{\sigma,\tau}$.

We will need the following technical statement:



FIG. 4.5. The tree $\mathbf{t}^{p,q}_{\sigma,\tau}$. Here $0 \le p \le r$, $2 \le q \le n-1$, $\sigma \in \mathrm{Sh}_{p,r-p}$ and $\tau \in \mathrm{Sh}_{q,n-q}$

Lemma 4.13. *Let n, r be a pair of integers with $r \ge 0$, $n \ge 2$, $w_{n,i}$ be a vector in the basis (4.17), and*

$$\big(b_1 \dots b_r\, w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^*$$

be the basis vector of $\mathsf{Ger}^\vee(r+n)$ dual to $b_1 \dots b_r\, w_{n,i}(b_{r+1}, \dots, b_{r+n}) \in \Lambda^{-2}\mathsf{Ger}(r+n)$. Then

$$\partial^{\mathrm{Cobar}} \mathbf{s}\, (b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n}))^* = \tag{4.23}$$

$$- \sum_{\substack{2 \le p \le r \\ \sigma \in \mathrm{Sh}_{p,r-p}}} (-1)^{|w_{n,i}|} \big(\mathbf{t}^{below}_\sigma ; \mathbf{s}\, (b_1 \dots b_{r-p+1} w_{n,i}(b_{r-p+2}, \dots, b_{r-p+1+n}))^* \otimes \mathbf{s}\, (b_1 \dots b_p)^*\big)$$

$$- \sum_{\substack{0 \le p \le r-1 \\ \sigma \in \mathrm{Sh}_{p,r-p}}} \big(\mathbf{t}^{above}_\sigma ; \mathbf{s}\, (b_1 \dots b_{r-p+1})^* \otimes \mathbf{s}\, (b_1 \dots b_p w_{n,i}(b_{p+1}, \dots, b_{p+n}))^*\big)$$

$$- \sum_{\substack{2-n \le p \le r-1 \\ \sigma \in \mathrm{Sh}_{p,r-p}}} \sum_{k=1}^{n} (-1)^{|w_{n,i}|} \big(\mathbf{t}^{k}_\sigma ; \mathbf{s}\, (b_1 \dots b_p w_{n,i}(b_{p+1}, \dots, b_{p+n}))^* \otimes \mathbf{s}\, (b_1 \dots b_{r-p+1})^*\big)$$

$$- \sum_{\substack{0 \le p \le r \\ 2 \le q \le n-1}} \sum_{\substack{\sigma \in \mathrm{Sh}_{p,r-p} \\ \tau \in \mathrm{Sh}_{q,n-q}}} \sum_{i_1, i_2} (-1)^{|w_{n-q+1,i_1}||w_{q,i_2}| + |w_{n-q+1,i_1}|} f^i_{i_1 i_2} \big(\mathbf{t}^{p,q}_{\sigma,\tau} \, ; \, \mathbf{s}\, (b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1}))^*$$

$$\otimes\, \mathbf{s}\, (b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q}))^*\big)\,,$$

where the coefficients $f^i_{i_1 i_2} \in \mathbb{K}$ are defined by the equation

$$\tau(w_{n-q+1,i_1} \circ_1 w_{q,i_2}) = \sum_{i \in I_n} f^i_{i_1 i_2} w_{n,i}\,. \tag{4.24}$$

Proof. The statement of the lemma follows from these equations:

$$\Delta_{\mathbf{t}^{below}_\sigma} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* = \tag{4.25}$$

$$(b_1 \dots b_{r-p+1} w_{n,i}(b_{r-p+2}, \dots, b_{r-p+1+n}))^* \otimes (b_1 \dots b_p)^*\,,$$

$$\Delta_{\mathbf{t}^{above}_\sigma} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* = \tag{4.26}$$

$$(b_1 \dots b_{r-p+1})^* \otimes (b_1 \dots b_p w_{n,i}(b_{p+1}, \dots, b_{p+n}))^*\,,$$

$$\Delta_{\mathbf{t}^{k}_\sigma} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* = \tag{4.27}$$

$$(b_1 \dots b_p w_{n,i}(b_{p+1}, \dots, b_{p+n}))^* \otimes (b_1 \dots b_{r-p+1})^*\,,$$

and

$$\Delta_{\mathbf{t}^{p,q}_{\sigma,\tau}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* = \tag{4.28}$$

$$\sum_{\substack{i_1 \in I_{n-q+1},\\ i_2 \in I_q}} (-1)^{|w_{n-q+1,i_1}| \cdot |w_{q,i_2}|} f^{i}_{i_1 i_2} (b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1}))^*$$

$$\otimes\, (b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q}))^*\,,$$

where the coefficients $f^{i}_{i_1 i_2}$ are defined by equation (4.24).

Let us prove that equation (4.28) holds.

In general, we have

$$\Delta_{\mathbf{t}^{p,q}_{\sigma,\tau}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* =$$

$$\sum_{\substack{j_1 \in I_{n+p-q+1},\\ j_2 \in I_{r-p+q}}} g^{i}_{j_1, j_2} w^*_{n+p-q+1, j_1} \otimes w^*_{r-p+q, j_2}\,, \qquad g^{i}_{j_1 j_2} \in \mathbb{K}\,. \tag{4.29}$$

It is not hard to see that, if the basis vector $w_{n+p-q+1,j_1}$ is not of the form

$$b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1})$$

or the basis vector w_{r-p+q,j_2} is not of the form

$$b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q})\,,$$

then

$$\big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(\mu_{\mathbf{t}^{p,q}_{\sigma,\tau}}(w_{n+p-q+1,j_1} \otimes w_{r-p+q,j_2}\Big) = 0\,.$$

In other words, $g^{i}_{j_1 j_2} = 0$ unless the basis vector $w_{n+p-q+1,j_1}$ is of the form

$$b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1})$$

and the basis vector w_{r-p+q,j_2} is of the form

$$b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q})\,.$$

Hence,

$$\Delta_{\mathbf{t}^{p,q}_{\sigma,\tau}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* = \tag{4.30}$$

$$\sum_{\substack{i_1 \in I_{n-q+1},\\ i_2 \in I_q}} \widetilde{f}^{i}_{i_1 i_2} (b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1}))^* \otimes (b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q}))^*$$

for some coefficients $\widetilde{f}^{i}_{i_1 i_2} \in \mathbb{K}$.

On the other hand, we have

$$\Delta_{\mathbf{t}^{p,q}_{\sigma,\tau}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1})$$

$$\otimes\, b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q})\Big) =$$

$$\begin{gathered}\big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(\mu_{\mathbf{t}^{p,q}_{\sigma,\tau}} \big(b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1}) \\ \otimes\, b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q})\big)\Big) = \\ (-1)^{\varepsilon_{i_1 i_2}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(b_{\sigma(1)} \dots b_{\sigma(p)} \\ w_{n-q+1,i_1}\big(b_{\sigma(p+1)} \dots b_{\sigma(r)} w_{q,i_2}(b_{r+\tau(1)}, \dots, b_{r+\tau(q)}), b_{r+\tau(q+1)}, \dots, b_{r+\tau(n)}\big)\,,\Big)\end{gathered} \tag{4.31}$$

where the sign factor $(-1)^{\varepsilon_{i_1 i_2}}$ comes from permuting the $\Lambda^{-2}\mathsf{Ger}$-monomial w_{q,i_2} with brackets $\{\ ,\ \}$.

Since in the variables $b_1, \dots, b_r$ in the $\Lambda^{-2}\mathsf{Ger}$-monomial $b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})$ only enter $\Lambda^{-1}\mathsf{Lie}$ words of length 1, we have

$$\begin{gathered}\Delta_{\mathbf{t}^{p,q}_{\sigma,\tau}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1}) \\ \otimes\, b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q})\Big) =\end{gathered}$$

$$\begin{gathered}(-1)^{\varepsilon_{i_1 i_2}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(b_{\sigma(1)} \dots b_{\sigma(r)} \\ w_{n-q+1,i_1}\big(w_{q,i_2}(b_{r+\tau(1)}, \dots, b_{r+\tau(q)}), b_{r+\tau(q+1)}, \dots, b_{r+\tau(n)}\big)\Big) = \\ (-1)^{\varepsilon_{i_1 i_2}} \big(b_1 \dots b_r w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^* \Big(b_1 \dots b_r \\ w_{n-q+1,i_1}\big(w_{q,i_2}(b_{r+\tau(1)}, \dots, b_{r+\tau(q)}), b_{r+\tau(q+1)}, \dots, b_{r+\tau(n)}\big)\Big) = f^i_{i_1 i_2}\,.\end{gathered} \tag{4.32}$$

Thus $\widetilde{f}^i_{i_1 i_2} = (-1)^{|w_{n-q+1,i_1}||w_{q,i_2}|} f^i_{i_1 i_2}$ and equation (4.28) holds.

Using the identification $\mathsf{Ger}^\vee = \Lambda^{-2}\mathsf{Ger}^*$ in the similar way, it is easy to prove that equations (4.25), (4.26), (4.27) also hold.

Lemma 4.13 follows. □

To prove Proposition 4.11 we need to show that the element $\alpha_{\mathfrak{G}}$ (4.19) satisfies the Maurer-Cartan equation

$$\partial^{\mathrm{Cobar}} \alpha_{\mathfrak{G}} + \partial^{\mathrm{Tw}} \alpha_{\mathfrak{G}} + \alpha_{\mathfrak{G}} \bullet \alpha_{\mathfrak{G}} = 0 \tag{4.33}$$

in the dg Lie algebra (4.20).

Equation (4.33) unfolds as follows:

$$\begin{gathered}\sum_{n\ge 2,\, r\ge 0}\ \sum_{i\in I_n} \mathbf{s}^{2r}\, \partial^{\mathrm{Cobar}}\Big(\mathbf{s}\, \big(b_1 \dots b_r\, w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^*\Big) \otimes w_{n,i} + \\ \sum_{n\ge 2,\, r\ge 0}\ \sum_{i\in I_n} \partial^{\mathrm{Tw}}\ \Big(\mathbf{s}^{2r+1}\, \big(b_1 \dots b_r\, w_{n,i}(b_{r+1}, \dots, b_{r+n})\big)^*\Big) \otimes w_{n,i} \\ + \sum_{\substack{0\le p\le r \\ 2\le q\le n-1}}\ \sum_{\substack{\sigma\in \mathrm{Sh}_{p,r-p} \\ \tau\in\mathrm{Sh}_{q,n-q}}}\ \sum_{i_1,i_2} (-1)^{|w_{n-q+1,i_1}|(|w_{q,i_2}|+1)}\, \big(\mathbf{t}^{p,q}_{\sigma,\tau};\, \mathbf{s}^{2p+1}\, (b_1 \dots b_p w_{n-q+1,i_1}(b_{p+1}, \dots, b_{p+n-q+1}))^* \\ \otimes\, \mathbf{s}^{2(r-p)+1}\, (b_1 \dots b_{r-p} w_{q,i_2}(b_{r-p+1}, \dots, b_{r-p+q}))^*\big)\ \otimes\ f^i_{i_1 i_2} w_{n,i} = 0\,,\end{gathered} \tag{4.34}$$

where the coefficients $f^i_{i_1 i_2}$ are defined by equations (4.24).

Let us now use Lemma 4.13.

The contribution to

$$\sum_{n\geq 2,\, r\geq 0} \sum_{i\in I_n} \mathbf{s}^{2r}\, \partial^{\mathrm{Cobar}}\Big(\mathbf{s}\,\big(b_1\dots b_r\, w_{n,i}(b_{r+1},\dots,b_{r+n})\big)^*\Big)\otimes w_{n,i}$$

coming from the last sum in the right hand side of (4.23) cancels with the third sum in equation (4.34).

The contributions to

$$\sum_{n\geq 2,\, r\geq 0} \sum_{i\in I_n} \mathbf{s}^{2r}\, \partial^{\mathrm{Cobar}}\Big(\mathbf{s}\,\big(b_1\dots b_r\, w_{n,i}(b_{r+1},\dots,b_{r+n})\big)^*\Big)\otimes w_{n,i}$$

coming from the remaining sums in the right hand side of (4.23) cancel the sum

$$\sum_{n\geq 2,\, r\geq 0} \sum_{i\in I_n} \partial^{\mathrm{Tw}}\,\Big(\mathbf{s}^{2r}\,\big(b_1\dots b_r\, w_{n,i}(b_{r+1},\dots,b_{r+n})\big)^*\Big)\otimes w_{n,i}\,.$$

Thus $\alpha_{\mathfrak{G}}$ satisfies (4.33) and Proposition 4.11 follows.

□

5. HOMOTOPY THEORETIC PROPERTIES OF THE TWISTING PROCEDURE

Let us prove that the functor Tw preserves quasi-isomorphisms.

Theorem 5.1. *Let* $\Lambda\mathsf{Lie}_\infty \to \mathcal{O} \xrightarrow{F} \mathcal{O}'$ *be a morphism of the under-category* $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads with* F *being a quasi-isomorphism. Then, the map* $\mathrm{Tw}\,(F) : \mathrm{Tw}\,\mathcal{O} \to \mathrm{Tw}\,\mathcal{O}'$ *is also a quasi-isomorphism.*

Proof. We want to show that $\mathrm{Tw}\,\mathcal{O}(n) \to \mathrm{Tw}\,\mathcal{O}'(n)$ is a quasi-isomorphism for every $n = 0, 1, 2, \dots$. This is equivalent to saying that the mapping cone

$$C := \mathrm{Tw}\,\mathcal{O}(n) \oplus \mathbf{s}\,\mathrm{Tw}\,\mathcal{O}'(n)$$

is acyclic for every n. There is a natural complete filtration $C = \mathcal{F}_0 \supset \mathcal{F}_1 \supset \cdots$ such that

$$\mathcal{F}_p = \prod_{r\geq p} \mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathcal{O}(r+n)) \oplus \prod_{r\geq p} \mathbf{s}\,\mathsf{Hom}_{S_r}(\mathbf{s}^{-2r}\mathbb{K}, \mathcal{O}'(r+n))\,.$$

Note that the associated graded complex for this filtration is

$$\bigoplus_p \mathcal{F}_p/\mathcal{F}_{p+1} \cong \bigoplus_p \mathsf{Hom}_{S_p}(\mathbf{s}^{-2p}\mathbb{K}, \mathcal{O}(p+n)\oplus \mathbf{s}\,\mathcal{O}'(p+n)).$$

The complex $\mathcal{O}(p+n)\oplus \mathbf{s}\,\mathcal{O}'(p+n)$ is the mapping cone of $\mathcal{O}(p+n)\to\mathcal{O}'(p+n)$ and hence acyclic by assumption. Since taking invariants with respect to a finite group action commutes with taking cohomology, we conclude that the associated graded is acyclic as well. Thus, by Lemma E.1 from Appendix E, the statement of the proposition follows. □

Example 5.2. *Since the canonical maps*

$$U_{\Lambda\mathsf{Lie}} : \Lambda\mathsf{Lie}_\infty \to \Lambda\mathsf{Lie}\,, \qquad U_{\mathsf{Ger}} : \mathsf{Ger}_\infty \to \mathsf{Ger}$$

are quasi-isomorphisms of dg operads, the morphisms

$$\mathrm{Tw}\,(U_{\Lambda\mathsf{Lie}}) : \mathrm{Tw}\,(\Lambda\mathsf{Lie}_\infty) \to \mathrm{Tw}\,(\Lambda\mathsf{Lie})\,,$$

and

$$\mathrm{Tw}\,(U_{\mathsf{Ger}}) : \mathrm{Tw}\,\mathsf{Ger}_\infty \to \mathrm{Tw}\,\mathsf{Ger}$$

are also quasi-isomorphisms of dg operads.

As we will see below in Section 5.1, many objects of the under-category $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads* satisfy the following remarkable property:

Definition 5.3. An arrow $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$ is called a *homotopy fixed point* for Tw if the counit map $\eta_{\mathcal{O}} : \mathrm{Tw}\,\mathcal{O} \to \mathcal{O}$ is a quasi-isomorphism.

If the map from $\Lambda\mathsf{Lie}_\infty$ is clear from the context, we will say, by abusing the notation, that $\mathcal{O}$ is a homotopy fixed point of Tw.

Theorem 5.1 implies that, if a dg operad $\mathcal{O}$ is a homotopy fixed point for Tw and a dg operad $\mathcal{O}'$ is quasi-isomorphic to $\mathcal{O}$, then $\mathcal{O}'$ is also a homotopy fixed point for Tw .

Example 5.4. *In Subsection 5.1.1 below, we will show that the operads* $\Lambda\mathsf{Lie}$ *and* Ger *are homotopy fixed points for* Tw . *Hence, by Theorem 5.1, the dg operads* $\Lambda\mathsf{Lie}_\infty$ *and* Ger_∞ *are also homotopy fixed points for* Tw .

Example 5.5. *In Section 9.3 below, we will show that the dg operad* Br *governing brace algebras (see Section 9) is a homotopy fixed point for* Tw*.*

5.1. **The distributive law and the functor Tw.** In this section, we describe a large class of dg operads which are simultaneously Tw-coalgebras and homotopy fixed points for Tw .

Let P be an arbitrary dg operad. Furthermore, let $\mathcal{O}$ be the collection of cochain complexes

$$\mathcal{O} = P \odot \Lambda\mathsf{Lie} \tag{5.1}$$

which is obtained by computing the plethysm (2.12) of P with $\Lambda\mathsf{Lie}$.

Let $\varsigma_{1,i}$ be the cycle $(1, 2, \dots, i)$ in S_{n+1}. It is not hard to see that imposing the relation

$$\{a_1, a_2\} \circ_2 \gamma = (-1)^{|\gamma|} \sum_{i=1}^{n} \varsigma_{1,i}(\gamma \circ_i \{a_1, a_2\}) \qquad \forall\ \gamma \in P(n) \tag{5.2}$$

and using the operad structures on P and on $\Lambda\mathsf{Lie}$ we get a natural dg operad structure on the collection (5.1). Following [21] and [19, Section 8.6], we say that the dg operad $\mathcal{O}$ is obtained from P and $\Lambda\mathsf{Lie}$ using the *distributive law* (5.2).

Example 5.6. *The operad* Ger *is obtained from the operad* Com *via the above construction.*

$$\mathsf{Ger} = \mathsf{Com} \odot \Lambda\mathsf{Lie}\,.$$

For dg operads obtained in this way, we have the following straightforward proposition:

Proposition 5.7. *Let $\mathcal{O}$ be the dg operad which is obtained via taking the plethysm* (5.1) *and imposing relation* (5.2)*. Then for every $\gamma \in \mathcal{O}(n)$ we have*

$$\{a_1, a_2\} \circ_2 \gamma = (-1)^{|\gamma|} \sum_{i=1}^{n} \varsigma_{1,i}(\gamma \circ_i \{a_1, a_2\}) \,. \tag{5.3}$$

Furthermore, for every $\mathcal{O}$-algebra V, the adjoint action $\{v, ~\}$ is a derivation of the $\mathcal{O}$-algebra structure on V for every $v \in V$. □

Remark 5.8. *Proposition 5.7 implies that the operad* BV *governing the Batalin-Vilkovisky (BV) algebra* [12] *is not an operad which is obtained via the above construction. Indeed, the unary operation δ on a BV algebra V satisfies the relation*

$$\delta\big(\{v_1, v_2\}\big) + \{\delta(v_1), v_2\} + (-1)^{|v_1|}\{v_1, \delta(v_2)\} = 0 \,.$$

Hence, in general, the adjoint action $\{v, ~\}$ is not a derivation of the BV algebra structure on V.

Let $\mathcal{O}$ be the dg operad which is obtained via taking the plethysm (5.1) and imposing relation (5.2). It is obvious that the dg operad $\mathcal{O}$ receives a natural embedding

$$\mathfrak{i} : \Lambda\mathsf{Lie} \hookrightarrow \mathcal{O} \,. \tag{5.4}$$

Composing $\mathfrak{i}$ with the canonical quasi-isomorphism (2.59) $U_{\Lambda\mathsf{Lie}} : \Lambda\mathsf{Lie}_{\infty} \to \Lambda\mathsf{Lie}$ we get a map of dg operads

$$\hat{\varphi} := \mathfrak{i} \circ U_{\Lambda\mathsf{Lie}} ~:~ \Lambda\mathsf{Lie}_{\infty} ~\to~ \mathcal{O} \,. \tag{5.5}$$

Hence, we may apply the twisting procedure to the pair $(\mathcal{O}, \hat{\varphi})$ and obtain a dg operad $\mathrm{Tw}\,\mathcal{O}$.

According to Section 3.4, the spaces

$$\mathrm{Tw}^{\oplus}\mathcal{O}(n) = \bigoplus \mathbf{s}^{2r}\big(\mathcal{O}(r+n)\big)^{S_r} \tag{5.6}$$

form a sub- dg operad of $\mathrm{Tw}\,\mathcal{O}$.

It turns our that the dg operad $\mathrm{Tw}^{\oplus}\mathcal{O}$ coincides with $\mathrm{Tw}\,\mathcal{O}$, provided the dg operad P satisfies a minor technical condition. Namely,

Proposition 5.9. *Let P be a dg operad for which there exists a integer N such that for each $n \geq 0$ and for every $v \in P(n)$*

$$|v| \geq N \,. \tag{5.7}$$

If the dg operad $\mathcal{O}$ is obtained via taking the plethysm (5.1) *and imposing relation* (5.2) *then*

$$\mathrm{Tw}^{\oplus}\mathcal{O} = \mathrm{Tw}\,\mathcal{O} \,. \tag{5.8}$$

Proof. Let m be an integer and n be a non-negative integer. Our goal is to prove every sum

$$\sum_{r=0}^{\infty} w_r \,, \qquad w_r \in \mathbf{s}^{2r}\big(\mathcal{O}(r+n)\big)^{S_r} \tag{5.9}$$

of a fixed degree m has only finitely many terms.

For every $r \ge 0$, the graded vector space $\mathbf{s}^{2r}\big(\mathcal{O}(r+n)\big)^{S_r}$ is spanned by vectors of the form

$$\sum_{\sigma \in S_r} \sigma(v; X_1, X_2, \dots, X_t)\,, \tag{5.10}$$

where $v \in P(t)$ for some t and X_i are vectors in $\Lambda\mathsf{Lie}(k_i)$ such that

$$\sum_{i=1}^{t} k_i = n + r\,. \tag{5.11}$$

Since a vector (5.10) carries degree m in

$$\mathbf{s}^{2r}\big(\mathcal{O}(r+n)\big)^{S_r}\,,$$

we obtain the following equation

$$m = 2r + |v| + \sum_{i=1}^{t}(1 - k_i)\,. \tag{5.12}$$

Combining (5.11) with (5.12) we get

$$m = r + |v| + t - n \tag{5.13}$$

and hence

$$r = m + n - t - |v|\,. \tag{5.14}$$

Since $t \ge 0$ and $|v| \ge N$, we deduce that

$$r \le m + n - N\,. \tag{5.15}$$

Thus the sum in (5.9) has indeed only finitely many terms. □

The following theorem is the central result of this section[9].

Theorem 5.10. *If an object* $(\mathcal{O}, \hat{\varphi})$ *of the under-category* $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads is obtained via taking the plethysm* (5.1) *of a dg operad* P *with* $\Lambda\mathsf{Lie}$ *and imposing relation* (5.2)*, then* $(\mathcal{O}, \hat{\varphi})$ *is canonically a* Tw *-coalgebra. If, in addition, the dg operad* P *satisfies the condition of Proposition 5.9, then* $\mathcal{O}$ *is a homotopy fixed point for* Tw *.*

Proof. Identity (5.3) implies that the canonical embedding

$$\mathrm{emb}_{\mathcal{O}} : \mathcal{O} \to \mathrm{Tw}\,\mathcal{O}$$

$$\big(\mathrm{emb}_{\mathcal{O}}(v)\big)(1_r) = \begin{cases} v & \text{if } r = 0\,, \\ 0 & \text{otherwise} \end{cases} \tag{5.16}$$

---

[9] An idea of this proof is borrowed from [28].

is compatible with the differentials ∂^{Tw} and $\partial^{\mathcal{O}}$. Furthermore, it is easy to see that the diagrams

$$\begin{tikzcd}
\mathcal{O} \arrow[r, "\mathrm{emb}_{\mathcal{O}}"] \arrow[dr, "\mathrm{id}"] & \mathrm{Tw}\,\mathcal{O} \arrow[d, "\eta_{\mathcal{O}}"] \\
 & \mathcal{O}
\end{tikzcd}
\qquad
\begin{tikzcd}
\mathcal{O} \arrow[r, "\mathrm{emb}_{\mathcal{O}}"] \arrow[d, "\mathrm{emb}_{\mathcal{O}}"'] & \mathrm{Tw}\,\mathcal{O} \arrow[d, "\mathrm{Tw}\,(\eta_{\mathcal{O}})"] \\
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathfrak{D}_{\mathcal{O}}"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}
\end{tikzcd}$$

commute. Thus $(\mathcal{O}, \hat{\varphi})$ is indeed a coalgebra over the comonad Tw .

Let us now observe that the map $\mathrm{emb}_{\mathcal{O}}$ (5.16) lands into the sub- dg operad $\mathrm{Tw}^{\oplus}\mathcal{O}$. Furthermore, if the dg operad P satisfies the condition of Proposition 5.9, then

$$\mathrm{Tw}^{\oplus}\mathcal{O} = \mathrm{Tw}\,\mathcal{O}\,.$$

Thus, we need to prove that the map

$$\mathrm{emb}_{\mathcal{O}}\Big|_{\mathcal{O}(n)} : \mathcal{O}(n) \hookrightarrow \mathrm{Tw}^{\oplus}\mathcal{O}(n) \tag{5.17}$$

is a quasi-isomorphism of cochain complexes for every n. For this purpose we consider the free $\mathcal{O}$-algebra

$$\mathcal{O}(a, a_1, a_2, \dots, a_n) \tag{5.18}$$

generated by n dummy variables $a_1, a_2, \dots, a_n$ of degree zero and one dummy variable a of degree 2. Next, we denote by δ the degree 1 derivation of the $\mathcal{O}$-algebra $\mathcal{O}(a, a_1, a_2, \dots, a_n)$ defined by the formulas:

$$\delta(a) = \frac{1}{2}\{a, a\}\,, \qquad \delta(a_i) = 0 \qquad \forall\ 1 \le i \le n\,. \tag{5.19}$$

Due to the Jacobi identity, we have $\delta^2 = 0$. Hence, δ is a differential on $\mathcal{O}$-algebra (5.18). We will combine δ with the differential $\partial^{\mathcal{O}}$ coming from $\mathcal{O}$ and consider the $\mathcal{O}$-algebra (5.18) with the differential

$$\partial^{\mathcal{O}} + \delta\,. \tag{5.20}$$

Let us denote by

$$\mathcal{O}'(a, a_1, a_2, \dots, a_n) \tag{5.21}$$

the subcomplex of (5.18) which is spanned by $\mathcal{O}$-monomials in which each variable from the set $\{a_1, a_2, \dots, a_n\}$ appears exactly once.

It is not hard to see that the formula

$$\psi(\mathbf{s}^{2r}\, v) = \frac{1}{r!}\, v \otimes a^{\otimes\, r} \otimes a_1 \otimes a_2 \otimes \dots \otimes a_n\,, \qquad v \in \big(\mathcal{O}(r+n)\big)^{S_r} \tag{5.22}$$

gives us an obvious isomorphism of graded vector spaces

$$\psi : \mathrm{Tw}^{\oplus}\mathcal{O}(n) = \bigoplus_{r=0}^{\infty} \big(\mathcal{O}(r+n)\big)^{S_r} \ \to\ \mathcal{O}'(a, a_1, a_2, \dots, a_n)\,. \tag{5.23}$$

Using identity (5.3), one can show that ψ intertwines the differentials ∂^{Tw} on $\mathrm{Tw}^{\oplus}\mathcal{O}(n)$ and (5.20) on (5.21). Hence, ψ (5.23) is an isomorphism of cochain complexes.

Let us denote by

$$\mathcal{O}'(a_1, a_2, \dots, a_n) \tag{5.24}$$

the subcomplex of the free $\mathcal{O}$-algebra $\mathcal{O}(a_1, a_2, \dots, a_n)$ which is spanned by monomials in which each variable from the set $\{a_1, a_2, \dots, a_n\}$ appears exactly once.

We observe that the assignment

$$v \mapsto v \otimes a_1 \otimes a_2 \otimes \dots \otimes a_n\,, \qquad v \in \mathcal{O}(n)$$

gives us an obvious identification between the cochain complex $\mathcal{O}(n)$ and (5.24).

We also observe that the composition $\psi \circ \mathrm{emb}_{\mathcal{O}}$ operates by the formula

$$\psi \circ \mathrm{emb}_{\mathcal{O}}(v) = v \otimes a_1 \otimes a_2 \otimes \dots \otimes a_n\,. \tag{5.25}$$

Thus, our goal is to show that the embedding

$$\mathcal{O}'(a_1, a_2, \dots, a_n) \hookrightarrow \mathcal{O}'(a, a_1, a_2, \dots, a_n) \tag{5.26}$$

is a quasi-isomorphism of cochain complexes.

For this purpose we introduce the free $\Lambda\mathsf{Lie}$-algebra

$$\Lambda\mathsf{Lie}(a, a_1, a_2, \dots, a_n) \tag{5.27}$$

generated by n dummy variables $a_1, a_2, \dots, a_n$ of degree zero and one dummy variable a of degree 2.

We consider the $\Lambda\mathsf{Lie}$-algebra $\Lambda\mathsf{Lie}(a, a_1, a_2, \dots, a_n)$ with the differential δ defined in (5.19).

In addition, we introduce two subspaces

$$\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n) \subset \Lambda\mathsf{Lie}(a, a_1, a_2, \dots, a_n) \tag{5.28}$$

and

$$\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n) \subset \Lambda\mathsf{Lie}(a, a_1, a_2, \dots, a_n)\,. \tag{5.29}$$

Here $\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)$ is spanned by $\Lambda\mathsf{Lie}$-monomials in $\Lambda\mathsf{Lie}(a, a_1, a_2, \dots, a_n)$ which involve each variable from the set $\{a_1, a_2, \dots, a_n\}$ at most once, and $\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)$ is spanned by $\Lambda\mathsf{Lie}$-monomials in $\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)$ which do not involve the variable a at all.

It is clear that both subspaces (5.28) and (5.29) are subcomplexes of (5.27). Moreover, the restriction of the differential δ to (5.28) is zero.

The proof of Theorem 5.10 is based on the following statement:

Lemma 5.11. *The embedding*

$$\mathrm{emb} : \Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n) \hookrightarrow \Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n) \tag{5.30}$$

is quasi-isomorphism of cochain complexes. In other words, for every cocycle $c \in \Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)$, *there exists a vector* $c_1 \in \Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)$ *such that*

$$c - \delta(c_1) \in \Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\,.$$

The proof of this lemma is somewhat technical so we give it in Subsection 5.1.2 below. To deduce the desired statement from Lemma (5.11), we observe that the cone of the embedding (5.26) is a direct summand in the cone of the embedding

$$P\big(\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\big) \hookrightarrow P\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big)\,. \tag{5.31}$$

Hence the map (5.26) is a quasi-isomorphism if so is the map (5.31).

To prove that the embedding (5.31) is a quasi-isomorphism we consider the following increasing filtration on the cochain complex $P\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big)$:

$$\dots \subset \mathcal{F}^m P\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big) \subset \mathcal{F}^{m+1} P\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big) \subset \dots\,, \tag{5.32}$$

where

$$\mathcal{F}^m P\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)$$

is spanned by $\mathcal{O}$-monomials w for which

$$\deg_a(w) - |w| \le m\,,$$

with $\deg_a(w)$ being the degree of w in a. Since the differential $\partial^{\mathcal{O}}$ does not change the degree in a and the differential δ raises it by 1 the total differential $\partial^{\mathcal{O}} + \delta$ is compatible with the filtration (5.32). Restricting (5.32) to the subcomplex $P\big(\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\big)$ we get the "silly" filtration

$$\mathcal{F}^m P\big(\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\big)^k = \begin{cases} P\big(\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\big)^k & \text{if } k \ge -m\,, \\ \mathbf{0} \quad \text{otherwise} \end{cases} \tag{5.33}$$

with the zero differential on the associated graded complex.

To describe the associated graded complex for (5.32) we denote by $\widetilde{P}$ the operad in $\mathsf{grVect}_{\mathbb{K}}$ which is obtained from P by forgetting the differential. It is clear from the construction that the associated graded complex

$$\mathrm{Gr}\, P\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big)$$

is isomorphic to the cochain complex

$$\widetilde{P}\big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big) \tag{5.34}$$

with the differential δ. Using the Künneth theorem and the fact that cohomology commutes with taking coinvariants, we deduce from Lemma (5.11) that the map

$$\Big(\widetilde{P}(n) \otimes \big(\Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\big)\Big)_{S_n} \to \Big(\widetilde{P}(n) \otimes \big(\Lambda\mathsf{Lie}'(a, a_1, a_2, \dots, a_n)\big)\Big)_{S_n}$$

is a quasi-isomorphism of cochain complexes for every n. Thus the embedding (5.31) induces a quasi-isomorphism on the level of associated graded complexes. Combining this observation with the fact that the filtrations (5.32) and (5.33) are locally bounded and cocomplete, we deduce, from

Lemma A.3 in [8, Appendix A], that the embedding (5.31) is also a quasi-isomorphism. Hence the map (5.26) is a quasi-isomorphism of cochain complexes and the theorem is proved. □

5.1.1. *The operads* $\Lambda\mathsf{Lie}$*,* Ger*,* $\Lambda\mathsf{Lie}_\infty$*, and* Ger_∞ *are homotopy fixed points for* Tw *.* Theorem 5.10 has the following useful Corollary:

Corollary 5.12. *The operads* $\Lambda\mathsf{Lie}$ *and* Ger *carry a canonical* Tw *-coalgebra structure. Moreover, these operads are homotopy fixed points for* Tw *.* □

Combining Corollary 5.12 with Theorem 5.1, we immediately conclude that[10]

Corollary 5.13. *The dg operads* $\Lambda\mathsf{Lie}_\infty$ *and* Ger_∞ *are homotopy fixed points for* Tw *.* □

Remark 5.14. *Let us recall that, due to Lemma 3.10 and Theorem 4.12, both dg operads* $\Lambda\mathsf{Lie}_\infty$ *and* Ger_∞ *carry canonical* Tw *-coalgebra structures. In fact, the canonical map of operads*

$$\mathfrak{T} : \Lambda\mathsf{Lie}_\infty \to \mathrm{Tw}\,\Lambda\mathsf{Lie}_\infty$$

was used to introduce the notion of Tw *-coalgebra.*

5.1.2. *Proof of Lemma 5.11.* The proof of Theorem 5.10 was based on Lemma 5.11. Here we give a proof of this lemma. We consider a non-empty ordered subset $\{i_1 < i_2 < \dots < i_k\}$ of $\{1, 2, \dots, n\}$ and denote by

$$\Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k}) \tag{5.35}$$

the subcomplex of $\Lambda\mathsf{Lie}'(a, a_1, \dots, a_n)$ which is spanned by $\Lambda\mathsf{Lie}$-monomials in $\Lambda\mathsf{Lie}(a, a_{i_1}, \dots, a_{i_k})$ involving each variable in the set $\{a_{i_1}, \dots, a_{i_k}\}$ exactly once.

It is clear that $\Lambda\mathsf{Lie}'(a, a_1, \dots, a_n)$ splits into the direct sum of subcomplexes:

$$\Lambda\mathsf{Lie}'(a, a_1, \dots, a_n) = \mathbb{K}\langle a, \{a, a\}\rangle \ \oplus \bigoplus_{\{i_1 < i_2 < \dots < i_k\}} \Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k})\,, \tag{5.36}$$

where the summation runs over all non-empty ordered subsets $\{i_1 < i_2 < \dots < i_k\}$ of $\{1, 2, \dots, n\}$.

It is not hard to see that the subcomplex $\mathbb{K}\langle a, \{a, a\}\rangle$ is acyclic. Thus our goal is to show that every cocycle in $\Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k})$ is cohomologous to cocycle in the intersection

$$\Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k}) \cap \Lambda\mathsf{Lie}'(a_1, a_2, \dots, a_n)\,.$$

To prove this fact we consider the tensor algebra

$$T\big(\mathbb{K}\langle \mathbf{s}^{-1} a, \mathbf{s}^{-1} a_{i_1}, \mathbf{s}^{-1} a_{i_2}, \dots, \mathbf{s}^{-1} a_{i_{k-1}}\rangle\big) \tag{5.37}$$

in the variables $\mathbf{s}^{-1} a, \mathbf{s}^{-1} a_{i_1}, \mathbf{s}^{-1} a_{i_2}, \dots, \mathbf{s}^{-1} a_{i_{k-1}}$ and denote by

$$T'(\mathbf{s}^{-1} a, \mathbf{s}^{-1} a_{i_1}, \mathbf{s}^{-1} a_{i_2}, \dots, \mathbf{s}^{-1} a_{i_{k-1}}) \tag{5.38}$$

the subspace of (5.37) which is spanned by monomials involving each variable from the set

$$\{\mathbf{s}^{-1} a_{i_1}, \mathbf{s}^{-1} a_{i_2}, \dots, \mathbf{s}^{-1} a_{i_{k-1}}\}$$

---

[10] The same result for the operad Lie_∞ was obtained in [5, Section 7] by J. Chuang and A. Lazarev.

exactly once.

It is not hard to see that the formula

$$\nu(x_{j_1} \otimes x_{j_2} \otimes \dots \otimes x_{j_N}) = \{\mathbf{s}\, x_{j_1}, \{\mathbf{s}\, x_{j_2}, \{\dots \{\mathbf{s}\, x_{j_N}, a_{i_k}\}..\} \tag{5.39}$$

defines an isomorphism of the graded vector spaces

$$\nu : T'(\mathbf{s}^{-1}\, a, \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}) \xrightarrow{\cong} \Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k})\,.$$

Let us denote by δ_T a degree 1 derivation of the tensor algebra (5.37) defined by the equations

$$\delta_T(\mathbf{s}^{-1}\, a_{i_t}) = 0\,, \qquad \delta_T(\mathbf{s}^{-1}\, a) = \mathbf{s}^{-1}\, a \otimes \mathbf{s}^{-1}\, a\,. \tag{5.40}$$

It is not hard to see that $(\delta_T)^2 = 0$. Thus, δ_T is a differential on the tensor algebra (5.37).

The subspace (5.38) is obviously a subcomplex of (5.37). Furthermore, using the following consequence of Jacobi identity

$$\{a, \{a, X\}\} = -\frac{1}{2}\{\{a, a\}, X\}\,, \qquad \forall\; X \in \Lambda\mathsf{Lie}(a, a_1, \dots, a_n),$$

it is easy to show that

$$\delta \circ \nu = \nu \circ \delta_T\,.$$

Thus ν is an isomorphism from the cochain complex

$$\big(T'(\mathbf{s}^{-1}\, a, \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}), \delta_T\big)$$

to the cochain complex

$$\big(\Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k}), \delta\big)\,.$$

To compute cohomology of the cochain complex

$$\Big(T\big(\mathbb{K}\langle \mathbf{s}^{-1}\, a, \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}\rangle\big), \delta_T\Big) \tag{5.41}$$

we observe that the truncated tensor algebra

$$\underline{T}_{\mathbf{s}^{-1}\, a} := \underline{T}\big(\mathbb{K}\langle \mathbf{s}^{-1}\, a\rangle\big) \tag{5.42}$$

form an acyclic subcomplex of (5.41).

We also observe that the cochain complex (5.41) splits into the direct sum of subcomplexes

$$T\big(\mathbb{K}\langle \mathbf{s}^{-1}\, a, \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}\rangle\big) = T\big(\mathbb{K}\langle \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}\rangle\big) \oplus \tag{5.43}$$

$$\bigoplus_{m \ge 2,\, p_1, \dots, p_m} V_{a_\bullet}^{\otimes\, p_1} \otimes \underline{T}_{\mathbf{s}^{-1}\, a} \otimes V_{a_\bullet}^{\otimes\, p_2} \otimes \underline{T}_{\mathbf{s}^{-1}\, a} \otimes \dots \otimes V_{a_\bullet}^{\otimes\, p_{m-1}} \otimes \underline{T}_{\mathbf{s}^{-1}\, a} \otimes V_{a_\bullet}^{\otimes\, p_m}\,,$$

where $V_{a_\bullet}$ is the cochain complex

$$V_{a_\bullet} := \mathbb{K}\langle \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}\rangle$$

with the zero differential and the summation runs over all combinations $(p_1, \dots, p_m)$ of integers satisfying the conditions

$$p_1, p_m \ge 0, \qquad p_2, \dots, p_{m-1} \ge 1\,.$$

By Künneth's theorem all the subcomplexes

$$V_{a_\bullet}^{\otimes\, p_1} \otimes \underline{T}_{\mathbf{s}^{-1}\, a} \otimes V_{a_\bullet}^{\otimes\, p_2} \otimes \underline{T}_{\mathbf{s}^{-1}\, a} \otimes \dots \otimes V_{a_\bullet}^{\otimes\, p_{m-1}} \otimes \underline{T}_{\mathbf{s}^{-1}\, a} \otimes V_{a_\bullet}^{\otimes\, p_m}$$

are acyclic. Hence for every cocycle c in (5.41) there exists a vector c_1 in (5.41) such that

$$c - \delta_T(c_1) \in T\big(\mathbb{K}\langle \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}} \rangle\big)\,.$$

Combining this observation with the fact that the subcomplex (5.38) is a direct summand in (5.41), we conclude that, for every cocycle c in (5.38) there exists a vector c_1 in (5.38) such that

$$c - \delta_T(c_1) \in T'(\mathbf{s}^{-1}\, a, \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}}) \ \cap\ T\big(\mathbb{K}\langle \mathbf{s}^{-1}\, a_{i_1}, \mathbf{s}^{-1}\, a_{i_2}, \dots, \mathbf{s}^{-1}\, a_{i_{k-1}} \rangle\big)\,.$$

Since the map ν (5.39) is an isomorphism from the cochain complex (5.38) with the differential δ_T to the cochain complex (5.35) with the differential δ, we deduce that every cocycle in (5.35) is cohomologous to a unique cocycle in the intersection

$$\Lambda\mathsf{Lie}''(a, a_{i_1}, \dots, a_{i_k}) \cap \Lambda\mathsf{Lie}'(a_1, \dots, a_n)\,.$$

Therefore every cocycle in $\Lambda\mathsf{Lie}'(a, a_1, \dots, a_n)$ is cohomologous to a unique cocycle in the subcomplex

$$\Lambda\mathsf{Lie}'(a_1, \dots, a_n)\,.$$

Lemma 5.11 is proved. $\square$

5.1.3. *The operads* As *and* As_∞ *are* Tw *-coalgebras and are homotopy fixed points for* Tw *.* Recall that As is the operad which governs associative algebras without unit.

We have the obvious map of operads

$$\mathsf{Lie} \to \mathsf{As}$$

and hence the maps of operads

$$\mathfrak{i} : \Lambda\mathsf{Lie} \to \Lambda\mathsf{As}\,, \tag{5.44}$$

and

$$\Lambda\mathsf{Lie}_\infty \to \Lambda\mathsf{As}\,. \tag{5.45}$$

In other words, the operad $\Lambda\mathsf{As}$ is naturally an object of the under-category $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads*.

Let us show that

Proposition 5.15. *The operad* $\Lambda\mathsf{As}$ *is naturally a* Tw *-coalgebra. Furthermore,* $\Lambda\mathsf{As}$ *is a homotopy fixed point for the functor* Tw *.*

Remark 5.16. *Before we proceed to the proof, we should remark that the associativity law for* $\Lambda\mathsf{As}$*-algebras has a "funny" sign factor. Namely, a* $\Lambda\mathsf{As}$*-algebra structure on a graded vector space* V *is a degree* -1 *binary operation* $\cdot$ *on* V *satisfying the associativity condition*

$$(v_1 \cdot v_2) \cdot v_3 = -(-1)^{|v_1|} v_1 \cdot (v_2 \cdot v_3)\,.$$

Proof. Although the operad $\Lambda\mathsf{As}$ is not obtained[11] via taking a plethysm of $\Lambda\mathsf{Lie}$ with another operad, it is not hard to see that for every vector $v \in \Lambda\mathsf{As}(n)$

$$\mathfrak{i}(\{a_1, a_2\}) \circ_2 v = (-1)^{|v|} \sum_{i=1}^{n} \varsigma_{1,i}\big(v \circ_i \mathfrak{i}(\{a_1, a_2\})\big)\,, \tag{5.46}$$

where $\varsigma_{1,i}$ is the cycle $(1, 2, \dots, i)$.

Using this identity, it is easy to see that the canonical embedding

$$\mathrm{emb} : \Lambda\mathsf{As} \to \mathrm{Tw}\ \Lambda\mathsf{As}$$

$$\big(\mathrm{emb}(v)\big)(1_r) = \begin{cases} v & \text{if } r = 0\,, \\ 0 & \text{otherwise} \end{cases} \tag{5.47}$$

is compatible with the differentials, i.e.

$$\partial^{\mathrm{Tw}}\big(\mathrm{emb}(v)\big) = 0\,, \qquad \forall\ v \in \Lambda\mathsf{As}(n)\,.$$

A straightforward verification shows that the map emb (5.47) satisfies the axioms of the Tw-coalgebra.

To show that $\Lambda\mathsf{As}$ is a homotopy fixed point for Tw, we consider the free $\Lambda\mathsf{As}$-algebra $\Lambda\mathsf{As}(a, a_1, \dots, a_n)$ in n dummy variables $a_1, \dots, a_n$ of degree 0 and one dummy variable a of degree 2.

The algebra $\Lambda\mathsf{As}(a, a_1, \dots, a_n)$ carries the differential δ defined by the formulas:

$$\delta(a) = a \cdot a\,, \qquad \delta(a_i) = 0\,, \qquad \forall\ 1 \le i \le n\,. \tag{5.48}$$

Next, we denote by $\Lambda\mathsf{As}'(a, a_1, \dots, a_n)$ the subcomplex of $\Lambda\mathsf{As}(a, a_1, \dots, a_n)$ which is spanned by $\Lambda\mathsf{As}$-monomials in which each variable from the set $\{a_1, \dots, a_n\}$ appears exactly once. It is not hard to see that the cochain complexes

$$\mathrm{Tw}\ \Lambda\mathsf{As}(n) \qquad \text{and} \qquad \Lambda\mathsf{As}'(a, a_1, \dots, a_n)$$

are isomorphic and, moreover, the natural embedding

$$\Lambda\mathsf{As}(n) \hookrightarrow \Lambda\mathsf{As}'(a, a_1, \dots, a_n)$$

induces an isomorphism on the level of cohomology.

This observation implies that the embedding emb (5.47) is a quasi-isomorphism of dg operads. Hence $\Lambda\mathsf{As}$ is indeed a homotopy fixed point for the functor Tw. □

According to Section 3.5, we may ask the same questions about the operad As keeping in mind the canonical map $\mathsf{Lie} \to \mathsf{As}$. For this case, Proposition 5.15 gives us the following obvious corollary:

Corollary 5.17. *The operad* As *is naturally a* Tw*-coalgebra. Furthermore,* As *is a homotopy fixed point for the functor* Tw. □

---

[11] Instead, $\Lambda\mathsf{As}$ carries a natural filtration, such that $\mathrm{Gr}\ \Lambda\mathsf{As} \cong \Lambda\mathsf{Com} \odot \Lambda\mathsf{Lie}$.

Let us denote by As_∞ the dg operad which governs A_∞-algebras, i.e.

$$\mathsf{As}_\infty = \mathrm{Cobar}(\Lambda \mathsf{coAs})\,. \tag{5.49}$$

We claim that

Corollary 5.18. *The dg operad* As_∞ *is naturally a* Tw *-coalgebra. Furthermore,* As_∞ *is a homotopy fixed point for the functor* Tw *.*

Proof. The second claim is an obvious consequence of Theorem 5.1 and Corollary 5.17. So it remains to prove the first claim.

Because of sign factor, it is more convenient to prove that the dg operad

$$\Lambda^{-1}\mathsf{As}_\infty = \mathrm{Cobar}(\mathsf{coAs}) \tag{5.50}$$

is a Tw-coalgebra. Then the desired statement will follow from the arguments of Section 3.5.

Our goal is to produce a map of dg operads

$$c : \mathrm{Cobar}(\mathsf{coAs}) \to \mathrm{Tw}\,\mathrm{Cobar}(\mathsf{coAs}) \tag{5.51}$$

and verify axioms of the coalgebra over Tw .

For this purpose we recall that $\mathsf{As}(n)$ has the canonical basis

$$\Big\{ a_{\sigma(1)} a_{\sigma(2)} \dots a_{\sigma(n)} \Big\}_{\sigma \in S_n}\,, \tag{5.52}$$

where $a_1, a_2, \dots, a_n$ are dummy variables of degree zero.

Next, we denote by $(a_{\sigma(1)} a_{\sigma(2)} \dots a_{\sigma(n)})^*$ vectors of the dual basis in $\mathsf{coAs}(n) = \big(\mathsf{As}(n)\big)^*$ and observe that the formula $(n \ge 2, r \ge 0)$

$$\alpha\big((a_{\sigma(1)} a_{\sigma(2)} \dots a_{\sigma(n)})^*\big)(1_r) = \sum_{\lambda \in \mathrm{Sh}_{r,n}} \sum_{\tau \in S_r} \mathbf{s}\, \lambda \, (a_{\tau(1)} \dots a_{\tau(r)} a_{r+\sigma(1)} a_{r+\sigma(2)} \dots a_{r+\sigma(n)})^* \tag{5.53}$$

defines a degree 1 vector in the dg Lie algebra $\mathrm{Conv}(\mathsf{coAs}_\circ, \mathrm{Tw}\,\mathrm{Cobar}(\mathsf{coAs}))$. Here

$$\mathbf{s}\, \lambda \, (a_{\tau(1)} \dots a_{\tau(r)} a_{r+\sigma(1)} a_{r+\sigma(2)} \dots a_{r+\sigma(n)})^*$$

is viewed as a vector in $\mathbf{s}\,\mathsf{coAs}(r+n) \subset \mathrm{Cobar}(\mathsf{coAs})$.

A direct computation shows that α is a Maurer-Cartan element of the dg Lie algebra

$$\mathrm{Conv}\big(\mathsf{coAs}_\circ, \mathrm{Tw}\,\mathrm{Cobar}(\mathsf{coAs})\big)$$

Hence, by Theorem 2.6, α gives us a map of dg operads (5.51).

It is easy to check that this map satisfies all the axioms of the coalgebra over the comonad Tw . So we leave the verification of these axioms to the reader. □

Remark 5.19. *In Section 7 of* [5]*, J. Chuang and A. Lazarev also proved that* As *and* As_∞ *are homotopy fixed points for the functor* Tw *.*

5.1.4. *Example: a* Tw *-coalgebra which is not a homotopy fixed point for* Tw *.* We constructed a large class of Tw-coalgebras each of which is a homotopy fixed point for Tw . Let us now give an example of a Tw-coalgebra which is not a homotopy fixed point for Tw . Let P be an operad in the category $\mathsf{grVect}_{\mathbb{K}}$ such that $P(n) \neq 0$ for all $n \geq 1$. (For example, the operad As would work.) Let

$$\hat{\varphi} : \Lambda\mathsf{Lie}_{\infty} \to P$$

be the zero map. Then the dg operads $\mathrm{Tw}\, P$ and $\mathrm{Tw}\,(\mathrm{Tw}\, P)$ with the spaces

$$\mathrm{Tw}\, P(n) = \prod_{r \geq 0} \mathbf{s}^{2r} \big(P(r+n) \big)^{S_r} \tag{5.54}$$

and

$$\mathrm{Tw}\,(\mathrm{Tw}\, P)(n) = \prod_{r,s \geq 0} \mathbf{s}^{2r+2s} \big(P(r+s+n) \big)^{S_r \times S_s} \tag{5.55}$$

carry the zero differentials. The operad $\mathrm{Tw}\, P$ is a Tw-coalgebra by construction, however,

Proposition 5.20. *The operad* $\mathrm{Tw}\, P$ *is not a homotopy fixed point for* Tw *.*

Proof. Let r, n, s be a positive integers and let v be a non-zero vector in $P(r+s+n)$. Such a vector exists, since $P(n) \neq 0$ for all $n \geq 1$. The sum

$$\sum_{\sigma \in S_r \times S_s} \mathbf{s}^{2r+2s} \sigma(v) \tag{5.56}$$

can be viewed as a vector in $\mathrm{Tw}\,(\mathrm{Tw}\, P)(n)$. Since the differential on $\mathrm{Tw}\,(\mathrm{Tw}\, P)$ is non-zero, then vector (5.56) is a non-trivial cocycle. On the other hand, since $r \geq 1$, the counit map

$$\eta_{\mathrm{Tw}\, P} : \mathrm{Tw}\,(\mathrm{Tw}\, P) \to \mathrm{Tw}\, P$$

sends vector (5.56) to zero. Thus $\eta_{\mathrm{Tw}\, P}$ is not a quasi-isomorphism. $\square$

6. TWISTING $\mathcal{O}$-ALGEBRA STRUCTURES BY MAURER-CARTAN ELEMENTS

Let $\mathcal{O}$ be an object of the under-category $\Lambda\mathsf{Lie}_{\infty} \downarrow \mathit{Operads}$. Following Section 3.2 we consider the category $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$ whose objects are pairs

$$(V, \alpha)\,,$$

where V is an $\mathcal{O}$-algebra equipped with a complete descending filtration[12] and $\alpha \in \mathcal{F}_1 V^2$ is a Maurer-Cartan element. Morphisms in $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$ are morphisms of filtered $\mathcal{O}$-algebras $f : V \to V'$ which send α to α'.

Recall that Theorem 3.7 yields a functor

$$\mathfrak{F} : \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}} \to \mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}} \tag{6.1}$$

---

[12] As above, we assume that the $\mathcal{O}$-algebra structure on V is compatible with the filtration (3.2).

from the category $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$ to the category $\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}}$ of filtered $\mathrm{Tw}\,\mathcal{O}$-algebras. More precisely, the functor $\mathfrak{F}$ assigns to a pair (V, α) the cochain complex V^{α} with the differential ∂^{α} (3.22) and the $\mathrm{Tw}\,\mathcal{O}$-algebra structure defined by equation (3.23).

If the arrow $\hat{\varphi} : \Lambda\mathsf{Lie}_{\infty} \to \mathcal{O}$ is a Tw-coalgebra then we have a morphism of dg operads

$$c : \mathcal{O} \to \mathrm{Tw}\,\mathcal{O} \tag{6.2}$$

which induces a functor

$$c^* : \mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}} \to \mathrm{Alg}_{\mathcal{O}} \tag{6.3}$$

from the category $\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}}$ to the category $\mathrm{Alg}_{\mathcal{O}}$ of $\mathcal{O}$-algebras.

Composing (6.1) and (6.3) we get the functor

$$\mathrm{ATw} := c^* \circ \mathfrak{F} : \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}} \to \mathrm{Alg}_{\mathcal{O}}\,. \tag{6.4}$$

As a cochain complex $\mathrm{ATw}\,(V, \alpha)$ is V^{α} with the differential ∂^{α} (3.22) and the $\mathcal{O}$-algebra structure is induced by the map $c : \mathcal{O} \to \mathrm{Tw}\,\mathcal{O}$.

For a pair $(V, \alpha) \in \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$, we say that the $\mathcal{O}$-algebra $\mathrm{ATw}\,(V, \alpha)$ is obtained from V via *twisting by a Maurer-Cartan element* α. We refer to the construction of $\mathrm{ATw}\,(V, \alpha)$ as *the twisting procedure*[13] for $\mathcal{O}$-algebras.

Axioms of a coalgebra over the comonad Tw listed in Section 4.2 imply that the twisting procedure satisfies the following properties:

P1 Considered as the $\Lambda\mathsf{Lie}_{\infty}$-algebra, $\mathrm{ATw}\,(V, \alpha)$ is obtained via twisting the $\Lambda\mathsf{Lie}_{\infty}$ algebra V by the Maurer-Cartan element α.

P2 Twisting by the zero Maurer-Cartan element $\alpha = 0$ does not change the $\mathcal{O}$-algebra, i.e., $\mathrm{ATw}\,(V, 0) = V$.

P3 If α is a Maurer-Cartan element of V and α' is a Maurer-Cartan element of $\mathrm{ATw}\,(V, \alpha)$ then the twisted $\mathcal{O}$-algebras

$$\mathrm{ATw}\big(\mathrm{ATw}\,(V, \alpha), \alpha'\big) \qquad \text{and} \qquad \mathrm{ATw}\,(V, \alpha + \alpha')$$

coincide.

Indeed, Property **P1** follows easily from commutativity of diagram (4.12).

Equation (3.23) implies that, if $\alpha = 0$, then the $\mathrm{Tw}\,\mathcal{O}$-algebra structure on $V^{\alpha} = V$ factors through the counit map $\eta_{\mathcal{O}} : \mathrm{Tw}\,\mathcal{O} \to \mathcal{O}$. Thus Property **P2** holds because the composition (4.13) is the identity map on $\mathcal{O}$.

Finally, Property **P3** follows from commutativity of diagram (4.14).

Let us now consider functorial properties of ATw with respect to morphisms in the under-category $\Lambda\mathsf{Lie}_{\infty} \downarrow$ *Operads*.

---

[13] Note that this twisting procedure for $\mathcal{O}$-algebras is defined only if the dg operad $\mathcal{O}$ is a coalgebra over the comonad Tw.

For any arrow $\Psi : \mathcal{O} \to \mathcal{O}'$ which fits into the commutative diagram

$$\begin{tikzcd}
 & \Lambda\mathsf{Lie}_\infty \arrow[dl, "\hat{\varphi}"'] \arrow[dr, "\hat{\varphi}'"] & \\
\mathcal{O} \arrow[rr, "\Psi"] & & \mathcal{O}'
\end{tikzcd} \tag{6.5}$$

we can obviously extend the functor

$$\Psi^* : \mathrm{Alg}_{\mathcal{O}'} \to \mathrm{Alg}_{\mathcal{O}}$$

to

$$\Psi^* : \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}'} \to \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}} \,. \tag{6.6}$$

Furthermore, we observe that the diagram

$$\begin{tikzcd}
\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}'} \arrow[r, "\Psi^*"] \arrow[d, "\mathfrak{F}"] & \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}} \arrow[d, "\mathfrak{F}"] \\
\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}'} \arrow[r, "(\mathrm{Tw}\,\Psi)^*"] & \mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}}
\end{tikzcd} \tag{6.7}$$

commutes because the construction of the functor $\mathfrak{F}$ (3.29) is functorial in $\mathcal{O}$.

If the dg operads $\mathcal{O}$ and $\mathcal{O}'$ are Tw-coalgebras and the map Ψ is compatible with the Tw-coalgebra structures then the diagram

$$\begin{tikzcd}
\mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}'} \arrow[r, "(\mathrm{Tw}\,\Psi)^*"] \arrow[d, "(c')^*"] & \mathrm{Alg}^{\mathrm{filtr}}_{\mathrm{Tw}\,\mathcal{O}} \arrow[d, "c^*"] \\
\mathrm{Alg}_{\mathcal{O}'} \arrow[r, "\Psi^*"] & \mathrm{Alg}_{\mathcal{O}}
\end{tikzcd} \tag{6.8}$$

commutes.

Thus, combining commutative diagrams (6.7) and (6.8) we arrive at the following functorial property of the twisting procedure.

Theorem 6.1. *If Ψ is a map of Tw-coalgebras $\mathcal{O}$ and $\mathcal{O}'$ then the diagram*

$$\begin{tikzcd}
\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}'} \arrow[r, "\Psi^*"] \arrow[d, "\mathrm{ATw}_{\mathcal{O}'}"] & \mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}} \arrow[d, "\mathrm{ATw}_{\mathcal{O}}"] \\
\mathrm{Alg}_{\mathcal{O}'} \arrow[r, "\Psi^*"] & \mathrm{Alg}_{\mathcal{O}}
\end{tikzcd} \tag{6.9}$$

commutes. □

At the end of this section, we observe that, under a mild technical condition, the functor ATw preserves quasi-isomorphisms. More precisely,

Proposition 6.2. *Let* $\mathcal{O}$ *be a dg operad equipped with a* Tw *-coalgebra structure and let* $f : (V, \alpha) \to (V', \alpha')$ *be a morphism in* $\mathrm{Alg}^{\mathrm{MC}}_{\mathcal{O}}$ *. Let us assume that the filtrations on* V *and* V' *are bounded from the left. If* f *is a quasi-isomorphism of the underlying cochain complexes* (V, ∂) *and* (V', ∂') *then* f *is also a quasi-isomorphism from* $\mathrm{ATw}\,(V, \alpha)$ *to* $\mathrm{ATw}\,(V', \alpha')$ *.*

Proof. The cone

$$C = V^{\alpha} \oplus \mathbf{s}(V')^{\alpha'}$$

of the morphism $f : V^{\alpha} \to (V')^{\alpha'}$ carries the obvious descending filtration

$$\mathcal{F}_m C := \mathcal{F}_m V^{\alpha} \,\oplus\, \mathbf{s}\mathcal{F}_m (V')^{\alpha'} .$$

This filtration is complete and bounded from the left.

Furthermore, the associated graded complex

$$\mathrm{Gr}\, C$$

is isomorphic to the cone

$$V \oplus \mathbf{s}V'$$

of the morphism $f : V \to V'$.

Hence $\mathrm{Gr}\, C$ is acyclic and, by Lemma E.1, the cone of the morphism $f : V^{\alpha} \to (V')^{\alpha'}$ is also acyclic.

Proposition 6.2 follows. □

7. The operad of brace trees BT

The remaining four sections of the paper are devoted to an application of the developed machinery to Deligne's conjecture. We will introduce an auxiliary operad BT, define the brace operad Br as a suboperad of $\mathrm{Tw}\,\mathsf{BT}$, and finally, prove Theorem 1.1 stated in the Introduction.

Let us start with defining the operad of brace trees BT.

To define the space $\mathsf{BT}(n)$ we introduce an auxiliary set $\mathcal{T}(n)$. An element of this set is a planar tree T equipped with a bijection between $\{1, 2, \dots, n\}$ and the set

$$V(T) \setminus \{\text{root vertex}\}$$

of non-root vertices. For $n = 0$ the set $\mathcal{T}(n)$ is empty.

We call elements of $\mathcal{T}(n)$ *brace trees*. Examples of brace trees are shown on figures 7.1 and 7.2. On figures, non-root vertices are depicted by white circles with the corresponding numbers inscribed.



Fig. 7.1. A brace tree $T' \in \mathcal{T}(6)$



Fig. 7.2. The brace tree $T_{\mathrm{id}} \in \mathcal{T}(1)$

The n-th space $\mathsf{BT}(n)$ of BT consists of linear combinations of elements in $\mathcal{T}(n)$. The structure of a graded vector space on $\mathsf{BT}(n)$ is obtained by declaring that each non-root edge carries degree -1. In other words, for every $T \in \mathcal{T}(n)$ we have

$$|T| = 1 - |E(T)|\,, \tag{7.1}$$

where $E(T)$ is the set of all edges of T. A simple combinatorics shows that for every brace tree $T \in \mathcal{T}(n)$ we have

$$|T| = 1 - n\,. \tag{7.2}$$

Hence, the graded vector space $\mathsf{BT}(n)$ is concentrated in the single degree $1-n$. (In particular, the operad BT may only carry the zero differential.)

Since $\mathcal{T}(0)$ is empty, we have

$$\mathsf{BT}(0) = \mathbf{0}\,. \tag{7.3}$$

Furthermore, since in $\mathcal{T}(1)$ we have only element T_{id} (see figure 7.2),

$$\mathsf{BT}(1) = \mathbb{K}\,. \tag{7.4}$$

7.1. **The operad structure on BT.** Let $T \in \mathcal{T}(n)$, $T' \in \mathcal{T}(k)$ and $1 \le i \le k$. Our goal is to define the output of the elementary insertion $T' \circ_i T \in \mathsf{BT}(n+k-1)$.

Let v_i be the non-root vertex of T' with label i. If v_i is a leaf (i.e. v_i does not have incoming edges) then the vector $T' \circ_i T \in \mathsf{BT}(n+k-1)$ is, up to sign, represented by a brace tree T'' which is obtained from T' by erasing the vertex v_i and gluing the brace tree T via identifying the root edge of T with the edge originating at v_i. After this operation we relabel elements of the set

$$V(T'') \setminus \{\text{root vertex}\} = \big(V(T) \setminus \{\text{root vertex}\}\big) \sqcup (V(T') \setminus \{v_i\})$$

in the obvious way. Let us now consider the case when v_i has $q \ge 1$ incoming edges. Since T' is a planar tree, these q incoming edges are totally ordered. So we denote them by $e_1, e_2, \dots, e_q$ keeping in mind that

$$e_1 < e_2 < \dots < e_q\,. \tag{7.5}$$

The desired vector $T' \circ_i T \in \mathsf{BT}(n+k-1)$ is represented by the sum

$$T' \circ_i T = \sum_{\alpha} (-1)^{f(\alpha)} T_\alpha \tag{7.6}$$

where T_α is obtained from T' and T following these steps:

- first, we erase the vertex v_i and glue the brace tree T via identifying the root edge T with the edge originating from v_i ,
- second, we attach the edges $e_1, e_2, \dots, e_q$ to vertices in the set

$$V(T) \setminus \{\text{root vertex}\}\,. \tag{7.7}$$

- finally, we relabel elements of the set

  $$V(T'') \setminus \{\text{root vertex}\} = \big(V(T) \setminus \{\text{root vertex}\}\big) \sqcup (V(T') \setminus \{v_i, \text{root vertex}\})$$

  in the obvious way.

Ways of connecting the edges $e_1, e_2, \dots, e_q$ to vertices in the set (7.7) should satisfy the following condition

Condition 7.1. *The restriction of the total order on the set* $E(T_\alpha)$ *of* T_α *to the subset* $\{e_1, e_2, \dots, e_q\}$ *should coincide with the order* (7.5) *.*

This condition can be reformulated in geometric terms as follows. If we choose a small tubular neighborhood of the tree T (drawn on the plane) and walk along its boundary starting from the root vertex in the clockwise direction then we will cross the edges in this order: first, we will cross e_1, second, we will cross e_2, third, we will cross e_3, and so on.

The summation in (7.6) goes over all ways α of connecting the edges $e_1, e_2, \dots, e_q$ to vertices in the set (7.7) satisfying Condition 7.1.

To define the sign factors $(-1)^{f(\alpha)}$ in (7.6) we extend the total orders on the sets $E(T')$ and $(E(T) \setminus \{\text{root edge}\})$ to the disjoint union

$$E(T') \sqcup (E(T) \setminus \{\text{root edge}\}) \tag{7.8}$$

by declaring that all elements of $E(T')$ are smaller than elements of $E(T) \setminus \{\text{root edge}\}$.

Next we observe that the set (7.8) is naturally isomorphic to the set $E(T_\alpha)$ of edges of T_α . On the other hand, the set $E(T_\alpha)$ carries a possibly different total order coming from planar structure on T_α .

So the factor $(-1)^{f(\alpha)}$ is the sign of the permutation which connects these total orders on the set

$$E(T_\alpha) \cong E(T') \sqcup (E(T) \setminus \{\text{root edge}\})\,.$$

Example 7.2. *Let* T' *(resp.* $T_{\circ\circ}$*) be the brace tree depicted on figure 7.1 (resp. figure 7.3). The result of the insertion* $T' \circ_2 T$ *is the sum of brace trees shown on figure 7.4.*



FIG. 7.3. The brace tree $T_{\circ\circ} \in \mathcal{T}(2)$



FIG. 7.4. The vector $T' \circ_2 T \in \mathsf{BT}(7)$

The symmetric group S_n acts on $\mathsf{BT}(n)$ in the obvious way by rearranging the labels. It is not hard to see that operations (7.6) together with this action give us an operad structure on BT with the identity element represented by the brace tree T_{id} depicted on figure 7.2.

8. THE OPERAD Tw BT

The operad BT receives a natural map from $\Lambda\mathsf{Lie}$

$$\varphi : \Lambda\mathsf{Lie} \to \mathsf{BT}\,. \tag{8.1}$$

Since the operad $\Lambda\mathsf{Lie}$ is generated by the binary bracket operation $\{\cdot,\cdot\} \in \Lambda\mathsf{Lie}(2)$, the map (8.1) is uniquely determined by its value $\varphi(\{\cdot,\cdot\})$, which equals

$$\varphi(\{\cdot,\cdot\}) = T_{\circ\circ} + \sigma_{12} T_{\circ\circ}\,, \tag{8.2}$$

where $T_{\circ\circ}$ is the brace tree depicted on figure 7.3 and σ_{12} is the transposition in S_2. The desired compatibility with the Jacobi relation

$$\varphi(\{\cdot,\cdot\}) \circ_1 \varphi(\{\cdot,\cdot\}) + \text{cyclic permutations}\,(1,2,3) = 0 \tag{8.3}$$

can be checked by a straightforward computation.

In this section we give an explicit description for the twisted version $\mathrm{Tw}\,\mathsf{BT}$ of BT corresponding to the map (8.2). As a graded vector space,

$$\mathrm{Tw}\,\mathsf{BT}(n) = \prod_{r=0}^{\infty} \mathbf{s}^{2r} \big(\mathsf{BT}(r+n)\big)^{S_r}\,. \tag{8.4}$$

Using the observation that for every m the space $\mathsf{BT}(m)$ is concentrated in the single degree $1-m$ we conclude that the subspace $\mathrm{Tw}\,\mathsf{BT}^p(n)$ of degree p vectors in $\mathrm{Tw}\,\mathsf{BT}(n)$ is spanned by vectors of the form

$$\sum_{\sigma \in S_r} \sigma(T) \tag{8.5}$$

where T is an arbitrary brace tree in $\mathcal{T}(r+n)$ and $r = p+n-1$. In particular,

$$\mathrm{Tw}\,\mathsf{BT} = \mathrm{Tw}^{\oplus}\mathsf{BT} \tag{8.6}$$

(cf. Section 3.4).

To represent vectors (8.5), it is convenient to extend the set $\mathcal{T}(n)$ to another auxiliary set $\mathcal{T}^{\mathrm{tw}}(n)$. An element of $\mathcal{T}^{\mathrm{tw}}(n)$ is a planar tree T equipped with the following data:

- a partition of the set $V(T)$ of vertices
$$V(T) = V_{\mathrm{lab}}(T) \sqcup V_{\nu}(T) \sqcup V_{root}(T)$$
into the singleton $V_{root}(T)$ consisting of the root vertex, the set $V_{\mathrm{lab}}(T)$ consisting of n vertices, and the set $V_{\nu}(T)$ consisting of vertices which we call *neutral*;
- a bijection between the set $V_{\mathrm{lab}}(T)$ and the set $\{1, 2, \dots, n\}$;

We also call elements of $\mathcal{T}^{\mathrm{tw}}(n)$ brace trees. Figures 7.3, 8.1, 8.2 show examples of brace trees in $\mathcal{T}^{\mathrm{tw}}(2)$. Figures 8.3 and 8.4 show examples of a brace tree in $\mathcal{T}^{\mathrm{tw}}(1)$. On figures, neutral vertices of a brace tree in $\mathcal{T}^{\mathrm{tw}}(n)$ are depicted by black circles and vertices in $V_{\mathrm{lab}}(T)$ are depicted by white circles with the corresponding numbers inscribed.



FIG. 8.1. A brace tree $T_{\cup} \in \mathcal{T}^{\mathrm{tw}}(2)$



FIG. 8.2. A brace tree $T_{\cup^{opp}} \in \mathcal{T}^{\mathrm{tw}}(2)$



FIG. 8.3. A brace tree $T_{\bullet 1} \in \mathcal{T}^{\mathrm{tw}}(1)$



FIG. 8.4. A brace tree $T_{1\bullet} \in \mathcal{T}^{\mathrm{tw}}(1)$

We have the obvious bijection between brace trees in $\mathcal{T}^{\mathrm{tw}}(n)$ with r neutral vertices and linear combinations (8.5). This bijection assigns to a brace tree T' with r neutral vertices the linear combination (8.5) where T is obtained from T' by labeling the neutral vertices by $1, 2, \dots, r$ in any possible way and shifting the labels for vertices in $V_{\mathrm{lab}}(T)$ up by r.

In virtue of this bijection, the n-th space $\mathrm{Tw}\,\mathsf{BT}(n)$ of $\mathrm{Tw}\,\mathsf{BT}$ is the space of (finite) linear combinations of brace trees in $\mathcal{T}^{\mathrm{tw}}(n)$. Furthermore, each brace tree $T' \in \mathcal{T}^{\mathrm{tw}}(n)$ carries the degree

$$|T| = 2|V_{\nu}(T)| - |E(T)| + 1\,, \tag{8.7}$$

where $E(T)$ is the set of all edges of T. In other words, each non-root edge carries degree -1 and each neutral vertex carries degree 2.

Using this description of $\mathrm{Tw}\,\mathsf{BT}$ it is easy to define the elementary insertions in terms of brace trees. Indeed, let $T \in \mathcal{T}^{\mathrm{tw}}(n)$, $T' \in \mathcal{T}^{\mathrm{tw}}(k)$, $1 \le i \le k$ and let v_i be the vertex in $V_{\mathrm{lab}}(T')$ with label i. If v_i is a leaf (i.e. v_i does not have incoming edges) then the vector $T' \circ_i T \in \mathrm{Tw}\,\mathsf{BT}(n+k-1)$ is represented by a tree T'' which is obtained from T' by erasing the vertex v_i and gluing the tree T

via identifying the root edge of T with the edge originating at v_i. After this operation we relabel elements of the set $V_{\mathrm{lab}}(T'') = V_{\mathrm{lab}}(T) \sqcup (V_{\mathrm{lab}}(T') \setminus \{v_i\})$ in the obvious way.

Let us now consider the case when v_i has $q \ge 1$ incoming edges. Since the tree T' is planar these q incoming edges are totally ordered. So we denote them by $e_1, e_2, \dots, e_q$ keeping in mind that

$$e_1 < e_2 < \dots < e_q\,. \tag{8.8}$$

The desired vector $T' \circ_i T \in \mathrm{Tw}\,\mathsf{BT}(n+k-1)$ is represented by the sum

$$T' \circ_i T = \sum_{\alpha} (-1)^{f(\alpha)} T_{\alpha} \tag{8.9}$$

where T_α is obtained from T' and T following these steps:

- first, we erase the vertex $v_i \in V_{\mathrm{lab}}(T')$ and glue the tree T via identifying the root edge T with the edge originating from v_i,
- second, we attach the edges $e_1, e_2, \dots, e_q$ to vertices in the set $V_{\mathrm{lab}}(T) \sqcup V_\nu(T)$
- finally, we relabel elements of the set $V_{\mathrm{lab}}(T'') = V_{\mathrm{lab}}(T) \sqcup (V_{\mathrm{lab}}(T') \setminus \{v_i\})$ in the obvious way.

Ways of connecting the edges $e_1, e_2, \dots, e_q$ to vertices in the set $V_{\mathrm{lab}}(T) \sqcup V_\nu(T)$ should obey the obvious analog of Condition 7.1:

Condition 8.1. *The restriction of the total order on the set $E(T_\alpha)$ of T_α to the subset $\{e_1, e_2, \dots, e_q\}$ should coincide with the order* (8.8).

The summation in (8.9) goes over all ways α of connecting the edges $e_1, e_2, \dots, e_q$ to vertices in the set $V_{\mathrm{lab}}(T) \sqcup V_\nu(T)$ satisfying Condition 8.1. The sign factor $(-1)^{f(\alpha)}$ is define in the same way as for elementary insertions in BT.

Example 8.2. *Let $T_{\circ\circ}$ (resp. $T_{\cup}$) be the brace tree in $\mathcal{T}^{\mathrm{tw}}(2)$ depicted on figure 7.3 (resp. figure 8.1). Then the vector $T \circ_1 T_{\cup} \in \mathrm{Tw}\,\mathsf{BT}(3)$ equals to the sum shown on figure 8.5.*



FIG. 8.5. The elementary insertion $T \circ_1 T_{\cup}$



FIG. 8.6. The brace tree $T_{\bullet\bullet} \in \mathcal{T}^{\mathrm{tw}}(0)$

Remark 8.3. *Let us recall that* $\mathsf{BT}(0) = \mathbf{0}$. *Hence, due to Remark 3.6, the graded vector space of the dg Lie algebra*

$$\mathcal{L}_{\mathsf{BT}} = \mathrm{Conv}(\Lambda^2\mathsf{coCom}, \mathsf{BT}) \tag{8.10}$$

is canonically identified with $\mathbf{s}^{-2}\mathrm{Tw}\,\mathsf{BT}(0)$ *Thus, we can use brace trees in* $\mathcal{T}^{\mathrm{tw}}(0)$ *to represent vectors in the dg Lie algebra* (8.10).

However, the degree formula should be adjusted as follows: a brace tree $T \in \mathcal{T}^{\mathrm{tw}}(0)$ *representing a vector in* $\mathcal{L}_{\mathsf{BT}}$ *carries the degree*

$$|T| = 2|V_\nu(T)| - |E(T)| - 1\,. \tag{8.11}$$

For example, the vector $T_{\bullet\bullet}$ *depicted on figure 8.6 is the Maurer-Cartan element in* $\mathcal{L}_{\mathsf{BT}}$ *corresponding to the map* (8.1). *This vector carries degree* 1 *.*

8.1. **The differential on** $\mathrm{Tw}\,\mathsf{BT}$. Following the general procedure, the differential $\partial(T)$ of a brace tree $T \in \mathrm{Tw}\,\mathsf{BT}(n)$ is given by the formula

$$\partial(T) = T_{\bullet\bullet}\cdot T + \kappa(T_{\bullet\bullet}) \circ_1 T - (-1)^{|T|} \sum_{i=1}^{n} T \circ_i \kappa(T_{\bullet\bullet})\,. \tag{8.12}$$

Here $T_{\bullet\bullet}\cdot$ denotes the auxiliary action (3.13) of the Lie algebra $\mathcal{L}_{\mathsf{BT}}$ on the operad $\mathrm{Tw}\,\mathsf{BT}$ and

$$\kappa(T_{\bullet\bullet}) = T_{\bullet 1} + T_{1\bullet}\,, \tag{8.13}$$

where $T_{\bullet 1}$ and $T_{1\bullet}$ are the brace trees depicted on figures 8.3 and 8.4, respectively. Unfolding the definition of the auxiliary action (3.13) and using (8.13) we get

$$\partial(T) = -(-1)^{|T|} \sum_{v \in V_\nu(T)} T_v \circ_{n+1} T_{\bullet\bullet} + T_{1\bullet} \circ_1 T + T_{\bullet 1} \circ_1 T \tag{8.14}$$

$$-(-1)^{|T|} \sum_{i=1}^{n} T \circ_i T_{1\bullet} - (-1)^{|T|} \sum_{i=1}^{n} T \circ_i T_{\bullet 1}\,.$$

where T_v is a brace tree in $\mathcal{T}^{\mathrm{tw}}(n+1)$ which is obtained from T by replacing the neutral vertex v by a "white" vertex with label $n+1$. In other words, $T_{\bullet\bullet}\cdot T$ is the sum over insertions of the brace tree $T_{\bullet\bullet}$ (see fig. 8.6) into neutral vertices of T with appropriate signs. The expression $T_{1\bullet} \circ_1 T$ is the sum of brace trees which are obtained from T by attaching the single edge with a neutral vertex on one end to all non-root vertices of T . The brace tree $T_{\bullet 1} \circ_1 T$ is obtained from the tree T by dividing the root edge into two parts and inserting a neutral vertex in the middle. The expression $T \circ_i T_{1\bullet}$ (resp. the expression $T \circ_i T_{\bullet 1}$) is obtained from T by inserting $T_{1\bullet}$ (resp. $T_{\bullet 1}$) into the vertex of T with label i .



FIG. 8.7. The brace tree $T_{\bullet\bullet 1} \in \mathcal{T}^{\mathrm{tw}}(1)$

Example 8.4. *Let $T_{\circ\circ}$, $T_{\cup}$, $T_{\cup^{opp}}$, $T_{\bullet 1}$ be the brace trees depicted on figures 7.3, 8.1, 8.2, and 8.3. Direct computations show that $T_{\cup}$ and $T_{\cup^{opp}}$ are ∂-closed and $T_{\circ\circ}$ is not ∂-closed. Instead, we have*

$$\partial(T_{\circ\circ}) = T_{\cup^{opp}} - T_{\cup}\,. \tag{8.15}$$

In other words, $T_{\cup}$ is cohomologous to $T_{\cup^{opp}}$.

For the brace tree $T_{\bullet 1}$ we have

$$\partial(T_{\bullet 1}) = T_{\bullet\bullet 1}\,. \tag{8.16}$$

where $T_{\bullet\bullet 1}$ is the brace tree depicted on figure 8.7.

The operad $\mathrm{Tw}\,\mathsf{BT}$ acts naturally on the Hochschild cochain complex $C^\bullet(A)$ of any (flat) A_∞-algebra A. This action is described in Appendix B.

9. THE OPERAD OF BRACES Br À LA KONTSEVICH-SOIBELMAN

In this section we define a suboperad Br of $\mathrm{Tw}\,\mathsf{BT}$. The operad Br coincides with what M. Kontsevich and Y. Soibelman called the "minimal operad" in [18]. By slightly abusing notation we will call Br the operad of braces and we will refer to algebras over Br as brace algebras. At the end of this section we will show that the embedding $\mathsf{Br} \hookrightarrow \mathrm{Tw}\,\mathsf{BT}$ is a quasi-isomorphism of operads.

We will say that a brace tree $T \in \mathcal{T}^{\mathrm{tw}}(n)$ is *admissible* if T satisfies the following condition:

Condition 9.1. *Every neutral vertex of T has at least 2 incoming edges.*

Thus figures 7.3, 8.1, and 8.2 show examples of admissible brace trees while figures 8.3, 8.4, 8.6 show examples of inadmissible brace trees.

We define $\mathsf{Br}(n)$ as the subspace of $\mathrm{Tw}\,\mathsf{BT}(n)$ which is spanned by admissible brace trees. It is easy to see that, in $\mathcal{T}^{\mathrm{tw}}(0)$ there are no admissible brace trees. Hence,

$$\mathsf{Br}(0) = \mathbf{0}. \tag{9.1}$$

Furthermore, in $\mathcal{T}^{\mathrm{tw}}(1)$, there is exactly one admissible brace tree. This brace tree is depicted on figure 7.2. Thus

$$\mathsf{Br}(1) = \mathbb{K}\,. \tag{9.2}$$

We claim that

Proposition 9.2. *The sub-collection $\big\{\mathsf{Br}(n)\big\}_{n\ge 0}$ is a suboperad of* $\mathrm{Tw}\,\mathsf{BT}$.

Proof. It is obvious that for admissible brace trees $T' \in \mathcal{T}^{\mathrm{tw}}(k)$ and $T \in \mathcal{T}^{\mathrm{tw}}(n)$

$$T' \circ_i T\,, \qquad 1 \le i \le k$$

are linear combinations of admissible brace trees. Thus it remains to prove that for every brace tree ∂T is a linear combination of admissible brace trees. Concretely, we have to show that in ∂T there occur no brace trees with 1- or 2-valent neutral vertices.

Let $e(T)$ be the number of all edges of T. The vertex splitting operations produce $2e(T)$ terms with a 2-valent neutral vertex, 2 for each of the $e(T)$ edges. Concretely,



Here the left term comes from splitting of the lower gray vertex and the right term comes from splitting of the upper gray vertex. The upper gray vertex can be either labeled or neutral, in particular, it may have valency 1. The lower gray vertex may be either a labeled vertex, or a neutral vertex, or the root vertex. (Of course, in the latter case the lower vertex has valency 1.) The remainder of the brace tree is omitted. Comparing the signs factors, we see that these two terms cancel each other, so the differential does not produce valence 2 neutral vertices.

Next consider valence 1 neutral vertices. Again, the corresponding brace trees come in pairs:



Here the left term comes from splitting of the shown gray (i.e., white or black) vertex and the right term comes from the linear combination $\kappa(T_{\bullet\bullet}) \circ_1 T$ in (8.12). Again, checking the signs, we see that these two terms cancel each other and no valence 1 neutral vertex is produced.

□

9.1. **A simpler description of the differential on** Br**.** Let T be an admissible brace tree in $\mathcal{T}^{\mathrm{tw}}(n)$, v be a vertex in $V_{\mathrm{lab}}(T)$ carrying label i, and $\kappa(T_{\bullet\bullet})$ be the sum $T_{\bullet 1} + T_{1 \bullet}$ of inadmissible brace trees depicted on figures 8.3 and 8.4. We denote by $\partial_v(T)$ a vector in $\mathsf{Br}(n)$ which is obtained from the sum

$$T \circ_i \kappa(T_{\bullet\bullet})$$

by omitting all non-admissible brace trees. Let v be a neutral vertex of T. To define $\partial_v(T)$ we change the color of vertex v to white and assign to v label $n+1$. We denote this new brace tree by T_v and obtain $\partial_v(T)$ from the sum

$$T_v \circ_{n+1} T_{\bullet\bullet}$$

via omitting all non-admissible brace trees.

Finally, we declare that the differential ∂ on Br is the vector $\partial(T) \in \mathsf{Br}(n)$ is represented by the sum

$$\partial(T) = \sum_{v \in V_{\mathrm{lab}}(T) \sqcup V_{\nu}(T)} \pm \partial_v(T)\,, \tag{9.3}$$

where the sign factors are determined in the same way as in formula (8.12).

9.2. **The inclusion $\mathsf{Br} \hookrightarrow \mathrm{Tw}\,\mathsf{BT}$ is a quasi-isomorphism.** The goal of this section is to prove the following theorem

Theorem 9.3. *The operad map*

$$\mathsf{Br} \hookrightarrow \mathrm{Tw}\,\mathsf{BT}$$

is a quasi-isomorphism.

Proof. Let T be a brace tree in $\mathcal{T}^{\mathrm{tw}}(n)$ and let $\nu_{\le 2}(T)$ be the total number of neutral vertices of valence ≤ 2. For example, a brace tree T is admissible if and only if

$$\nu_{\le 2}(T) = 0\,.$$

We observe that for every brace tree $T \in \mathcal{T}^{\mathrm{tw}}(n)$ the linear combination

$$\partial(T)$$

involves brace trees T_i with $\nu_{\le 2}(T_i) = \nu_{\le 2}(T)$ or $\nu_{\le 2}(T_i) = \nu_{\le 2}(T) + 1$.

This observation allows us to introduce on $\mathrm{Tw}\,\mathsf{BT}$ the following increasing filtration

$$\dots \subset \mathcal{F}^{m-1}\mathrm{Tw}\,\mathsf{BT} \subset \mathcal{F}^{m}\mathrm{Tw}\,\mathsf{BT} \subset \mathcal{F}^{m+1}\mathrm{Tw}\,\mathsf{BT} \subset \dots\,, \tag{9.4}$$

where $\mathcal{F}^m\mathrm{Tw}\,\mathsf{BT}(n)$ consists of linear combinations of brace trees $T \in \mathcal{T}^{\mathrm{tw}}(n)$ satisfying the condition:

$$\nu_{\le 2}(T) - \deg(T) \le m\,. \tag{9.5}$$

The restriction of (9.4) on Br gives us the "silly" filtration

$$\mathcal{F}^m\mathsf{Br}(n)^k = \begin{cases} \mathsf{Br}(n)^k & \text{if } k \ge -m\,, \\ \mathbf{0} & \text{otherwise} \end{cases} \tag{9.6}$$

with the associated graded complex carrying the zero differential.

To describe the associated graded complex for $\mathrm{Tw}\,\mathsf{BT}$, we introduce the notion of a *core* for brace trees in $\mathcal{T}^{\mathrm{tw}}(n)$. Namely, for a brace tree $\Gamma \in \mathcal{T}^{\mathrm{tw}}(n)$, its core Γ' obtained by deleting all neutral bivalent vertices and connecting their incident edges. Here is an example:



It is easy to see that the associated graded complex

$$\mathrm{Gr}\,\mathrm{Tw}\,\mathsf{BT}(n)$$

splits into the direct sum over all possible cores with n labeled vertices

$$\bigoplus_{\text{cores }\Gamma'} \mathrm{Tw}\,\mathsf{BT}_{\Gamma'}\,, \tag{9.7}$$

where $\mathrm{Tw}\,\mathsf{BT}_{\Gamma'}$ is spanned by brace trees with the given core Γ'.

Furthermore, for each core Γ' the subcomplex $\mathrm{Tw}\,\mathsf{BT}_{\Gamma'}$ is isomorphic to the tensor product

$$\mathrm{Tw}\,\mathsf{BT}_{\Gamma'} \cong \bigotimes_{\text{edges } e \text{ of } \Gamma'} V_e\,,$$

where

$$V_e = \begin{cases} \mathbb{K} \xrightarrow{id} \mathbb{K} \xrightarrow{0} \mathbb{K} \xrightarrow{id} \mathbb{K} \cdots & \text{if } e \text{ connects to a univalent neutral vertex,} \\ \mathbb{K} \xrightarrow{0} \mathbb{K} \xrightarrow{id} \mathbb{K} \xrightarrow{0} \mathbb{K} \cdots & \text{otherwise.} \end{cases}$$

The various copies of $\mathbb{K}$ correspond to strings of bivalent neutral vertices of various lengths:



As before, the shown gray vertices stand for either labeled or neutral vertices. Any lower gray vertex may also be the root vertex. The rest of the brace tree is omitted, and just indicated by some stubs at the gray vertices. Clearly the cohomology of V_e is trivial if e connects to a univalent neutral vertex and $\mathbb{K}$ otherwise. The representative in the latter case is a single edge (i.e., a string of zero neutral vertices).

Therefore the cochain complex $\mathrm{Tw}\,\mathsf{BT}_{\Gamma'}$ is acyclic if Γ' contains at least one univalent neutral vertex and

$$H^\bullet\big(\mathrm{Tw}\,\mathsf{BT}_{\Gamma'}\big) \cong \mathbb{K}\langle\{\Gamma'\}\rangle \tag{9.8}$$

if the core Γ' does not have univalent neutral vertices. Hence the embedding

$$\mathsf{Br}(n) \hookrightarrow \mathrm{Tw}\,\mathsf{BT}(n)$$

induces a quasi-isomorphism on the level of associated graded complexes.

Combining this observation with the fact that the filtrations (9.4) and (9.6) are cocomplete and locally bounded, we deduce Theorem 9.3 from [8, Lemma A.3]. □

9.3. **The dg operad Br is a homotopy fixed point of Tw.** Various solutions of the Deligne conjecture on Hochschild cochain complex [1], [2], [9], [22], [18], [26], [27] imply that the dg operad Br is quasi-isomorphic to the dg operad

$$C_{-\bullet}(E_2, \mathbb{K})$$

of singular chains for the little disc operad E_2.

Combining this statement with the formality [17], [25] for the dg operad $C_{-\bullet}(E_2, \mathbb{K})$, we conclude that the dg operad Br (as well as $\mathrm{Tw}\,\mathsf{BT}$) is quasi-isomorphic[14] to the operad Ger.

Thus Theorem 5.1 and Corollary 5.12 imply that

Corollary 9.4. *The dg operads* Br *and* $\mathrm{Tw}\,\mathsf{BT}$ *are homotopy fixed points for* Tw . □

9.4. **Br is a Tw-coalgebra.** Consider the composition

$$\mathsf{Br} \to \mathrm{Tw}\,\mathsf{BT} \to \mathrm{Tw}\,\mathrm{Tw}\,\mathsf{BT}.$$

We claim that the image actually lands in $\mathrm{Tw}\,\mathsf{Br} \subset \mathrm{Tw}\,\mathrm{Tw}\,\mathsf{BT}$. Indeed, neither the first nor the second map creates new neutral vertices, and hence no neutral vertices of valence 1 or 2 in particular. Hence we have a map

$$c : \mathsf{Br} \to \mathrm{Tw}\,\mathsf{Br}.$$

It follows almost immediately from the Tw colagebra axioms for $\mathrm{Tw}\,\mathsf{BT}$ that this map endows Br with the structure of a Tw-coalgebra. Summarizing, we arrive at the following result.

Lemma 9.5. Br *is a sub* Tw*-coalgebra of* $\mathrm{Tw}\,\mathsf{BT}$. □

10. Every solution of Deligne's conjecture is homotopic to a one that is compatible with twisting

We are now ready to prove Theorem 1.1, which roughly states that every solution of Deligne's conjecture is homotopic to a one that is compatible with twisting.

We will deduce this result from the following more general statement:

Theorem 10.1. *Let* $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}$, $\Lambda\mathsf{Lie}_\infty \to \mathcal{O}'$ *be two objects of the under-category* $\Lambda\mathsf{Lie}_\infty \downarrow$ *Operads and let* $\mathcal{O} \xrightarrow{F} \mathrm{Tw}\,\mathcal{O}'$ *be a morphism of dg operads compatible with the maps from* $\Lambda\mathsf{Lie}_\infty$.

---

[14] In Appendix C we give direct proof of the fact that $H^\bullet(\mathsf{Br}) \cong \mathsf{Ger}$.

If $(\mathcal{O}, c)$ *is a* Tw *-coalgebra and the dg operads* $\mathcal{O}$, $\mathrm{Tw}\,\mathcal{O}'$ *are homotopy fixed points for* Tw *, then the morphism*

$$F' := (\mathrm{Tw}\,(\eta_{\mathcal{O}'} \circ F)) \circ c \colon \mathcal{O} \to \mathrm{Tw}\,\mathcal{O}' \tag{10.1}$$

is homotopy equivalent to F*. Furthermore,* F' *is a morphism of* Tw *-coalgebras.*

This theorem can be restated as follows: if $\mathcal{O}$ and $\mathrm{Tw}\,\mathcal{O}'$ are homotopy fixed points for Tw (with $\mathcal{O}$ being a Tw -coalgebra) then each homotopy class of maps of dg operads $\mathcal{O} \to \mathrm{Tw}\,\mathcal{O}'$ has at least one representative which respects the Tw -coalgebra structures on $\mathcal{O}$ and $\mathrm{Tw}\,\mathcal{O}'$. Furthermore, such a representative may be written down explicitly.

Proof. For the homotopy theory of dg operads in the generality required for this proof we refer the reader to [10]. In particular, the notion of homotopy between operad morphisms is borrowed from *loc. cit.*

Abbreviate $f := \eta_{\mathcal{O}'} \circ F$ and consider the following commutative diagram:

$$\begin{tikzcd}
\mathcal{O} \arrow[r, "F"] \arrow[dr, "f"'] & \mathrm{Tw}\,\mathcal{O}' \arrow[d, "\eta_{\mathcal{O}'}"] \\
 & \mathcal{O}'
\end{tikzcd} \tag{10.2}$$

Applying the functor Tw to (10.2) we obtain the commutative diagram:

$$\begin{tikzcd}
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,F"] \arrow[dr, "\mathrm{Tw}\,f"'] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}' \arrow[d, "\mathrm{Tw}\,\eta_{\mathcal{O}'}"] \\
 & \mathrm{Tw}\,\mathcal{O}'
\end{tikzcd} \tag{10.3}$$

Next consider the following diagram:

$$\begin{tikzcd}
\mathrm{Tw}\,\mathcal{O} \arrow[rr, "\mathrm{Tw}\,F"] \arrow[d, bend right, "\eta_{\mathcal{O}}"'] \arrow[drr, "\mathrm{Tw}\,f" description] & & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}' \arrow[d, "\mathrm{Tw}\,\eta_{\mathcal{O}'}"'] \arrow[d, bend left, "\eta_{\mathrm{Tw}\,\mathcal{O}'}"] \\
\mathcal{O} \arrow[u, "c"'] \arrow[rr, "F"] \arrow[r, phantom, "?" near start] & & \mathrm{Tw}\,\mathcal{O}'
\end{tikzcd} \tag{10.4}$$

Our goal is to show that the triangle marked with "?" is homotopy commutative. We will do this by showing that all the other cells (including the big cell around the diagram) homotopy commute. We saw above that the upper right triangle commutes. Furthermore the big outer square commutes because η is a natural transformation. The small cell on the left commutes since $\eta_{\mathcal{O}} \circ c = id_{\mathcal{O}}$ by the definition of a Tw coalgebra. Finally consider the small cell on the right. Note that, by the defining property of the counit, the following diagram commutes:

$$\begin{tikzcd}
\mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}' \arrow[r, "\mathrm{Tw}\,\eta_{\mathcal{O}'}"] & \mathrm{Tw}\,\mathcal{O}' \\
\mathrm{Tw}\,\mathcal{O}' \arrow[u, "\mathfrak{D}_{\mathcal{O}'}"] \arrow[r, "\mathfrak{D}_{\mathcal{O}'}"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}' \arrow[u, "\eta_{\mathrm{Tw}\,\mathcal{O}'}"']
\end{tikzcd}$$

All arrows are quasi-isomorphisms since $\mathrm{Tw}\,\mathcal{O}'$ is a homotopy fixed point of Tw. Hence the morphisms $\mathrm{Tw}\,\eta_{\mathcal{O}'}$ and $\eta_{\mathrm{Tw}\,\mathcal{O}'}$ are homotopic. This shows that the triangle ? in the previous diagram is homotopy commutative.

It remains to prove that F' (10.1) is a morphism of Tw-coalgebras.

For this purpose we consider the following diagram

$$\begin{tikzcd}
\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,(c)"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O} \arrow[r, "\mathrm{Tw}\,\mathrm{Tw}\,(f)"] & \mathrm{Tw}\,\mathrm{Tw}\,\mathcal{O}' \\
\mathcal{O} \arrow[u, "c"] \arrow[r, "c"] & \mathrm{Tw}\,\mathcal{O} \arrow[u, "\mathfrak{D}_{\mathcal{O}}"] \arrow[r, "\mathrm{Tw}\,(f)"] & \mathrm{Tw}\,\mathcal{O}' \arrow[u, "\mathfrak{D}_{\mathcal{O}'}"]
\end{tikzcd} \tag{10.5}$$

The left square of diagram (10.5) commutes since $c : \mathcal{O} \to \mathrm{Tw}\,\mathcal{O}$ is the comultiplication map for the Tw-coalgebra $\mathcal{O}$. The right square of diagram (10.5) commutes since $\mathfrak{D}$ is a natural transformation from Tw to $\mathrm{Tw}\,\mathrm{Tw}$. Thus the ambient rectangle forms a commutative diagram.

On the other hand, the composition $\mathrm{Tw}\,(f) \circ c$ of the lower horizontal arrows is F' and the composition $\mathrm{Tw}\,\mathrm{Tw}\,(f) \circ \mathrm{Tw}\,(c)$ of upper horizontal arrows is $\mathrm{Tw}\,(F')$. Hence F' is indeed a morphism of Tw-coalgebras.

Theorem 10.1 is proved. □

In order to prove Theorem 1.1 we need a technical property of operad maps from Ger_∞ to BT.

So let $f : \mathsf{Ger}_\infty \to \mathsf{BT}$ be a map of dg operads and let

$$\alpha \in \mathrm{Conv}(\mathsf{Ger}^\vee, \mathsf{BT}) \cong \prod_{n \ge 2} \Big(\mathsf{BT}(n) \otimes \Lambda^{-2}\mathsf{Ger}(n) \Big)^{S_n} \tag{10.6}$$

be the Maurer-Cartan element which corresponds to f.

Using the fact that every vector in $\mathsf{BT}(n)$ carries degree $1-n$, we deduce that α has the form

$$\alpha = \sum_{n \ge 2,\, i} \sum_{\sigma \in S_n} \sigma\Big(X_{n,i} \otimes v^1_{n,i} v^2_{n,i} \Big)\,, \tag{10.7}$$

where $X_{n,i} \in \mathsf{BT}(n)$ and

$$v^1_{n,i} v^2_{n,i} \in \mathbf{s}^2 S^2(\Lambda^{-1}\mathsf{Lie}(b_1, \dots, b_n)) \cap \Lambda^{-2}\mathsf{Ger}(n)\,.$$

We claim that

Proposition 10.2. *If* $f : \mathsf{Ger}_\infty \to \mathsf{BT}$ *is an operad map satisfying the condition*

$$f(\mathbf{s}\, 1^{\mathfrak{c}}_2) = T_{\circ\circ} + \sigma_{12}(T_{\circ\circ})\,, \tag{10.8}$$

with $T_{\circ\circ}$ being the brace tree shown on figure 7.3, then the Maurer-Cartan element α does not involve monomials of the form

$$T \otimes b_1 v(b_2, \dots b_n)\,, \qquad n \geq 3\,,$$

where v is a $\Lambda^{-1}\mathsf{Lie}$-monomial and T is a brace tree with vertex ① having valency 1 *or* 2*.*

We will postpone the technical proof of this proposition to Section 10.1 and complete the proof of Theorem 1.1.

According to Section 4.3, Ger_∞ is a Tw-coalgebra. Furthermore, by Corollaries 5.13 and 9.4 the dg operads Ger_∞ and $\mathrm{Tw}\,\mathsf{BT}$ are homotopy fixed points for Tw. Thus, given an operad quasi-isomorphism $F : \mathsf{Ger}_\infty \to \mathsf{Br}$ we can apply Theorem 10.1 to $\mathcal{O} = \mathsf{Ger}_\infty$, $\mathcal{O}' = \mathsf{BT}$ and the composition

$$\widetilde{F} : \mathsf{Ger}_\infty \xrightarrow{F} \mathsf{Br} \hookrightarrow \mathrm{Tw}\,\mathsf{BT}\,.$$

Since $\widetilde{F}$ is a quasi-isomorphism from Ger_∞ to $\mathrm{Tw}\,\mathsf{BT}$, the composition

$$f = \eta_{\mathsf{BT}} \circ \widetilde{F} : \mathsf{Ger}_\infty \to \mathsf{BT}$$

satisfies condition (10.8). Hence, applying Proposition 10.2, we conclude that for every basis vector $v \in \Lambda^{-1}\mathsf{Lie}(n-1)$ the linear combination

$$\eta_{\mathsf{BT}} \circ \widetilde{F}\big(\, \mathbf{s}\,(b_1 v(b_2, \dots b_n))^* \,\big) \in \mathsf{BT}(n)$$

does not involve brace trees with vertex ① having valency ≤ 2.

This observation implies that the composition

$$\widetilde{F}' = \mathrm{Tw}\,(f) \circ c : \mathsf{Ger}_\infty \to \mathrm{Tw}\,\mathsf{BT}$$

factors through the sub-operad Br.

Since $\widetilde{F}'$ is homotopy equivalent to $\widetilde{F}$, and $\widetilde{F}'$ is compatible with the Tw-coalgebra structures, Theorem 1.1 follows.

□

10.1. **The proof of Proposition 10.2.** The proof of Theorem 1.1 is based on technical Proposition 10.2. Here we give a proof of this proposition.

Condition (10.8) implies that α has the form

$$\alpha = \alpha_0 + \sum_{n \geq 3,\, i} \; \sum_{\sigma \in S_n} \sigma\Big(X_{n,i} \otimes v^1_{n,i} v^2_{n,i}\Big)\,, \tag{10.9}$$

where, as above, $X_{n,i} \in \mathsf{BT}(n)$,

$$v^1_{n,i} v^2_{n,i} \in \mathbf{s}^2 S^2(\Lambda^{-1}\mathsf{Lie}(b_1, \dots, b_n)) \cap \Lambda^{-2}\mathsf{Ger}(n)\,,$$

and α_0 is the degree 1 vector

$$\alpha_0 := T_{\circ\circ} \otimes b_1 b_2 + \sigma_{12}(T_{\circ\circ}) \otimes b_1 b_2\,. \tag{10.10}$$

Recall that, due to Proposition 2.8, the map Av (2.51) gives us an isomorphism to the pre-Lie algebra

$$\mathrm{Conv}(\mathsf{Ger}^\vee, \mathsf{BT}) \cong \prod_{n\ge 2} \Big(\mathsf{BT}(n) \otimes \Lambda^{-2}\mathsf{Ger}(n) \Big)^{S_n}$$

from the pre-Lie algebra

$$\prod_{n\ge 2} \Big(\mathsf{BT}(n) \otimes \Lambda^{-2}\mathsf{Ger}(n) \Big)_{S_n} \tag{10.11}$$

with the pre-Lie product $\bullet'$ given by equation (2.54). So, in our consideration, we may replace $\mathrm{Conv}(\mathsf{Ger}^\vee, \mathsf{BT})$ by the pre-Lie algebra (10.11).

Let us denote by $\mathcal{T}'(n)$ the set of all brace trees for which the labeling agrees with the natural order on the set of vertices. For example, the brace tree $T_{\circ\circ}$ on figure 7.3 belongs to $\mathcal{T}'(2)$ and $\sigma_{12}(T_{\circ\circ})$ does not belong to $\mathcal{T}'(2)$.

Since $\mathsf{BT}(n)$ is a free S_n-module generated elements of $\mathcal{T}'(n)$ we conclude that

$$\Big(\mathsf{BT}(n) \otimes \Lambda^{-2}\mathsf{Ger}(n) \Big)_{S_n} \cong \mathbb{K}\langle \mathcal{T}'(n) \rangle \otimes \Lambda^{-2}\mathsf{Ger}(n)\,. \tag{10.12}$$

Hence the vector space (10.12) has the following natural basis

$$\Big\{ T \otimes w_{n,i} \Big\}_{T \in \mathcal{T}'(n),\, i \in I_n}\,, \tag{10.13}$$

where $w_{n,i}$ are vectors of the basis (4.17) for $\Lambda^{-2}\mathsf{Ger}(n)$.

We also observe that, the $\Lambda^{-1}\mathsf{Lie}$-monomials

$$\Big\{ v_\tau = \{b_{\tau(1)}, \{b_{\tau(2)}, \dots b_{\tau(n-2)}, b_{n-1}\}\}..\} \Big\}_{\tau \in S_{n-2}} \tag{10.14}$$

form a basis in the vector space $\Lambda^{-1}\mathsf{Lie}(n-1)$ and

$$\Big\{ b_i v_\tau(b_1, \dots, \widehat{b_i}, \dots, b_n) \Big\}_{1\le i \le n;\ \tau \in S_{n-2}} \tag{10.15}$$

is a subset of the basis (4.17) for $\Lambda^{-2}\mathsf{Ger}(n)$.

Our goal is to show that the decomposition of the Maurer-Cartan element $\mathrm{Av}^{-1}(\alpha)$ with respect to the basis (10.13) does not involve vectors of the form

$$T \otimes b_i v_\tau(b_1, \dots, \widehat{b_i}, \dots, b_n)\,, \tag{10.16}$$

where T is a brace tree in $\mathcal{T}'(n)$ with vertex ⓘ being either univalent or bivalent.

For this purpose we introduce an operation $V_i^{\circ\!-\!\circ}$ whose input is a brace tree in $\mathcal{T}'(n)$ with vertex ⓘ being either univalent or bivalent. The construction of the output $V_i^{\circ\!-\!\circ}(T)$ depends on whether vertex ⓘ is univalent or bivalent.

If vertex ⓘ of T is univalent then $V_i^{\circ\!-\!\circ}(T)$ is a brace tree in $\mathcal{T}'(n+1)$ which is obtained from T via

- shifting labels of T which are $> i$ up by 1,
- creating a vertex with label $i+1$, and
- attaching the vertex with label $i+1$ to the vertex with label i by a single edge.

If vertex ⓘ of T is bivalent then $V_i^{\circ\!-\!\circ}(T)$ is a brace tree in $\mathcal{T}'(n+1)$ which is obtained from T via

- shifting labels of T which are $> i$ up by 1,
- creating a vertex with label $i+1$,
- detaching the edge terminating at the vertex with label i and attaching it to the vertex with label $i+1$,
- and finally, attaching the vertex with label $i+1$ to the label i by a single edge.

Unfolding the definition, it is not hard to see that for every vector of the form (10.16)

$$\alpha_0 \bullet' T \otimes b_i v_\tau(b_1, \dots, \widehat{b_i}, \dots, b_n) + T \otimes b_i v_\tau(b_1, \dots, \widehat{b_i}, \dots, b_n) \bullet' \alpha_0 = \tag{10.17}$$

$$\pm V_i^{\circ\!-\!\circ}(T) \otimes b_i b_{i+1} v_\tau(b_1, \dots, b_{i-1}, b_{i+2}, \dots, b_n) + \dots\,,$$

where $\dots$ denote the linear combination of the remaining vectors in the basis for (10.12).

It is clear that, if vv' (resp. w) is a vector in $\Lambda^{-2}\mathsf{Ger}(n)$ (resp. in $\Lambda^{-2}\mathsf{Ger}(m)$) with the $\Lambda^{-1}\mathsf{Lie}$-words v, v' having length ≥ 2, then the pre-Lie products

$$\big(T_1 \otimes vv'\big) \bullet' \big(T_2 \otimes w\big)$$

and

$$\big(T_2 \otimes w\big) \bullet' \big(T_1 \otimes vv'\big)$$

do not involve tensor factors of the form $b_i b_{i+1} v_\tau(b_1, \dots, b_{i-1}, b_{i+2}, \dots, b_n)$.

It is also clear that, if the vertex with label j_1 (resp. label j_2) of a brace tree $T_1 \in \mathcal{T}'(n)$ (resp. $T_2 \in \mathcal{T}'(m)$) has valency ≥ 3, then for every pair of basis vectors $v_\tau \in \Lambda^{-1}\mathsf{Lie}(n-1)$, $v_{\tau'} \in \Lambda^{-1}\mathsf{Lie}(m-1)$, the decomposition of the pre-Lie product

$$\big(T_1 \otimes b_{j_1} v_\tau(b_1, \dots, \widehat{b_{j_1}}, \dots, b_n)\big) \bullet' \big(T_2 \otimes b_{j_2} v_{\tau'}(b_1, \dots, \widehat{b_{j_2}}, \dots, b_m)\big)$$

in the basis (10.13) does not involve vectors of the form

$$V_i^{\circ\!-\!\circ}(T) \otimes b_i b_{i+1} v_\tau(b_1, \dots, b_{i-1}, b_{i+2}, \dots, b_n)\,. \tag{10.18}$$

These observations imply that, if the decomposition of the element $\mathrm{Av}^{-1}(\alpha)$ in the basis (10.13) involves a vector of the form (10.16), then

$$\mathrm{Av}^{-1}(\alpha) \bullet' \mathrm{Av}^{-1}(\alpha) \neq 0\,.$$

On the other hand, $\mathrm{Av}^{-1}(\alpha)$ is a Maurer-Cartan element of (10.12). Thus, the proposition follows.

□

10.2. **Proof of Corollary 1.2.** In this section, we give a proof of Corollary 1.2 of Theorem 1.1 stated in the Introduction.

Let us assume that the quasi-isomorphism

$$F : \mathsf{Ger}_\infty \to \mathsf{Br}$$

is compatible with the Tw-coalgebra structures on Ger_∞ and Br. This means that the diagram

$$\begin{tikzcd}
\mathrm{Tw}\,\mathsf{Ger}_\infty \arrow[r, "\mathrm{Tw}\,F"] & \mathrm{Tw}\,\mathsf{Br} \\
\mathsf{Ger}_\infty \arrow[r, "F"] \arrow[u, "c_{\mathsf{Ger}_\infty}"] & \mathsf{Br} \arrow[u, "c_{\mathsf{Br}}"']
\end{tikzcd} \tag{10.19}$$

commutes.

Following Theorem 3.7 and using the Maurer-Cartan element α of $C^\bullet(A)$, we form an operad map

$$\mathfrak{a}^\alpha_A : \mathrm{Tw}\,\mathsf{Br} \to \mathsf{End}_{C^\bullet(A)^\alpha} . \tag{10.20}$$

Unfolding definitions, we see that the composition

$$\mathfrak{a}^\alpha_A \circ \mathrm{Tw}\,(F) \circ c_{\mathsf{Ger}_\infty} : \mathsf{Ger}_\infty \to \mathsf{End}_{C^\bullet(A)^\alpha}$$

gives us the Ger_∞-structure F^α_A and the composition

$$\mathfrak{a}^\alpha_A \circ c_{\mathsf{Br}} \circ F : \mathsf{Ger}_\infty \to \mathsf{End}_{C^\bullet(A)^\alpha}$$

give us the Ger_∞-structure F_{A^α}.

Thus, since diagram (10.19) commutes, the Ger_∞-structures F_{A^α} and F^α_A coincide and the second statement of Corollary 1.2 is proved.

Let us now consider the case when

$$F : \mathsf{Ger}_\infty \to \mathsf{Br}$$

is an arbitrary solution of Deligne's conjecture (not necessarily compatible with Tw-coalgebra structures).

Due to Theorem 1.1, F is homotopy equivalent to a quasi-isomorphism

$$F' : \mathsf{Ger}_\infty \to \mathsf{Br}$$

that is compatible with the Tw-coalgebra structures on Ger_∞ and Br.

Hence diagram (10.19) commutes up to homotopy. Therefore, the two (possibly different) Ger_∞-structures F_{A^α} and F^α_A are indeed homotopy equivalent.

Corollary 1.2 is proved.

□

APPENDIX A. THE OPERAD Ger

A Gerstenhaber algebra is a graded vector space V equipped with a commutative (and associative) product (without identity) and a degree -1 binary operation $\{\,,\,\}$ which satisfies the following relations:

$$\{v_1, v_2\} = (-1)^{|v_1||v_2|}\{v_2, v_1\}\,, \tag{A.1}$$

$$\{v, v_1 v_2\} = \{v, v_1\}v_2 + (-1)^{|v_1||v|+|v_1|}v_1\{v, v_2\}\,, \tag{A.2}$$

$$\{\{v_1, v_2\}, v_3\} + (-1)^{|v_1|(|v_2|+|v_3|)}\{\{v_2, v_3\}, v_1\} + (-1)^{|v_3|(|v_1|+|v_2|)}\{\{v_3, v_1\}, v_2\} = 0\,. \tag{A.3}$$

To define spaces of the operad Ger governing Gerstenhaber algebras we introduce the free Gerstenhaber algebra Ger_n in n dummy variables $a_1, a_2, \dots, a_n$ of degree 0. Next we set $\mathsf{Ger}(0) = \mathbf{0}$ and $\mathsf{Ger}(1) = \mathbb{K}$. And then we declare that, for $n \ge 2$, $\mathsf{Ger}(n)$ is spanned by monomials of Ger_n in which each dummy variable a_i appears exactly once.

The symmetric group S_n acts on $\mathsf{Ger}(n)$ in the obvious way by permuting the dummy variables. It is also clear how to define elementary insertions.

Example A.1. *Let us consider the monomials* $u = \{a_2, a_3\}a_1\{a_4, a_5\} \in \mathsf{Ger}(5)$ *and* $w = \{a_1, a_2\} \in \mathsf{Ger}(2)$ *and compute the insertions* $u \circ_2 w$, $u \circ_4 w$ *and* $w \circ_1 u$. *We get*

$$\begin{aligned}
u \circ_2 w &= -\{\{a_2, a_3\}, a_4\}a_1\{a_5, a_6\}\\
u \circ_4 w &= \{a_2, a_3\}a_1\{\{a_4, a_5\}, a_6\}\\
w \circ_1 u &= \{\{a_2, a_3\}a_1\{a_4, a_5\}, a_6\} = \{a_6, \{a_2, a_3\}a_1\{a_4, a_5\}\}\\
&= \{a_6, \{a_2, a_3\}\}a_1\{a_4, a_5\} - \{a_2, a_3\}\{a_6, a_1\}\{a_4, a_5\}\\
&\qquad - \{a_2, a_3\}a_1\{a_6, \{a_4, a_5\}\}\,.
\end{aligned}$$

It is easy to see that the operad Ger is generated by the vectors $a_1 a_2, \{a_1, a_2\} \in \mathsf{Ger}(2)$.

APPENDIX B. ACTION OF THE OPERAD Tw BT ON THE HOCHSCHILD COCHAIN COMPLEX OF AN A_∞-ALGEBRA

Let A be a cochain complex. We form another cochain complex

$$C^{\bullet}(A) = \bigoplus_{m \ge 0} \mathbf{s}^m\, \mathsf{Hom}(A^{\otimes m}, A) \tag{B.1}$$

with the differential ∂_A coming from A.

Let us show that (B.1) is a naturally an algebra over BT. To define an action BT on (B.1) we observe that the collection

$$\{\mathsf{Hom}(A^{\otimes m}, A)\}_{m \ge 0} \tag{B.2}$$

is an naturally an operad, i.e. the endomorphism operad of A.



FIG. B.1. A brace tree $T \in \mathcal{T}(4)$

Using this observation, we will construct an auxiliary map

$$\varrho : \bigoplus_{N \ge 1} \Big(\mathsf{BT}(N) \otimes \big(C^\bullet(A)\big)^{\otimes N} \Big)_{S_N} \to A \tag{B.3}$$

Given a brace tree $T \in \mathcal{T}(N)$ and N homogeneous vectors

$$P_i \in \mathbf{s}^{m_i}\, \mathsf{Hom}(A^{\otimes m_i}, A)\,, \qquad 1 \le i \le N \tag{B.4}$$

we decorated non-root vertices of T with vectors (B.4) following this rule: the vertex with label i is decorated by the vector P_i.

We say that such a decoration is *admissible* if for every $1 \le i \le N$ the vertex with label i has exactly m_i incoming edges. Otherwise, we say that the decoration is *inadmissible.* In particular, if a decoration is admissible, then each leaf of T is decorated by a vector in

$$\mathsf{Hom}(A^{\otimes 0}, A) = A\,.$$

Since (B.2) is an operad, each brace tree $T \in \mathcal{T}(N)$ with an admissible decoration gives us a vector in A. We denote this vector in A by

$$\mathfrak{m}(T; P_1, P_2, \dots, P_N)\,. \tag{B.5}$$

For example, if T the brace tree depicted on figure B.1, $P_1 \in \mathbf{s}\mathsf{Hom}(A, A)$, $P_2 \in A$, $P_3 \in \mathbf{s}^2\mathsf{Hom}(A^{\otimes 2}, A)$, and $P_4 \in A$ then we have

$$\mathfrak{m}(T; P_1, P_2, P_3, P_4) = P_3(P_1(P_2), P_4)\,.$$

We can now define the map ϱ (B.3).

If T is a brace tree in $\mathcal{T}(N)$ and cochains $P_1, P_2, \dots, P_N$ give us an inadmissible decoration of T then we set

$$\varrho(T, P_1, \dots, P_N) = 0\,. \tag{B.6}$$

Otherwise, we declare that

$$\varrho(T, P_1, \dots, P_N) := (-1)^{\varepsilon(T; P_1, \dots, P_N)} \mathfrak{m}(T; P_1, \dots, P_N) \tag{B.7}$$

where the sign factor $(-1)^{\varepsilon(T; P_1, \dots, P_N)}$ comes from permutation on the set

$$\{E(T) \setminus \{\text{root edge}\}\} \sqcup \{P_1, \dots, P_N\}$$

which we perform when we decorate the tree T with vectors $P_1, P_2, \dots, P_N$.



FIG. B.2. The brace corolla $T_k \in \mathcal{T}(k+1)$

For example, if T is the brace tree depicted on figure B.1, $P_1 \in \mathsf{s}\mathsf{Hom}(A,A)$, $P_2 \in A$, $P_3 \in \mathsf{s}^2\mathsf{Hom}(A^{\otimes 2}, A)$, and $P_4 \in A$ then

$$\varepsilon(T; P_1, \dots, P_N) = |P_3|(|P_1| + |P_2|) + 3|P_3| + 2|P_1| + |P_2|\,.$$

The sign factor $(-1)^{|P_3|(|P_1|+|P_2|)}$ appears because we need to switch from the order (P_1, P_2, P_3, P_4) to the order (P_3, P_1, P_2, P_4) ; the sign factor $(-1)^{3|P_3|}$ appears because P_3 "jumps over" three edges of brace tree T. Similarly, P_1 (resp. P_2) "jumps over" two edges (resp. one edge) of the brace tree T.

We can now define how a brace tree $T \in \mathcal{T}(n)$ acts on n homogeneous vectors

$$P_i \in \mathsf{s}^{m_i}\,\mathsf{Hom}(A^{\otimes\, m_i}, A)\,, \qquad 1 \le i \le n\,. \tag{B.8}$$

For this purpose form the linear combinations ($k \ge 0$)

$$T_k \circ_1 T\,, \tag{B.9}$$

where T_k is the brace corolla shown on figure B.2. and

$$k = \sum_{i=1}^{n} m_i + 1 - n\,. \tag{B.10}$$

Next we set

$$T(P_1, \dots, P_n; a_1, \dots, a_k) = \varrho(T_k \circ_1 T, P_1, \dots, P_n, a_1, \dots, a_k), \tag{B.11}$$

where $a_1, \dots, a_k$ are viewed as vectors in

$$\mathsf{Hom}(A^{\otimes\, 0}, A)\,.$$

Note that, the vectors $a_1, \dots, a_k$ will decorate leaves (of brace trees in the linear combination $T_k \circ_1 T$) with labels $n+1, n+2, \dots, n+k$. Moreover, if equation (B.10) were not satisfied then all decorations of $T_k \circ_1 T$ would be inadmissible.

Tedious but straightforward computations show that formula (B.11) indeed defines an action of the operad BT on (B.1).

Example B.1. *Let T be the brace tree in $\mathcal{T}(4)$ depicted on figure B.1 and*

$$\begin{gathered} P_1 \in \mathsf{s}^2\,\mathsf{Hom}(A \otimes A, A)\,, \qquad P_2 \in \mathsf{s}\,\mathsf{Hom}(A, A)\,, \\ P_3 \in \mathsf{s}^3\,\mathsf{Hom}(A^{\otimes\, 3}, A)\,, \quad P_4 \in \mathsf{Hom}(A^{\otimes\, 0}, A) = A\,. \end{gathered} \tag{B.12}$$



FIG. B.3. The brace trees which contribute to $T(P_1, P_2, P_3, P_4)$

The vector $T(P_1, P_2, P_3, P_4)$ *belongs to*

$$\mathsf{s}^3\mathsf{Hom}(A^{\otimes 3}, A)$$

The linear combination $T_3 \circ_1 T$ *contains a lot of terms. On figure B.3 we list all brace trees (with signs) which get admissible decorations by the vectors*

$$P_1, P_2, P_3, P_4, a_1, a_2, a_3 \tag{B.13}$$

Thus we get

$$T\big(P_1, P_2, P_3, P_4\big)(a_1, a_2, a_3) = -(-1)^{\varepsilon_1} P_3(a_1, P_1(a_2, P_2(a_3)), P_4) + \\ (-1)^{\varepsilon_2} P_3(a_1, P_1(P_2(a_2), a_3), P_4) - (-1)^{\varepsilon_3} P_3(P_1(a_1, P_2(a_2)), a_3, P_4)\,, \tag{B.14}$$

where

$$\varepsilon_1 = \varepsilon(P_3, a_1, P_1, a_2, P_2, a_3, P_4) + 6|P_3| + 5|a_1| + 4|P_1| + 3|a_2| + 2|P_2| + |a_3|\,,$$
$$\varepsilon_2 = \varepsilon(P_3, a_1, P_1, P_2, a_2, a_3, P_4) + 6|P_3| + 5|a_1| + 4|P_1| + 3|P_2| + 2|a_2| + |a_3|\,,$$
$$\varepsilon_3 = \varepsilon(P_3, P_1, a_1, P_2, a_2, a_3, P_4) + 6|P_3| + 5|P_1| + 4|a_1| + 3|P_2| + 2|a_2| + |a_3|\,,$$

and $\varepsilon(\dots)$ *is the sign of the permutation of vectors* $P_1, P_2, P_3, P_4, a_1, a_2, a_3$ *from their standard position* (B.13)*.*

Since BT receives the operad map (8.1) from $\Lambda\mathsf{Lie}$, the cochain complex (B.1) is naturally a $\Lambda\mathsf{Lie}$ algebra. The bracket is the famous Gerstenhaber bracket introduced in [13].

Let us observe that the cochain complex $C^\bullet(A)$ (B.1) is equipped with the obvious decreasing filtration:

$$\mathcal{F}_q C^\bullet(A) = \Big\{ P \in C^\bullet(A) \;\big|\; P(a_1, a_2, \dots, a_k) = 0 \;\forall\; k \le q \Big\} \tag{B.15}$$

and the action of BT on $C^\bullet(A)$ is compatible with this filtration. Thus we may apply the general procedure of twisting described in Section 3 to the completion

$$\hat{C}^\bullet(A) := \prod_{m \ge 0} \mathsf{s}^m\, \mathsf{Hom}(A^{\otimes m}, A) \tag{B.16}$$

of the BT-algebra $C^\bullet(A)$ with respect to the filtration (B.15).

According to Theorem 3.9, filtered $\mathrm{Tw}\,\mathsf{BT}$-algebra structures on $\hat{C}^\bullet(A)$ are in bijection with Maurer-Cartan elements α in $\hat{C}^\bullet(A)$ of the form

$$\alpha = \sum_{k\geq 2} \alpha_k\,, \qquad \alpha_k \in \mathsf{s}^k \mathsf{Hom}(A^{\otimes k}, A) \tag{B.17}$$

i.e. flat A_∞-structures on the cochain complex A.

Moreover, for every Maurer-Cartan element (B.17), the cochain complex $\hat{C}^\bullet(A)$ with the twisted differential (3.22) is exactly the (completed) Hochschild cochain complex of the A_∞-algebra A.

Thus we conclude that, for every flat A_∞-algebra A, its completed Hochschild cochain complex is equipped with a natural action of the operad $\mathrm{Tw}\,\mathsf{BT}$ and hence with a natural action of the operad Br.

Remark B.2. *Sometimes the Maurer-Cartan element* (B.17) *satisfies the condition*

$$\alpha_k = 0$$

for all $k \geq N$ for some given number N. (For example, if A is an associative algebra.) In this case, the action of $\mathrm{Tw}\,\mathsf{BT}$ *(and hence the action of* Br*) is well defined before completing the Hochschild cochain complex.*

Appendix C. A direct computation of the cohomology of Br

In this section, we prove that $H^\bullet(\mathsf{Br}) \cong \mathsf{Ger}$. A proof of this fact was sketched, for example, in [18, Theorem 4]. Here, we decided to give a different proof of this fact for two reasons: first, the sketch given in [18] has gaps, and second, our proof is "topology independent".

Theorem C.1 (Theorem 4, [18]). *The cohomology of the operad* Br *is the Gerstenhaber operad:*

$$H^\bullet(\mathsf{Br}) \cong \mathsf{Ger}\,.$$

The cohomology class $\{a_1, a_2\} \in \mathsf{Ger}(2)$ is represented by the sum $T_{\{\ ,\ \}} =$  *and the cohomology class $a_1 a_2 \in \mathsf{Ger}(2)$ is represented by*

$$T_\wedge = \frac{1}{2}\Bigg($$  $$\Bigg)\,.$$

Remark C.2. *In fact, mapping the generator of* $\Lambda\mathsf{Lie}$ *to the element indicated in Theorem C.1, we obtain a map*

$$\mathsf{j} : \Lambda\mathsf{Lie} \to \mathsf{Br}. \tag{C.1}$$

To see this, one has to check that the Jacobi identity holds in Br. *Theorem C.1 implies that this map is injective and that the image is given by the closed elements of* Br *which do not contain neutral vertices. Note, however, that* j *does not extend to an operad map* $\mathsf{Ger} \to \mathsf{Br}$.

We show that $H^\bullet(\mathsf{Br}(n)) = \mathsf{Ger}(n)$ by induction on n. For $n = 1$ there is nothing to show. So suppose we know that $H^\bullet(\mathsf{Br}(j)) = \mathsf{Ger}(j)$ for $j = 1, 2 \dots, n-1$ and let us tackle the statement for $j = n$. We split

$$\begin{tikzcd}
\mathsf{Br}(n) & = & V_\circ(n) \arrow[loop above, "\delta_0"] \arrow[rr, bend left, "\delta_1"] & \oplus & V_\bullet(n) \arrow[loop above, "\delta_0"]
\end{tikzcd}$$

Here $V_\bullet(n)$ is the subspace of $\mathsf{Br}(n)$ spanned by brace trees whose lowest non-root vertex is neutral, while $V_\circ(n)$ is the subspace of $\mathsf{Br}(n)$ spanned by brace trees whose lowest non-root vertex is labeled. The arrows indicate the several components of the differential. For example, $V_\bullet(n)$ is a subcomplex of $\mathsf{Br}(n)$. Clearly δ_1 induces a map, called $H(\delta_1)$, on δ_0-cohomologies:

$$H(V_\circ(n), \delta_0) \xrightarrow{H(\delta_1)} H(V_\bullet(n), \delta_0)\,. \tag{C.2}$$

Furthermore, it is not hard to see that

$$H(\mathsf{Br}(n)) = \big(\ker H(\delta_1)\big) \oplus \big(\operatorname{coker} H(\delta_1)\big). \tag{C.3}$$

C.1. **Computing $H(V_\circ(n), \delta_0)$.** Let us observe that the linear dual $\mathsf{Br}(n)^*$ can be canonically identified with $\mathsf{Br}(n)$ as vector space and the differential on $\mathsf{Br}(n)^*$ is given by edge contractions instead of vertex splitting. Using this observation we will freely switch back and forth between the cochain complexes under consideration and their linear duals.

Let us begin by computing $H(V_\circ^*(n), \delta_0^*)$.

Claim C.3. *We claim that*

$$H(V_\circ^*(n), \delta_0^*) \cong \mathbf{s}^{n-1}\mathbb{K}^{n!} \cong \mathbf{s}^{n-1}\mathbb{K}[S_n] \tag{C.4}$$

The class corresponding to a permutation $\lambda \in S_n$ is represented by a brace tree T^n_λ shown on figure C.3.

Proof. We proceed by induction on n. For $n = 1$ the statement is clear. Otherwise split:

$$\begin{tikzcd}
V_\circ^* & = & W_1 \arrow[loop above, "\delta_0'"] \arrow[rr, bend left, "\delta_1'"] & \oplus & W_{\geq 2} \arrow[loop above, "\delta_0'"]
\end{tikzcd}$$

Here W_1 is spanned by brace trees in which the lowest non-root vertex has exactly one child and $W_{\geq 2}$ is spanned by brace trees in which the lowest non-root vertex has at least two children. It is easy to see that δ_1' is surjective and that its kernel is spanned by brace trees whose lowest non-root vertex has a labeled vertex as a child. The complex $(\ker \delta_1', \delta_0')$ is isomorphic to $(V_\circ^*(n-1), \delta_0^*)$.

Thus the induction hypothesis implies the desired statement. □

Remark C.4. *Note that every brace tree $T \in \mathsf{Br}(n)^*$ without neutral vertices is automatically closed and in particular δ_0^*-closed. Hence, by Claim C.3, there exists a vector $T' \in V_\circ^*(n)$, such* *that* $T - \delta_0^* T'$ *is a linear combination of string-like brace trees, i.e. brace trees of the form* T^n_λ *(see figure C.3).*

C.2. **Computing** $H(V_\bullet(n), \delta_0)$**.** Let us prove the following claim:

Claim C.5. *For the complex* $(V_\bullet(n), \delta_0)$ *we have*

$$H(V_\bullet(n), \delta_0) \cong \mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n) \;\oplus\; \mathbf{s}\big(\Lambda\mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n)\big)\,, \tag{C.5}$$

where $\odot$ *denotes the plethysm* (2.12) *of collections.*

Proof. Let us filter $V_\bullet(n)$ by the number of children of the lowest non-root vertex.

$$\mathbf{0} = \mathcal{F}^1 V_\bullet(n) \subset \mathcal{F}^2 V_\bullet(n) \subset \mathcal{F}^3 V_\bullet(n) \subset \dots \subset \mathcal{F}^n V_\bullet(n) = V_\bullet(n). \tag{C.6}$$

Here $\mathcal{F}^p V_\bullet(n)$ is spanned by brace trees whose lowest non-root vertex has $\le p$ children. Let us consider the spectral sequence associated to this filtration. The first differential, say d_0, splits vertices except for the lowest non-root vertex. Hence,

$$\mathrm{Gr}\, V_\bullet(n) \cong \big(\mathbf{s}\Lambda\mathsf{coAs}_\circ \odot \mathsf{Br}\big)(n)\,, \tag{C.7}$$

where $\mathbf{s}\Lambda\mathsf{coAs}_\circ$ is the collection with

$$\mathbf{s}\Lambda\mathsf{coAs}_\circ(m) = \begin{cases} \mathbf{s}^{2-m}\mathbb{K}[S_m] \otimes \mathrm{sgn}_m & \text{if } m \ge 2\,, \\ \mathbf{0} \quad \text{otherwise}\,. \end{cases} \tag{C.8}$$

Therefore, by inductive hypothesis, we conclude that

$$E^1 V_\bullet(n) := H(\mathrm{Gr}\, V_\bullet(n), d_0) \cong \big(\mathbf{s}\Lambda\mathsf{coAs}_\circ \odot \mathsf{Ger}\big)(n) \tag{C.9}$$

The differential d_1 on $E^1 V_\bullet(n)$ splits the lowest non-root vertex producing a neutral child node with two children. To describe this cochain complex, we consider the free Gerstenhaber algebra Ger_n in n dummy variables $a_1, a_2, \dots, a_n$ of degree zero. We forget the bracket $\{\ ,\ \}$ on Ger_n can view it merely as the free commutative algebra (without unit)

$$\mathsf{Ger}_n = \mathsf{Com}(\Lambda\mathsf{Lie}_n)$$

generated by the free $\Lambda\mathsf{Lie}$-algebra $\Lambda\mathsf{Lie}_n$ in the dummy variables $a_1, a_2, \dots, a_n$

Next, we introduce the cofree coassociative coalgebra

$$\mathsf{coAs}(\mathbf{s}^{-1}\,\mathsf{Ger}_n) = \bigoplus_{q \ge 1} \big(\mathbf{s}^{-1}\,\mathsf{Ger}_n\big)^{\otimes q} \tag{C.10}$$

and equip it with the coderivation $\mathfrak{d}$ defined by the equation[15]

$$p \circ \mathfrak{d}(\mathbf{s}^{-1}\, v_1 \otimes \dots \otimes \mathbf{s}^{-1}\, v_q) = \begin{cases} (-1)^{|v_1|+1} v_1 v_2 & \text{if } q = 2\,, \\ 0 \quad \text{otherwise}\,, \end{cases} \tag{C.11}$$

---

[15] Note that, since the coalgebra (C.10) is cofree, any coderivation $\mathfrak{d}$ is uniquely determined by its composition with the projection (C.12).

where $v_i \in \mathsf{Ger}_n$ and p is the canonical projection;

$$p : \mathsf{coAs}(\mathbf{s}^{-1}\,\mathsf{Ger}_n) \to \mathbf{s}^{-1}\,\mathsf{Ger}_n\,. \tag{C.12}$$

It is easy to see that the coderivation $\mathfrak{d}$ has degree 1. Moreover, due to associativity of the multiplication on Ger_n, we have

$$\mathfrak{d}^2 = 0\,.$$

In other words, $\mathfrak{d}$ is a differential on the coalgebra (C.10).

For our purposes we need the following truncation of the cochain complex $\mathbf{s}^2\mathsf{coAs}(\mathbf{s}^{-1}\,\mathsf{Ger}_n)$

$$\mathbf{s}^2 T'(\mathbf{s}^{-1}\,\mathsf{Ger}_n) = \bigoplus_{q\ge 2} \mathbf{s}^2\big(\mathbf{s}^{-1}\,\mathsf{Ger}_n\big)^{\otimes\, q} \tag{C.13}$$

with the differential $\mathfrak{d}'$ given by the formula:

(C.14)

$$\mathfrak{d}'\big(\mathbf{s}^2(\mathbf{s}^{-1}\,v_1\otimes\mathbf{s}^{-1}\,v_2\otimes\cdots\otimes\mathbf{s}^{-1}\,v_q)\big) = \begin{cases} \mathbf{s}^2\mathfrak{d}(\mathbf{s}^{-1}\,v_1\otimes\mathbf{s}^{-1}\,v_2\otimes\cdots\otimes\mathbf{s}^{-1}\,v_q) & \text{if } q>2\,, \\ 0 \quad \text{if } q=2\,, \end{cases} \qquad v_i \in \mathsf{Ger}_n\,.$$

It is not hard to see that $E^1 V_\bullet(n)$ (C.9) is isomorphic to the subspace of $\mathbf{s}^2 T'(\mathbf{s}^{-1}\,\mathsf{Ger}_n)$ which is spanned by tensor monomials

$$\mathbf{s}^2(\mathbf{s}^{-1}\,v_1\otimes\mathbf{s}^{-1}\,v_2\otimes\cdots\otimes\mathbf{s}^{-1}\,v_q)\,, \qquad v_i\in\mathsf{Ger}_n\,, \qquad 2\le q\le n$$

in which each variable from the set $\{a_1, a_2, \dots, a_n\}$ appears exactly once. It is easy to see that this subspace is a subcomplex with respect to $\mathfrak{d}'$ and, moreover, the differential d_1 coincides with the restriction of $\mathfrak{d}'$ up to a total sign.

Since the augmentation

$$\dots \xrightarrow{\mathfrak{d}} \big(\mathbf{s}^{-1}\,\mathsf{Ger}_n\big)^{\otimes\,2} \xrightarrow{\mathfrak{d}} \mathbf{s}^{-1}\,\mathsf{Ger}_n \xrightarrow{0} \mathbb{K} \tag{C.15}$$

of the cochain complex (C.10) computes the Hochschild homology

$$HH_{-\bullet}(S(\Lambda\mathsf{Lie}_n), \mathbb{K})$$

of the free commutative algebra $S(\Lambda\mathsf{Lie}_n)$ (with unit) with the trivial coefficients, we conclude that

$$H^\bullet(\mathsf{coAs}(\mathbf{s}^{-1}\,\mathsf{Ger}_n), \mathfrak{d}) = \bigoplus_{q\ge1} S^q(\mathbf{s}^{-1}\,\Lambda\mathsf{Lie}_n)\,, \tag{C.16}$$

and the cohomology class of the symmetric word $(\mathbf{s}^{-1}\,v_1, \mathbf{s}^{-1}\,v_2, \dots, \mathbf{s}^{-1}\,v_q) \in S^q(\mathbf{s}^{-1}\,\Lambda\mathsf{Lie}_n)$ is represented by the cocycle:

$$\sum_{\sigma\in S_q} (-1)^{\varepsilon(\sigma, v_1,\dots,v_q)} \big(\mathbf{s}^{-1}\,v_{\sigma(1)}, \mathbf{s}^{-1}\,v_{\sigma(2)}, \dots, \mathbf{s}^{-1}\,v_{\sigma(q)}\big)\,,$$

where the sign factors $(-1)^{\varepsilon(\sigma, v_1,\dots,v_q)}$ are determined by the Koszul rule.

Using this observation, it is not hard to see that

$$H^\bullet\big(\mathbf{s}^2 T'(\mathbf{s}^{-1}\,\mathsf{Ger}_n), \mathfrak{d}'\big) \cong \bigoplus_{q\ge2} S^q(\Lambda\mathsf{Lie}_n) \;\oplus\; \bigoplus_{q\ge2} \mathbf{s}^2 S^q(\mathbf{s}^{-1}\,\Lambda\mathsf{Lie}_n)\,. \tag{C.17}$$

On the other hand, $E^1 V_\bullet(n)$ is isomorphic to the direct summand of the cochain complex $\big(\mathbf{s}^2 T'(\mathbf{s}^{-1}\,\mathsf{Ger}_n), \mathfrak{d}'\big)$.

Thus we conclude that the following three facts:

- First,

$$E^2 V_\bullet(n) \cong \mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n) \;\oplus\; \mathbf{s}\big(\Lambda\mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n)\big)\,. \tag{C.18}$$

- Second, the class corresponding to the vector

$$\mathbf{s}^2(\mathbf{s}^{-1}\,v_1, \dots, \mathbf{s}^{-1}\,v_q) \in \mathbf{s}\Lambda\mathsf{Com} \odot \Lambda\mathsf{Lie}(n), \qquad q \geq 2$$

  is represented in the associated graded $\mathrm{Gr}\, V_\bullet$ by the cochain

$$\mu\left(\Big(\sum_{\sigma\in S_q}(-1)^{|\sigma|} T_q^\bullet\Big) \otimes \mathfrak{j}(v_1) \otimes \mathfrak{j}(v_2) \otimes \dots \mathfrak{j}(v_q)\right)\,, \tag{C.19}$$

  where μ is the operadic multiplication

$$\mu : \mathsf{Br}(q) \otimes \mathsf{Br}(m_1) \otimes \dots \otimes \mathsf{Br}(m_q) \to \mathsf{Br}(m_1 + \dots + m_q)$$

  and $T_q^\bullet$ is the brace tree shown on figure C.1. Note that the element (C.19) is not closed in $V_\bullet$, only in the associated graded. To obtain a cocycle in $V_\bullet$, a correction term in $\mathcal{F}^{q-1}$ has to be added. We will see below that such a term may always be found.

- Finally, classes corresponding to vectors in $\mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n)$ are represented by cocycles which are obtained by composing the vectors $T_{\{\ ,\ \}}$ and $T_\wedge$ from the statement of Theorem C.1. For example, the vector $\{a_1, a_3\}a_2$ is represented in $E^2 V_\bullet(n)$ by the linear combination shown on figure C.2



FIG. C.1. The brace tree $T_q^\bullet \in \mathcal{T}^{\mathrm{tw}}(q)$



FIG. C.2. A representative of the class $\{a_1, a_3\}a_2$

We claim that the spectral sequence abuts at this point. In other words,

$$E^\infty V_\bullet(n) = E^2 V_\bullet(n)\,. \tag{C.20}$$

Indeed, closed (in $\mathsf{Br}(n)$) representatives for elements of $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n)$ may be given, and hence all higher differentials are zero on $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n)$. Note that

$$\mathsf{s}\big(\Lambda\mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n)\big) \tag{C.21}$$

is concentrated in cohomological degree $2-n$. Hence higher differentials in the spectral sequence must map (some subspace of) (C.21) to (some quotient of) $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n)$. Furthermore there is an action of the symmetric group S_p on $\mathrm{Gr}^p\, V_\bullet(n)$ by permuting the subtrees of the lowest vertex. The space (C.21) is a subspace of the symmetric pieces $(\mathrm{Gr}^p\, V_\bullet(n))^{S_p}$ while $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n)$ lives in the anti-symmetric part of $\mathrm{Gr}^2\, V_\bullet(n)$. Hence the following Lemma is almost evident.

Lemma C.6. *If $x \in (\mathrm{Gr}^p\, V_\bullet(n))^{S_p}$ is a cocycle of cohomological degree $> 2-n$, then x is exact, i.e., there is some $x' \in \mathrm{Gr}^p\, V_\bullet(n)$ whose coboundary is x. Moreover, x' may be chosen to be symmetric, i. e., $x' \in (\mathrm{Gr}^p\, V_\bullet(n))^{S_p}$.*

Proof. $\mathrm{Gr}^p\, V_\bullet(n)$ splits into a direct sum of subcomplexes, one for each isotypical component with respect to the S_p-action. By the above computation, only the symmetric piece and the antisymmetric piece contains any cohomology, and the former only in cohomological degree $2-n$, and the latter only for $p=2$ and in cohomological degrees between $3-n$ and 0. The lemma immediately follows. □

Let us consider a non-trivial cocycle

$$u \in \mathsf{s}\big(\Lambda\mathsf{Com} \odot \Lambda\mathsf{Lie}(n)/\Lambda\mathsf{Lie}(n)\big) \cap (\mathrm{Gr}^p\, V_\bullet(n))^{S_p}$$

and examine how the higher differentials in the spectral sequence act on u. We claim that all these higher differentials vanish. Equivalently, this means that a closed representative for u in $V_\bullet(n)$ may be found. u has a unique representative we call $u_p \in V_\bullet(n)$ that is a linear combination of trees whose lowest non-root vertex has valence p. Now $-\delta_0 u_p \in \mathcal{F}^{p-1}V_\bullet(n)$ represents some cocycle $u'_{p-1} \in \mathrm{Gr}^{p-1}\, V_\bullet(n)$ of cohomological degree $3-n$. By symmetry of u it is also symmetric and hence the coboundary of some symmetric element u_{p-1} by the preceding lemma. u_{p-1} has a unique representative we also call $u_{p-1} \in V_\bullet(n)$ that is a linear combination of trees whose lowest non-root vertex has valence $p-1$. Next, $-\delta_0(u_p + u_{p-1}) \in \mathcal{F}^{p-2}V_\bullet(n)$ represents some symmetric cocycle $u'_{p-2} \in \mathrm{Gr}^{p-2}\, V_\bullet(n)$ which is again the coboundary of some symmetric element u_{p-2} by the preceding lemma. Continuing in this manner, we obtain some closed element

$$u_p + u_{p-1} + \cdots + u_2 \in V_\bullet(n)$$

representing u. Hence Claim C.5 is shown.

□

Remark C.7. *One may, of course, dualize the statement of Claim C.5. The dual statement says that*

$$H(V^*_\bullet(n), \delta^*_0) \cong X \oplus U^*\,,$$

where $X \subset \mathsf{Ger}(n)^*$ is the kernel of $\mathsf{Ger}(n)^* \to \Lambda\mathsf{Lie}(n)^*$ and U^* is the linear dual of (C.21).

Let us also note that explicit representatives for all classes in the subspace

$$H(V_\bullet^*(n),\delta_0^*)_1 \subset H(V_\bullet^*(n),\delta_0^*)$$

spanned by classes containing only trees with exactly one neutral vertex may be given. Let $T^\sigma_{r_1,\dots,r_k}$ be the brace tree depicted in figure C.4, where

$$\sigma = \begin{pmatrix} 1 & \dots & r_1 & r_1+1 & \dots & r_1+r_2 & \dots & n-r_k+1 & \dots & n \\ i_1^1 & \dots & i_{r_1}^1 & i_1^2 & \dots & i_{r_2}^2 & \dots & i_1^k & \dots & i_{r_k}^k \end{pmatrix}$$

is any permutation in S_n with $n = r_1 + r_2 + \dots + r_k$. Each such tree is δ_0^*-closed and hence defines a cohomology class in $H(V_\bullet^*(n),\delta_0^*)_1$.

Lemma C.8. *The cohomology classes in $H(V_\bullet^*(n),\delta_0^*)$ represented by the trees $T^\sigma_{r_1,\dots,r_k}$ span the space $H(V_\bullet^*(n),\delta_0^*)_1$.*

Proof. An equivalent form of the statement is that if x is a cohomology class in $H(V_\bullet(n),\delta_0)$ representable by a linear combination of trees with exactly one neutral vertex, and if furthermore all $T^\sigma_{r_1,\dots,r_k}$ evaluate to zero on x, then x is trivial. To show this equivalent statement let us trace through the proof of Claim C.5. We may assume that x is contained either in the first or the second summand of (C.5). Assume that x is contained in the second summand and nontrivial. Then, by the proof of Claim C.5 we see that it is represented by a linear combination of graphs containing graphs of the form $T^\sigma_{r_1,\dots,r_k}$ for some σ and $r_1,\dots,r_k$. Hence the evaluation of $T^\sigma_{r_1,\dots,r_k}$ on x is non-zero.

Next, if x lives in the first summand of (C.5), an explicit representative for x may be written down, which is a linear combination of elements of the form

$$\sigma' \cdot \mu(T_\wedge \otimes L_1 \otimes L_2)$$

where μ denotes the operadic composition, $\sigma' \in S_n$ and L_1 and L_2 are elements of $\Lambda\mathsf{Lie} \subset \mathsf{Br}$. Since non-zero elements of $\Lambda\mathsf{Lie} \subset \mathsf{Br}$ always contain string-like graphs, the representative for x necessarily contains graphs as in figure C.4, and hence the evaluation of $T^\sigma_{r_1,\dots,r_k}$ on x is necessarily non-zero for some σ and $r_1,\dots,r_k$, unless $x = 0$.

□

C.3. **A technical claim.** To proceed further we need the following technical claim:

Claim C.9. *Let $\{T^n_\lambda\}_{\lambda\in S_n}$ be the family of brace trees depicted on figure C.3. Furthermore, let $T^\sigma_{r_1,\dots,r_k}$ be the brace tree depicted in figure C.4, as introduced above Lemma C.8. Then the vector*

$$\delta_1^*(T^\sigma_{r_1,\dots,r_k}) \tag{C.22}$$

is cohomologous in $(V_\circ^*,\delta_0^*)$ to

$$\sum_{\tau\in \mathrm{Sh}_{r_1,\dots,r_k}} (-1)^{|\tau|}\, T^n_{\sigma\circ\tau^{-1}}\,, \tag{C.23}$$

where $(-1)^{|\tau|}$ is the sign of the permutation τ.



FIG. C.3. The brace tree T^n_λ. Here $\lambda \in S_n$



FIG. C.4. The brace tree $T^{\sigma}_{r_1,\dots,r_k}$

Example C.10. *Before giving the proof, we illustrate the statement of Claim C.9 with an example shown on figure C.5. In the first step on figure C.5, we apply the operator δ^*_1. In the second step we add a coboundary term and compute the sum. In the third step, we add another coboundary term to the result and obtain an expression of the form* (C.23)*.*

Proof. Since the operators δ^*_1 and δ^*_0 commute with the action of the symmetric group, it suffices to prove the desired statement for $\sigma = \mathrm{id} \in S_n$.

The case $n = 2$ is very easy, so we leave it to the reader and take it as the base of our induction on n.

To perform the inductive step, we observe that the set of permutations

$$\left\{ \tau^{-1} \right\}_{\tau \in \mathrm{Sh}_{r_1,\dots,r_k}} \tag{C.24}$$

is the disjoint union of the subsets Σ_q, $q \ge 1$, where Σ_q consists of permutations of the form

$$\begin{pmatrix} 1 & 2 & \dots & n \\ t_q & \bullet & \dots & \bullet \end{pmatrix} \tag{C.25}$$



FIG. C.5. Illustration of Claim C.9

where

$$t_q = \begin{cases} 1 & \text{if } q = 1\,, \\ r_1 + \dots + r_{q-1} + 1 & \text{otherwise} \end{cases} \tag{C.26}$$

and $\bullet\ \dots\ \bullet$ is any unshuffle of the groups of indices:

$$(1, \dots, r_1),\ (r_1 + 1, \dots, r_1 + r_2),\ \dots,\ (t_q + 1, \dots, r_q),\ \dots,\ (n - r_k + 1, \dots, n).$$

Next, we denote by $X^q_{r_1,\dots,r_k}$, $Y^q_{r_1,\dots,r_k}$, and $T^{opp}_{\circ\circ}$ the brace trees shown on figures C.6, C.7, C.8, respectively.



FIG. C.6. The brace tree $X^q_{r_1,\dots,r_k} \in \mathcal{T}(n)$



FIG. C.7. The brace tree $Y^q_{r_1,\dots,r_k} \in \mathcal{T}^{\text{tw}}(n)$



FIG. C.8. The brace tree $T^{opp}_{\circ\circ} \in \mathcal{T}(2)$

We remark that

$$\delta_1^*(T^{\text{id}}_{r_1,\dots,r_k}) = \sum_{q=1}^{k} (-1)^{t_q-1} X^q_{r_1,\dots,r_k} \tag{C.27}$$

and for every $1 \le q \le k$

$$X^q_{r_1,\dots,r_k} - \delta_0^*(Y^q_{r_1,\dots,r_k}) = (t_q, t_q+1, \dots, n)\left(T^{opp}_{\circ\circ} \circ_1 \delta_1^*(T^{\mathrm{id}}_{r_1,\dots,r_q-1,\dots,r_k})\right). \tag{C.28}$$

By inductive assumption, the vector $\delta_1^*(T^{\mathrm{id}}_{r_1,\dots,r_q-1,\dots,r_k})$ is cohomologous in $(V_\circ^*, \delta_0^*)$ to

$$\sum_{\tau' \in \mathrm{Sh}_{r_1,\dots,r_q-1,\dots,r_k}} (-1)^{|\tau'|}\, T^{n-1}_{(\tau')^{-1}}\,. \tag{C.29}$$

Thus Claim C.9 is proved. □

C.4. **Analyzing the map** (C.2)**.** The goal of this subsection is to describe the kernel and the cokernel of the map (C.2). This way we will get the desired isomorphism $H^\bullet(\mathsf{Br}(n)) \cong \mathsf{Ger}(n)$.

More precisely,

Claim C.11. *For the map* (C.2) *we have*

$$\ker\Big(H(\delta_1) : H(V_\circ(n), \delta_0) \to H(V_\bullet(n), \delta_0)\Big) \cong \Lambda\mathsf{Lie}(n)$$

and

$$\mathrm{coker}\Big(H(\delta_1) : H(V_\circ(n), \delta_0) \to H(V_\bullet(n), \delta_0)\Big) \cong \mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n).$$

Proof. We will prove Claim C.11 by analyzing the dual map

$$H(\delta_1^*) : H(V_\bullet^*(n), \delta_0^*) \to H(V_\circ^*(n), \delta_0^*).$$

Recall that the right hand side has dimension $n!$ and classes are represented by string-like trees corresponding to permutations (see figure C.3).

Due to Claim C.9, the image of $H(\delta_1^*)$ in

$$H(V_\circ^*(n), \delta_0^*) \cong \mathbf{s}^{n-1}\mathbb{K}[S_n]$$

is spanned by classes which correspond to shuffle products in $\mathbb{K}[S_n]$.

Hence,

$$\mathrm{coker}\left(H(\delta_1^*)\right) \cong \Lambda\mathsf{Lie}(n)^*$$

or equivalently

$$\ker\left(H(\delta_1)\right) \cong \Lambda\mathsf{Lie}(n)\,.$$

Next let us examine the kernel of the map $\delta_1^* : H(V_\bullet^*, \delta_0^*) \to H(V_\circ^*, \delta_0^*)$. We already mentioned that δ_1^* is zero on all $H(V_\bullet^*, \delta_0^*)_r$ for $r \neq 1$. By Remark C.7 above, $H(V_\bullet^*, \delta_0^*)_1$ consists of two summands, namely the linear dual of (C.21) and a subspace which we call X_1 of $\mathsf{Ger}(n)^*$. Representatives of classes in X_1 are given by antisymmetric linear combinations of a tree with a neutral vertex with two child strings, and the same tree, but with the order of the strings flipped. Since the shuffle product is commutative, such a tree is mapped to zero by δ_1^*, hence $\ker(\delta_1^*) \supset X$. Now we are ready to show that

$$\ker\left(H(\delta_1^*)\right) \cong X \tag{C.30}$$

or equivalently,

$$\operatorname{coker}\big(H(\delta_1)\big) \cong \mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n)\,. \tag{C.31}$$

Indeed, by counting dimensions we get

$$dim\,\mathrm{im}(\delta_1^*) = dim(\mathsf{As}(n)) - dim(\mathsf{Lie}(n)) = n! - (n-1)!.$$

Of course $dim\,\mathrm{im}(\delta_1^*) = codim\,\ker(\delta_1^*)$. Finally

$$codim X = dim U = dim(\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n)) = n! - (n-1)! = codim\,\ker(\delta_1^*)\,,$$

where U is the space (C.21).

Thus, it follows that $\ker(\delta_1^*) = X$, and Claim C.11 is proved.

□

Now let us pull together all strings and finish the proof of Theorem C.1. The theorem states that the map

$$\mathsf{Ger} \to H(\mathsf{Br})$$

defined on generators as stated in the theorem is a map of operads and an isomorphism. Showing that it is a map of operads amounts to checking that the Gerstenhaber relations hold between the images of the generators. This is a relatively trivial verification. Let us turn to the statement that $\mathsf{Ger} \to H(\mathsf{Br})$ is an isomorphism, i.e., that $\mathsf{Ger}(j) \to H(\mathsf{Br}(j))$ is an isomorphism for all j. Assume inductively the statement holds for $j < n$ and let us show it for $j = n$. By Claim C.11 and the arguments above it we know that

$$H(\mathsf{Br}(n)) \cong \Lambda\mathsf{Lie}(n) \oplus \mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n) \cong \mathsf{Ger}(n),$$

but we still have to verify that the morphisms $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n) \to H(\mathsf{Br}(n))$ and $\Lambda\mathsf{Lie}(n) \to H(\mathsf{Br}(n))$ constructed in the proof above agree with the ones proposed in Theorem C.1. Let us begin with the morphism $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n) \to H(\mathsf{Br}(n))$. Above, we constructed it recursively by sending an element $A \wedge B$ to the (class represented by the) insertion of the images of A and B in into the element $T_\wedge$ given in the Theorem. Hence, under the induction hypotheses our two maps $\mathsf{Ger}(n)/\Lambda\mathsf{Lie}(n) \to H(\mathsf{Br}(n))$ agree. Let us consider the two maps $\Lambda\mathsf{Lie}(n) \to H(\mathsf{Br}(n))$. In the proof of Claim C.11 above we identified $\Lambda\mathsf{Lie}(n)$ as the δ-closed linear combinations of trees without neutral vertices. Moreover the latter space was identified with $\Lambda\mathsf{Lie}(n)$ by (trivially) identifying the dual space (string-like trees modulo shuffles) with $\Lambda\mathsf{Lie}(n)^*$, i.e., with linear combinations of permutations modulo shuffles. Hence we are done if we can verify that the two pairings

$$\Lambda\mathsf{Lie}(n)^* \otimes \Lambda\mathsf{Lie}(n) \to \mathbb{K}$$

that one can construct from the data given agree. One verifies this (for example) by choosing an explicit basis of $\Lambda\mathsf{Lie}(n)$ given by Lie words of the form $\{?, \{?, \cdots, \{?, a_1\} \cdots\}$ and tracing the maps.

Theorem C.1 is proven. □



Fig. D.1. The *first* neutral vertex of a brace tree.



Fig. D.2. The filtration on V_p we use comes from the number of valence zero gray vertices attached to to the very left of the first neutral vertex. In this example, that number is three.

Appendix D. Br is a homotopy Tw fixed point – combinatorial argument

We showed in Section 9.3 above that the operad Br is a homotopy fixed point of Tw , i. e., that $\mathrm{Tw}\,\mathsf{Br} \to \mathsf{Br}$ is a quasi-isomorphism. One may give an alternative proof of that fact by an elementary combinatorial argument, which we briefly sketch in this appendix.

Elements of $\mathrm{Tw}\,\mathsf{Br}$ can be seen as series of brace trees, for which some of the labeled vertices have been colored in, say, gray. The differential is schematically depicted in Figure D.4. The number of neutral ("black") vertices of brace trees yields a descending complete filtration

$$\mathrm{Tw}\,\mathsf{Br} = \mathcal{F}^0 \supset \mathcal{F}^1 \supset \mathcal{F}^2 \supset \cdots$$

where $\mathcal{F}^p$ is composed of series in trees with $\geq p$ neutral (black) vertices. Let us consider the associated graded $\mathrm{Gr}\,\mathcal{F}$. Its differential misses those terms of Figure D.4 that produce a black vertex. We claim that $V_p := \mathcal{F}^p/\mathcal{F}^{p+1}$ is acyclic for $p \geq 1$. To show this we need to use additional notation. For a brace tree, the *first* neutral vertex is the one hit first when going around he tree in clockwise order, see Figure D.1. We filter V_p by the number of valence zero gray vertices attached to the very left of the first internal vertex, see Figure D.2. Taking a spectral sequence, the first differential increases that number by one, see Figure D.3. It is easy to see that the cohomology under this differential is zero. It follows that V_p is acyclic as claimed. Hence the projection $\mathrm{Gr}\,\mathcal{F} \to \mathcal{F}^0/\mathcal{F}^1$ is a quasi-isomorphism. From this one can see that also the projection $\mathrm{Tw}\,\mathsf{Br} \to \mathrm{Tw}\,\mathsf{BT}$ is a quasi-isomorphism. But since $\mathsf{Br} \to \mathrm{Tw}\,\mathsf{BT}$ is a quasi-isomorphism by section 9.2, we are done.



FIG. D.3. The differential increases the number of valence zero gray vertices attached to to the very left of the first neutral vertex by at most one. The component that increases this number by exactly one is shown here. Note that if this number is odd, the differential acts as zero.



FIG. D.4. The differential on $\mathfrak{g}_1$ has two parts. The first (first row) is obtained by splitting off neutral vertices from neutral, gray, or labeled vertices and comes from the differential on Br. The second part (second row) comes from the bracket with ϕ in the deformation complex. The very last term contains the terms depicted in Figure D.3.

APPENDIX E. A LEMMA ON A FILTERED COMPLEX

Lemma E.1. *Let C be a cochain complex with a complete descending filtration which is bounded from the left. I.e.,*

$$C = \mathcal{F}_{-N} \supset \mathcal{F}_{-N+1} \supset \cdots \supset \mathcal{F}_0 \supset \mathcal{F}_1 \supset \cdots$$

and $C = \lim_p C/\mathcal{F}_p$. If the associated graded complex $\operatorname{Gr} C$ is acyclic then so is C.

Proof. Suppose $c \in C$ is a cocycle. Then there exists j such that $c \in \mathcal{F}_j$. Then by acyclicity of the associated graded complex, there is a $b_j \in \mathcal{F}_j$ such that $c - db_j \in \mathcal{F}_{j+1}$. Proceed in this manner to define b_{j+1}, b_{j+2} etc. Then set $b := \sum_j b_j$. The sum converges since C is complete with respect to the filtration. Again by completeness

$$c - db = \lim_{N\to\infty} c - \sum_{j\le N} db_j = 0.$$

Hence c is exact. □

REFERENCES

[1] M. Batanin and M. Markl, Crossed interval groups and operations on the Hochschild cohomology, arXiv:0803.2249.

[2] C. Berger and B. Fresse, Combinatorial operad actions on cochains, Math. Proc. Cambridge Philos. Soc., **137**, 1 (2004) 135–174.

[3] C. Berger and I. Moerdijk, Axiomatic homotopy theory for operads, Comment. Math. Helv. **78**, 4 (2003) 805–831; arXiv:math/0206094.

[4] J. Chuang and A. Lazarev, L-infinity maps and twistings, Homology Homotopy Appl. **13**, 2 (2011) 175–195.

[5] J. Chuang and A. Lazarev, Combinatorics and formal geometry of the master equation, arXiv:1205.5970.

[6] V.A. Dolgushev, Covariant and equivariant formality theorems, Adv. Math. **191**, 1 (2005)147–177.

[7] V.A. Dolgushev, A proof of Tsygan's formality conjecture for an arbitrary smooth manifold, Ph.D. thesis, M.I.T. (2005); arXiv:math/0504420.

[8] V.A. Dolgushev and C.L. Rogers, Notes on Algebraic Operads, Graph Complexes, and Willwacher's Construction, arXiv:1202.2937.

[9] V.A. Dolgushev, D.E. Tamarkin, and B.L. Tsygan, Proof of Swiss Cheese Version of Deligne's Conjecture, Int. Math. Res. Notices **2011** (2011) 4666–4746.

[10] V.A. Dolgushev and T.H. Willwacher, A Homotopy Theory for Colored dg Operads, in preparation.

[11] B. Fresse, Koszul duality of operads and homology of partition posets, in "Homotopy theory and its applications (Evanston, 2002)", Contemp. Math. **346** (2004) 115–215.

[12] I. Galvez-Carrillo, A. Tonks, and B. Vallette, Homotopy Batalin-Vilkovisky algebras, arXiv:0907.2246.

[13] M. Gerstenhaber, The cohomology structure of an associative ring, Annals of Math., **78** (1963) 267–288.

[14] E. Getzler and J.D.S. Jones, Operads, homotopy algebra and iterated integrals for double loop spaces, hep-th/9403055.

[15] V. Ginzburg and M. Kapranov, Koszul duality for operads, Duke Math. J. **76**, 1 (1994) 203–272.

[16] V. Hinich, Tamarkin's proof of Kontsevich formality theorem, Forum Math. **15**, 4 (2003) 591–614; math.QA/0003052.

[17] M. Kontsevich, Operads and motives in deformation quantization, Lett. Math. Phys. **48**, 1 (1999) 35–72.

[18] M. Kontsevich and Y. Soibelman, Deformations of algebras over operads and the Deligne conjecture, *Proceedings of the Moshé Flato Conference Math. Phys. Stud.* **21**, 255–307, Kluwer Acad. Publ., Dordrecht, 2000.

[19] J.-L. Loday and B. Vallette, Algebraic Operads, *Grundlehren der mathematischen Wissenschaften*, **346**, Springer-Verlag 2012.

[20] M. Markl, Operads and PROPs, *Handbook of algebra.* Vol. 5, 87–140, Elsevier/North-Holland, Amsterdam, 2008; arXiv:math/0601129.

[21] M. Markl, Distributive laws and Koszulness, Ann. Inst. Fourier (Grenoble) **46**, 2 (1996) 307–323.

[22] J. E. McClure and J. H. Smith, A solution of Deligne's Hochschild cohomology conjecture, Contemp. Math. **293** (2002) 153–193, Amer. Math. Soc., Providence, RI; math.QA/9910126.

[23] S. Merkulov and B. Vallette, Deformation theory of representations of prop(erad)s, J. Reine Angew. Math. **634** (2009), 51–106. and J. Reine Angew. Math. **636** (2009), 123–174, arXiv:0707.0889.

[24] D. Quillen, Rational homotopy theory, Ann. of Math. (2) **90** (1969) 205–295.

[25] D. Tamarkin, Formality of chain operad of little discs, Lett. Math. Phys. **66**, 1-2 (2003) 65–72.

[26] D. Tamarkin, What do DG categories form? Compos. Math. **143**, 5 (2007) 1335–1358; math.CT/0606553.

[27] A.A. Voronov, Homotopy Gerstenhaber algebras, *Proceedings of the Moshé Flato Conference* Math. Phys. Stud., 22, 307-331. Kluwer Acad. Publ., Dordrecht, 2000.

[28] T. Willwacher, M. Kontsevich's graph complex and the Grothendieck-Teichmüller Lie algebra, arXiv:1009.1654.

[29] A. Yekutieli, Continuous and Twisted L-infinity Morphisms, J. Pure Appl. Algebra **207**, (2006), 575–606.

Department of Mathematics, Temple University,
Wachman Hall Rm. 638
1805 N. Broad St.,
Philadelphia, PA, 19122 USA
E-mail address: **vald@temple.edu**

Harvard University,
Department of Mathematics,
One Oxford Street
Cambridge, MA, 02138, USA
E-mail address: **wilthoma@math.harvard.edu**